ON THE VARIATION OF THE FACTOR h IN THE EQUATION

$$\frac{1}{2}mv^2 = h\nu.$$

BY FERNANDO SANFORD.

SYNOPSIS.

Relation between the β - and γ -rays of Ra B; Theory.—The author has previously shown that by combining the equations for orbital motion of an electron about a central positive charge with the equation $\frac{1}{2}mv^2 = h\nu$, the magnitude of the central positive charge about which a radiating electron is revolving may be calculated, and that when this value of the central charge is equated with one derived from an equation of the form of Moseley's equation for the frequency of X-rays the wave-length of the shortest lines in the K-radiation of elements of atomic numbers from 35 to 58 may be calculated as accurately as they can be measured. Later measurements of wave-lengths of elements with higher atomic numbers show that the computed wave-lengths vary progressively from the measured wave-lengths as the atomic numbers increase. It is also known that Moseley's law for the frequency of X-rays varies in the same manner.

The equation from which the author calculates the wave-length is

$$\frac{2^{1/2}h^{3/2}c^{1/2}}{m^{1/2}\lambda^{1/2}en} = 2e(N - 3.6).$$

Since the right-hand member of this equation shows that the nuclear charge in the case of K-radiation increases by $2e$ in passing from one element to the one of next higher atomic number, it is not probable that this law breaks down at atomic number 58, and that the nuclear charges of elements of higher atomic number increase by variable quantities. The failure of the equation to apply to elements of higher atomic number must then be looked for in the left-hand member. In this member the only variable quantities (since λ is measured) are m , the mass of the electron, and possibly h , the Einstein factor. These quantities appear in the form of the ratio h^3/m , and the author has calculated values of h^3/m which substituted in his equation will give the correct wave-lengths for elements of higher atomic number. These values of h^3/m are then plotted against their corresponding wave-lengths and a smooth curve is obtained. Values of h^3/m are then taken from this curve for a number of the γ -ray wave-lengths measured by Rutherford and Andrade. These wave-lengths are then assigned to certain groups of β -rays from Radium B on grounds which seem to the author permissible. The mass, m , and the quantity h corresponding to the speeds of these β particles were then computed. The values of h computed in this way seem to be derivable from m by the equation $h = 4.25 \log m + 2.50 \cdot 10^{-27}$. Using this value of h , wave-lengths were computed for the fourteen groups of β -rays from Radium B whose speeds were measured by Rutherford, and nine of them were found to agree satisfactorily with wave-lengths in the γ -radiation of Radium B and Radium C, while two more of the measured wave-lengths seem to correspond with the speeds of β -particles from Radium C.

Calculation of Orbital Velocities of Radiating Electrons by Substituting $4.25 \log m + 2.50 \cdot 10^{-27}$ for h in the equation $\frac{1}{2}mv^2 = h\nu$.

Using a series of values of v from $0.1 \cdot 10^{10}$ to $2.2 \cdot 10^{10}$ corresponding values of

m were calculated from Bucherer's equation for the transverse mass of an electron, viz., $m = m_0/\sqrt{1 - v^2/c^2}$. From these values ν was computed using the equation $\nu = \frac{1}{2}mv^2/(4.25 \log m + 2.50)$, and these values of ν were plotted on a large scale as abscissas using the values of v from which they were derived as ordinates. From this curve were taken values of v corresponding to all the highest frequencies of K-radiation as measured by Siegbahn and his colleagues and Duane and his colleagues. The values of v obtained in this way are then compared with values calculated from the equation $v = 2.054(N - 1.66)$ and are found to agree satisfactorily with no tendency to vary progressively with the higher numbers, as do the values calculated by Duane using h as a constant.

It is concluded that since h may be replaced in the equation $\frac{1}{2}mv^2 = h\nu$ by a function of m with a great improvement in the accuracy of the equation, there is no longer ground for speaking of an energy quantum in this equation.

THE writer has shown elsewhere¹ that by combining the equation for orbital motion of an electron about a central positive charge, viz., $mv^2/R = Qe/R^2$, with the equation $\frac{1}{2}mv^2 = h\nu$ it is possible to compute the central positive charge, Q , of a radiating atom, and from this value of Q and a corresponding value computed by an equation of the form given by Moseley to determine the wave-lengths of all known characteristic X-ray bands. Also, by combining the equation for Q with the formulæ for computing the wave-length of the lines in the various spectral series it was shown that the unit electrical charge may be computed from any line in any known spectral series of hydrogen or helium in terms of the wave-length and the serial number of the line. It is also possible by the use of the same equation to calculate the value of the Einstein factor h from the convergence charge of the Balmer series in hydrogen.

The above mentioned computations are not based upon any theory of atomic structure except that radiating electrons are in orbital revolution about a central positive charge, and that the frequency of revolution of the electron determines the wave-frequency of its radiation. In the opinion of the writer these assumptions are not incompatible with any of the commoner theories of atomic structure, but are admittedly in conflict with the Bohr theory of radiation.

The expression for the value of the central charge, Q , derived from the above equations is

$$Q = \frac{2^{1/2}h^{3/2}c^{1/2}}{m^{1/2}\lambda^{1/2}e\pi}.$$

By equating this value of Q with a value derived from an equation of the form of the Moseley equation we may get an expression for the

¹ See especially On The Nuclear Charges of Atoms, PHYSICAL REVIEW, IX., 383 (1917); The Astronomical Atom and the Spectral Series of Hydrogen, Astrophysical Journal, XLVIII., 1 (1910); The Helium Spectrum and the Unit Electrical Charge, Astrophysical Journal, XLIX., 337 (1919).

value of λ . Thus for the characteristic K-radiation $Q = 2e(N - 3.6)$ where N is the atomic number. Then for this radiation

$$\frac{2^{1/2} h^{3/2} c^{1/2}}{m^{1/2} \lambda^{1/2} e \pi} = 2e(N - 3.6).$$

This equation has been found to apply to the elements from $N = 35$ to $N = 58$ so that the wave-lengths of the K-radiation from these elements may be calculated from this formula as accurately as they may be measured.¹

In the PHYSICAL REVIEW of January, 1920, the present writer undertook to show that if the speed of the β -particles sent off from Ra-B and Ra-C is related by Einstein's law to the accompanying γ -radiation then before the expression $\frac{1}{2}mv^2$ may be substituted for Ve in that equation the factor h , as well as m , must be assumed to vary with the speed of the β -particles.

Since the article referred to was written, two papers which have an important bearing on this question have appeared. In the *Physikalische Zeitschrift* of June 1, 1919, Siegbahn and Jönssen have published new determinations of the wave-lengths of highest frequency in the K-radiation series of a considerable number of elements from cadmium to uranium, and in the *Physical Review* of December, 1919, Duane and Hang-Fuh-Huh and Duane and Shimizu have extended the wave-length measurements on the absorption frequencies of the K-radiation to elements of both lower and higher atomic numbers than those whose frequencies were determined by Blake and Duane. These measurements when taken with the wave-length determinations of Siegbahn and Stenström in 1916 give a series of measurements of the short wave-lengths in K-radiation for most of the elements from atomic numbers $N = 12$ to $N = 92$.

When these newly determined wave-lengths are computed by our formula it is found that the correspondence is equally as good as for the measurements of Blake and Duane up to an atomic number 58, but that from this point onward the calculated values are greater than the measured values. Siegbahn and Jönssen call attention to the fact that from this point onward their wave-lengths do not follow Moseley's law, and they attribute the variation from this law to the variable mass of the electron at high speeds.

In the papers in the PHYSICAL REVIEW, Duane and his colleagues calculate a value of v for each of the elements whose wave-lengths they have measured by using the equation $\frac{1}{2}mv^2 = h\nu$ and computing the value of m at each speed by the Bucherer equation, $m = m_0/\sqrt{1 - (v^2/c^2)}$.

¹ See PHYSICAL REVIEW, XV., 68 (January, 1920).

They found that their values of v computed in this way bore a straight line relation to the atomic numbers, as does the square root of the frequencies in Moseley's equation, but that for atomic numbers above 58 their calculated values of v vary slightly, but progressively, from the values given by the atomic number relation.

It will be seen that in the equation

$$\frac{2^{1/2}h^{3/2}c^{1/2}}{m^{1/2}\lambda^{1/2}e\pi} = 2e(N - 3.6)$$

a variation in m can affect only the left-hand member of the equation, and that the only other quantity in this member which may possibly vary (since λ is measured) is the quantity h , or, in other words, the possible variable quantity in this member is the fraction h^3/m . Since the right-hand member of the equation merely shows that Q increases by $2e$ in passing from any element to the one of next higher atomic number, there is no reason for thinking that this law changes and that the increase of charge becomes variable for elements of higher atomic number than 58.

If we may rely upon the constancy of the right-hand member of the equation, we may compute values of h^3/m for each of the given wave-lengths and perhaps decide whether the variation in this side of the equation is due wholly to m , or whether h is also probably variable.

In Table I. Blake and Duane's wave-length measurements are given in the third column, those of Siegbahn and Jönsson in the fourth column, the wave-lengths calculated from our formula in the fifth column and the values of h^3/m required to give the measured wave-lengths in the sixth column. It may be seen that h^3/m remains constant to within the errors of measurement to about $N = 58$, from which point on it decreases, while the calculated values of the wave-length also show a systematic variation from about the same value of N .

When these values of h^3/m are plotted as ordinates against their corresponding wave-lengths as abscissas a smooth curve is obtained from which values of h^3/m may be taken for any wave-length down to $\lambda = 0.1 \cdot 10^{-8}$. If corresponding values of v could now be found it would be possible to calculate the respective values of h . This I have attempted to do in the following way:

All of the wave-length calculations and the values of h^3/m given in Table I. are based upon the assumption that the speeds of electrons expelled under the stimulus of ultra-violet light or of X-rays are the orbital speeds which they possessed at the instant of breaking away from their respective atoms. If this be the true explanation of their velocities, it seems almost certain that the same explanation must account for the speeds of the β -particles expelled by radioactive bodies. This would

TABLE I.

Element.	<i>N</i> .	λ_{10^8} , B. & D.	λ_{10^8} , S. & J.	λ_{10^8} , Calc.	h^2/m , 10^{24} .
Br.....	35	.9179		.924	309.5
Rb.....	37	.8143		.817	310.5
Sr.....	38	.7696		.771	311
Yt.....	39	.7255		.727	309.5
Zr.....	40	.6872		.689	311
Nb.....	41	.6543		.654	311
Mb.....	42	.6180		.618	311
Ru.....	44	.5584		.558	311
Rh.....	45	.5324		.532	311.5
Pd.....	46	.5075		.507	312.5
Ag.....	47	.4850		.484	312
Cd.....	48	.4632	.4629	.463	312
In.....	49	.4434		.443	312.5
Sn.....	50	.4242	.4231	.424	312
Sb.....	51	.4065		.409	312
Te.....	52	.3896	.3877	.389	310.5
I.....	53	.3737	.3715	.373	310.5
Cs.....	55	.3444	.3436	.345	310.5
Ba.....	56	.3307	.3306	.332	310.5
La.....	57	.3188	.3186	.320	310.5
Ce.....	58	.3073	.3064	.308	310.5
Pr.....	59		.2946	.297	309
Nd.....	60		.2835	.286	308
Sa.....	62		.2636	.2665	307.5
Eu.....	63		.2543	.2585	306.5
Gd.....	64		.2456	.2495	306.5
Dy.....	66		.2294	.234 +	305.5
Ho.....	67		.2214	.2265	304.5
Pt.....	78		.1578	.165	298.5
Au.....	79		.1524	.1605	296.5
Hg.....	80		.1479	.156	295
Tl.....	81		.1427	.152	292
Pb.....	82		.1385	.148	291.5
Bi.....	83		.1346	.1445	290
Th.....	90		.1127	.122	288
U.....	92		.1048	.1167	280

require that the Einstein photoelectric equation should hold for the energies of the β -rays and the frequencies of the accompanying γ -radiations from the same elements. Rutherford and his colleagues have given much time to the investigation of this question, and are of the opinion that such a relation does, indeed, hold and that some of the γ -rays from Ra-B and Ra-C are the true K- and L-radiation bands of these elements.

In an article on The Connexion Between β and γ Ray Spectra¹ Ruther-

¹ Phil. Mag., 28, 305-319.

ford gives the velocities relative to the velocity of light of fourteen groups of β -rays expelled from Ra-B and ten groups of β -rays from Ra-C. In an article by Rutherford and Andrade on The Spectrum of the Penetrating γ Rays from Radium B and Radium C in the same volume of the Philosophical Magazine is given a table of the wave-lengths of fourteen different lines in the γ -ray spectrum of these elements. The authors do not know to which element the various lines belong, but give their reasons for believing that the shortest wave-lengths are due to Radium C, and that the lines $\lambda = 1.59 \cdot 10^{-9}$ and $\lambda = 1.69 \cdot 10^{-9}$ are the characteristic K-radiation lines of Radium B.

In the data on the velocities of β -rays the groups of rays of speeds 1.905–2.100 are marked very strong, indicating that there are more β -rays of this speed than of any other unless it be 1.01. Since the lines of wave-length 1.59 and 1.69 are referred to as the strongest lines in the γ -ray spectrum, it seems reasonable to assign these lines to the group of β -rays whose speeds are from 1.9 to $2.10 \cdot 10^{-10}$.

If we calculate a value of m for a speed of $1.9 \cdot 10^{-10}$ we find $m = 1.16 \cdot 10^{-27}$. Since the wave-length 1.69 lies within the spectral region for which we have computed the value of h^3/m , we may take the value of this ratio from the curve, which was drawn for that region, where we find it to be $300 \cdot 10^{-54}$. Accordingly, $h^3 = 348 \cdot 10^{-81}$ and $h = 7.03 \cdot 10^{-27}$.

In order to express graphically the relation of h to m we must have this relation at one more point, at least, since we already have the initial values of the two quantities. By a slight extrapolation of our curve it was seen that the γ -ray line $\lambda = .99$ must give $h^3/m = 276$, approximately. For trial purposes this line was assigned to the highest speed β -particle from Radium-B, viz., $v = 2.469$. For this speed $m = 1.2 \cdot 10^{-27}$, and if $h^3/m = 276$, $h = 7.60$.

When these three values of h and m were compared they seemed to suggest that h varies as $\log m$. On plotting a curve with the values of h as abscissas and the values of $\log m$ as ordinates, a linear relation was seen to hold between the two quantities, so that for $h \cdot 10^{27}$ and $\log m \cdot 10^{28}$, $h = 4.25 \log m + 2.50 \cdot 10^{-27}$.

Assuming the validity of this equation it is possible to calculate m and h for each of the speeds given, and to compare the values of h^3/m obtained in this way with values taken from our curve for each of the wave-lengths measured in the spectra of Radium B and Radium C. This is done in Table II., where the velocities of the β -rays from Radium B are given in the first column, the corresponding values of m as computed from the equation $m = m_0 / \sqrt{1 - (v^2/c^2)}$ in the second column, h as derived from

the equation $h = 4.25 \log m + 2.50$ in the third column and h^3/m in the fourth column. In the fifth column are given values of λ corresponding to the values of h^3/m in the fourth column as taken from the curve drawn to show this relation. In the sixth column are given the values of λ as measured by Rutherford and Andrade in the γ -radiation of Radium B and Radium C.

TABLE II.

β Radiation from Radium B.				γ Radiation.	
v .	m .	h .	h^3/m .	λ Calc.	λ Meas.
2.469	1.59	7.60	276	.99	.71
2.450	1.56	7.55	278	1.03	.99
2.391	1.49	7.46	281	1.10	—
2.361	1.46	7.45	282	1.15	1.15
2.286	1.39	7.35	287	1.24	—
2.252	1.36	7.31	288	1.27	—
2.193	1.32	7.25	289.5	1.31	—
2.157	1.295	7.22	292	1.37	1.37
2.100	1.26	7.16	292	1.37	1.37
1.968	1.195	7.06	297	1.55	1.59
1.905	1.165	7.03	300	1.69	1.69
—	—	—	—	—	1.96
—	—	—	—	—	2.29
—	—	—	—	—	2.42
1.278	.995	6.74	307	2.62	2.62
1.242	.990	6.73	308	2.90	2.96
—	—	—	—	—	3.24
—	—	—	—	—	3.93
1.011	.956	6.66	309	4.28	4.28

A comparison of columns five and six of Table II. will show that nine of the fourteen measured wave-lengths correspond as closely as could be expected with the values calculated from the speeds of the β -particles on the assumption that $h = 4.25 \log m + 2.50$. For the three slowest groups of β -particles the values of h^3/m fall on a part of the curve where h^3/m varies but slightly with the wave-length, and where the values of λ derived from this curve must be more uncertain than for the faster β -particles.

The slowest speed given has been assigned to the wave-length 4.28 from the following considerations: The radiation of particles at this speed is marked "very strong," indicating that this group of β -particles corresponds to a characteristic radiation of the element. I have shown elsewhere¹ that the shortest wave-length of the characteristic L-radiation

¹ Electrical Charges of Atoms and Ions, Leland Stanford Junior University Publications, p. 70.

may be calculated from the equation

$$\lambda = \left[\frac{2.882 \cdot 10^{-12}}{e(N - 15.4)} \right]^2$$

for elements up to atomic number 71. This equation would give for Radium B ($N = 82$) a wave-length $\lambda = 4.33$, which is within the error of measurement the line 4.28.

The measured speeds of the β -particles from Radium C are with two exceptions much higher than those from Radium B, and the curve derived from X-ray wave-lengths may not safely be extrapolated far enough to give a probable estimate of their γ -ray wave-length. The slowest group of these particles has a velocity of $2.25 \cdot 10^{10}$, which would make its γ -ray wave-length about 1.27 from our computation. The next slowest group of β -particles has a speed of 2.604. This would give $m = 1.80$ and $h = 7.83$, making $h^3/m = 266$. This seems to correspond with the measured wave-length .71.

It seems not improbable that the γ -radiation from the other β -particles of Radium C was of too short wave-length to be measured by Rutherford and Andrade. With their spectrometer the wave-length taken as .71 was determined from a deviation of only $43'$, and the authors say, "It is surprising that the architecture of the crystal (rock salt) is sufficiently definite to resolve such short wave-lengths. This is especially the case when we consider that owing to the heat agitation of the atoms, the distance between the atoms must be continually varying over a range comparable with the wave-length of the radiation."

In a paper by F. Dessauer and E. Back published in *Berichte der Deutschen Physikalischen Gesellschaft*, May 30, 1919, the authors give the results of their attempt to find the shortest wave-length given by X-rays from any element. By using a potential of 170 k.v. and an exposure of 9 hours they found the apparent limit of X-rays at $\lambda = .57 \cdot 10^{-9}$. It is probable that all of the β -rays from Radium C except the two sets referred to would be accompanied by γ -rays of shorter wave-length than $.57 \cdot 10^{-9}$.

If these estimates of the relation between the speeds of β -rays and the wave-lengths of their accompanying γ -radiations are correct, it seems almost certain that the speeds of β -particles are the orbital speeds which they possessed at the instant of escaping from the atoms.

It also seems certain that for speeds as great as those of β -particles the product $\frac{1}{2}mv^2$ of the β -rays does not bear a constant ratio to the frequency of the accompanying γ -radiation.

It has already been mentioned that Duane and his colleagues have

calculated values of ν from the K-radiation of a large number of elements which values are related to the atomic numbers by an equation of the form of Moseley's equation for the square root of the frequencies. It will be seen from our fundamental equation for the atomic charge that this will necessarily be true if h^3/m is constant or bears a straight line relation to the atomic numbers. It was observed, however, that this law of Duane's holds only so far as the wave-length may be correctly calculated on the assumption of the constancy of h^3/m .

In order to test the same relation with values of ν calculated from our hypothesis, I have calculated values of m for a series of values of ν from $\nu = 0.1 \cdot 10^{10}$ to $\nu = 2.2 \cdot 10^{10}$, letting

$$m = \frac{9 \cdot 10^{-28}}{\sqrt{1 - \nu^2/c^2}}.$$

Then from the equation $\frac{1}{2}m\nu^2 = \nu(4.25 \log m + 2.50)$ I have calculated values of ν for each value of ν used in computing m . These values of ν were then plotted on a large scale as abscissas against the respective values of ν as ordinates. From this curve it is possible to obtain a value of ν accurate to the third significant figure for any value of ν from $\nu = 0.1 \cdot 10^{18}$ to $\nu = 40 \cdot 10^{18}$.

Values of ν corresponding to all the K-radiation frequencies measured by Siegbahn and Stenström, Siegbahn and Jönssen, Blake and Duane, Duane and Hang-Fuh-Huh and Duane and Shimizu were then taken from this curve and plotted against their respective atomic numbers. In the case of the Siegbahn and Stenström measurements the frequencies for the $K\beta$ lines were used throughout. It was found that the velocities calculated in this manner bear a straight line relation to the atomic numbers with no tendency to systematic departure on either side of the line for higher atomic numbers. This relation of velocity to atomic number may be expressed by the equation $\nu = 2.054(N - 1.66)$.

The relation between the values of ν calculated in the two ways is shown in Table III., in which the wave-length measurements of Siegbahn and his colleagues are given in the third column and those of Duane and his colleagues in the fourth column. The corresponding values of ν are given in the fifth column. Where two values of λ are available a mean value is used in computing ν . The sixth column gives ν from the equation

$$\nu = \sqrt{\frac{2\nu(4.25 \log m + 2.50)}{m}}$$

and the seventh column gives ν from the equation $\nu = 2.054(N - 1.66)$.

It will be seen that though the series runs to $N = 92$, the agreement

TABLE III.

$$v_1 = \sqrt{\frac{2\nu(4.25 \log m + 2.50)}{m}},$$

$$v_2 = 2.054(N - 1.66).$$

Element.	<i>N</i> .	λ Siegbahn and Others.	λ Duane and Others.	ν .	v_1 .	v_2 .
Mg.....	12	9.477		.3165	.213	.2124
Al.....	13	7.986		.3753	.233	.233
Si.....	14	6.759		.4436	.2535	.2535
P.....	15	5.808		.516	.274	.274
S.....	16	5.018		.5978	.295	.2946
Cl.....	17	4.394		.6826	.316	.3152
K.....	19	3.449		.8700	.357	.3564
Ca.....	20	3.086		.973	.377	.377
Sc.....	21	2.778		1.080	.3975	.3972
Ti.....	22	2.509		1.195	.418	.418
Va.....	23	2.281		1.315	.437	.438
Cr.....	24	2.079		1.443	.458	.458
Mn.....	25	1.902	1.8892	1.582	.480	.4795
Fe.....	26	1.743	1.7396	1.721	.500	.500
Co.....	27		1.6018	1.86	.520	.521
Ni.....	28		1.4890	2.015	.540	.541
Cu.....	29		1.3785	2.175	.560	.562
Zn.....	30		1.2963	2.315	.578	.582
Ga.....	31		1.1902	2.521	.602	.603
Ge.....	32		1.1146	2.693	.623	.623
As.....	33		1.0435	2.875	.644	.644
Se.....	34		.9790	3.065	.665	.664
Br.....	35		.9179	3.268	.685	.685
Rb.....	37		.8143	3.684	.726	.726
Sr.....	38		.7696	3.898	.745	.746
Yt.....	39		.7255	4.136	.767	.767
Zr.....	40		.6872	4.365	.788	.788
Nb.....	41		.6503	4.61	.808	.808
Mo.....	42		.6180	4.855	.830	.829
Ru.....	44		.5584	5.374	.871	.871
Rh.....	45		.5330	5.63	.891	.890
Pd.....	46		.5075	5.913	.912	.911
Ag.....	47		.4850	6.185	.9325	.932
Cd.....	48	.4629	.4632	6.48	.953	.952
In.....	49		.4434	6.766	.973	.973
Sn.....	50	.4231	.4242	7.08	.994	.994
Sb.....	51		.4605	7.384	1.014	1.014
Te.....	52	.3877	.3896	7.722	1.036	1.034
I.....	53	.3715	.3737	8.065	1.057	1.055
Cs.....	55	.3436	.3444	8.725	1.098	1.096
Ba.....	56	.3306	.3307	9.08	1.118	1.116
La.....	57	.3186	.3188	9.42	1.138	1.137
Ce.....	58	.3064	.3068	9.80	1.160	1.157
Pr.....	59	.2946		10.18	1.181	1.178

Element.	N .	λ Siegbahn and Others.	λ Duane and Others.	ν .	ν_1 .	ν_2 .
Nd.....	60	.2835	.2861	10.54	1.201	1.199
Sa.....	62	.2636		11.38	1.243	1.239
Eu.....	63	.2543		11.79	1.263	1.260
Gd.....	64	.2456		12.21	1.284	1.282
Tb.....	65		.2398	12.51	1.299	1.301
Dy.....	66	.2294	.2308	13.04	1.324	1.322
Ho.....	67	.2214		13.54	1.348	1.343
W.....	74		.1786	16.78	1.485	1.486
Os.....	76		.1683	17.82	1.531	1.528
Pt.....	78	.1578		19.01	1.572	1.568
Au.....	79	.1524	.1541	19.57	1.590	1.589
Hg.....	80	.1479	.1501	20.14	1.608	1.608
Tl.....	81	.1427		21.01	1.640	1.626
Pb.....	82	.1385	.1424	21.35	1.651	1.650
Bi.....	83	.1346		22.29	1.681	1.671
Th.....	90	.1127		26.61	1.814	1.815
U.....	92	.1048		28.60	1.866	1.835

between the two values of ν is very close and no progressive variation appears. In only one case, that of uranium, is the difference between the two values of ν as great as one per cent. From the equations here used the wave-length of the shortest K-radiation from uranium should be $\lambda = .1099 \cdot 10^{-8}$ instead of $.1048 \cdot 10^{-8}$, as given by Siegbahn and Jönssen. Dessauer and Back, in the paper already mentioned, attempted to find the wave-length of the $K\beta$ line of uranium, and concluded that it lies between $.104 \cdot 10^{-8}$ and $.154 \cdot 10^{-8}$. When extrapolated from their measured wave-length of the platinum line by Moseley's equation they found it to be between $.114$ and $.136 \cdot 10^{-8}$, but it is well known that Moseley's equation does not apply closely to elements of the higher atomic numbers. From Sommerfeld's formula this line would lie between $.131$ and $.146 \cdot 10^{-8}$.

The one other considerable deviation is in the case of thallium, for which our equation would give a wave-length $.1455 \cdot 10^{-8}$ instead of $.1427 \cdot 10^{-8}$, as determined by Siegbahn and Jönssen.

On the whole, the two values of ν are in very satisfactory agreement when we take into consideration that the most careful wave-length determinations in this region by different experimenters frequently differ in the third decimal place.

While it is probably too much to claim that the relations shown in Table II. and Table III. give an adequate test of the value of the expression here substituted for h in the equation $\frac{1}{2}mv^2 = h\nu$, they are, so far as is known to the writer, the only tests thus far made of the validity of

this equation. The equation $Ve = h\nu$ has frequently been tested by experiment, but the assumption that all the energy given to an electron in an electrostatic field appears in the form of kinetic energy of the electron has not been so tested. Whatever evidence may be derived from the relations here shown indicates that in this form h does not represent an energy quantum, but only a function of the mass of the electron.