and showed no edema except for the uvula. The structures of the under surface of the tongue were markedly swollen. On palpation with the little finger, it proved tender to the touch with the maximum intensity at a point opposite the second molar tooth. That point was rather harder and thicker than the rest. Temperature, 101.2. Pulse 90. Ether was administered, the patient became cyanotic, artificial respiration with traction upon the tongue had to be resorted to. The operation was performed in a sitting position by reflected light. The tongue was lifted and rolled over to the side, and an incision was made with a straight knife about one-half inch in length, in the most thickened spot, under and parallel to the side of the tongue. The opening was dilated with forceps and about two teaspoonfuls of foul pus were evacuated. A probe was put into the opening about two inches deep and did not show any pockets. A strip of gauze was put in the wound and left there. Cracked ice on tongue ordered.

Dr. Masten informed me that the patient made an uneventful recovery after the pus was evacuated, and the wound closed on the second day after the packing was removed. The infective agent, as suggested by Dr. Masten, must have been the staphylococcus on account of the absence of high fever and the large amount of pus found.

On my second visit, when the throat could be examined satisfactorily, the following interesting point was noticed: The uvula was still swollen and a red streak was running from it along the anterior pillar down to the tongue. The tonsils were not inflamed and the pillars not congested. Whether the primary infection started in the uvula, and from the glands around the uvula was transmitted by the way of the lymphatics along the anterior pillars to the sides of the tongue, or vice versa, the writer is not prepared to say, though it is more reasonable to assume the latter.

I wish to thank Dr. Charles H. Masten, of Worcester, for the information given and for the valuable assistance rendered at the operation.

REFERENCES.

- Butlin, H.: Diseases of the Tongue.
Keen's Surgery.
Deaver: Surgical Anatomy.
Gray's Anatomy.
Journal A. M. A., Vol. Ixv.

AN OPERATION FOR THE CORRECTION OF THE DEFORMITY DUE TO "OBSTETRICAL PARALYSIS."

By MARK H. ROGERS, M.D., BOSTON.

THIS case is reported to show the result of an operation which seems to have a very definite place in the correction of the deformity that results from the so-called "birth palsy." Lately there has been considerable discussion as to the exact lesion in these cases, whether the results are due to a tearing of the brachial plexus or whether these may be a congenital dislocation of the shoulder joint. Whatever is the underlying

Before operation.

cause and what treatment in the early stages should be employed are not considered in this report. The deformity that results is practically always the same, with the amount of paralysis of individual muscles varying in each case.

The usual deformity that follows brachial paralysis occurring at child-birth is essentially an inward rotation of the whole arm from the shoulder-joint, so that, when the hand lies normally against the side of the body, the elbow is carried away from the body. This is well shown in the first illustration, which was taken before the operation. The elbow-joint is rotated so that the anterior surface points toward the body instead of directly forward.

Without taking into consideration the extent of the paralysis, and we know that it varies to a great extent, the resulting deformity is practically always as described. As the child grows older the deformity becomes more conspicuous, and it is the carrying angle of the elbow that is most prominent. Another result of the deformity is seen when the patient attempts to raise the hand to the mouth. With the direction of the elbow-joint reversed, it is always necessary to raise the arm away from the body to bring the hand up. Also when the patient attempts to use the hand as in picking up any object or in the ordinary handshake, it is necessary first to rotate the whole arm and also to bring the elbow away from the body.

The operation here shown for the correction of the rotation has been in vogue in Germany,

Before operation.

and has been described by Vulpius and Lange. The result, as shown by the photographs of the case, is given because the operation is evidently not employed here very often, and it does seem to accomplish a very definite object.

The operation is simply an osteotomy of the upper portion of the humerus about two inches below the shoulder joint, followed by a one-quarter rotation of the whole arm below the line of fracture. An incision is made between the muscle planes sufficiently long to be sure that there is no nerve involvement. A clean cut osteotomy is done, and before the incision is closed the arm is brought out at a right angle to the body and the lower fragment is rotated outward, while traction is kept on the hand. The tip of the finger is held on the fracture to see that there is no overriding and that the position is good. The wound is closed and the arm and body are encased in a shoulder spica with the arm full-length at right angles and the palm of the hand pointing upwards. It is then treated as an ordinary fracture.

The results of the operation are a normal carrying position of the arm, with the elbow at the side, and a normal position of the hand. It is also a striking feature that it is possible to bring the hand to the mouth so that it is possible to use the hand normally while eating.

The only possible danger seems to be the possibility of non-union from poor position of the fragments. This is very slight and may be obviated by the use of a bone-plate if necessary. Three cases form the basis of this report, the

After operation.

photographs of one being shown here, and they have all given good and equal results. It is not an operation to be done in the early steps of the paralysis, but only after the deformity has been developed. It evidently allows a more normal use of the hand and arm, and also corrects a very prominent deformity, which is very noticeable and objectionable.

Medical Progress.

PROGRESS IN PEDIATRICS.

LOBAR PNEUMONIA IN INFANCY AND CHILDHOOD. A SUMMARY OF THE LITERATURE OF THE PAST TEN YEARS.

By JOHN LOVETT MORSE, A.M., M.D., BOSTON.

(Concluded from page 135.)

TREATMENT.

Fresh and Cold Air. Northrup, in a paper read before the Pediatric Section of the New York Academy of Medicine, January 14, 1904, entitled "The Treatment of Bronchopneumonia in Children," called attention to the importance of the hygienic treatment of diseases of the respiratory tract in childhood. He advocated fresh air, a large room with southern exposure and the bed in the middle of the room.