A CASE OF SYRINGOMA.

By S. NORMAN PAUL,
Sydney, N.S.W.

With Remarks by H. G. ADAMSON.

THE patient, J. S—, is a man, aged 46 years. Situated over the left side of the sternum, 2 in. inside the left nipple and reaching to the mid-sternal line, there are about twenty-five smooth, firm, raised growths, the largest 2·5 cm. across and raised 1·5 cm. The growths are sessile, intimately associated with the skin, and move with it. The bases of the smaller nodules coalesce to form an infiltrated plaque (Fig. 1). The two largest growths have been present for twenty-five years, three others for twelve years, and a number have appeared in the past three years. In colour some show a decided dark blue, the remainder being either pink or pinkish-blue. Dilated vessels are coursing over the two larger growths, which appear cystic. The blue colour has been noticed only in the past twelve months. There are no enlarged glands. The patient has dyspepsia and states that, after an attack of dyspepsia, for several months past, a new growth has appeared and become painful, but otherwise there is no pain. The whole mass was excised and the blue coloration was found to be caused by blood, which filled the larger and some of the smaller cysts. The section shows the growth to be situated in the corium, lobulated, surrounded by fibrous tissue, and appears to be derived from the sweat-glands.

Diagnosis: (?) Adenoma of the sweat-glands.

REMARKS.

Dr. Paul's case at once excites interest on account of its unusual character (Fig. 1). It would be impossible to predict its histological structure from its naked-eye appearances, but microscopical sections were prepared by Dr. Paul, and from them (Figs. 2 and 3) he made the diagnosis of Syringoma.

The group of new growths to which Syringoma belongs comprises a number of types. It includes growths made up of embryonic basal cells in various stages of differentiation towards gland tissue.

FIG. 1.—Photograph of a case of Syringoma (?).

FIG. 2.—Syringoma (?). Drawing of a section of a small nodule. Under the low-power magnification the section recalls that of an adenoid epithelioma of the type of Brooke's "benign cystic epithelioma" (tricho-epithelioma, so-called), except that it is not divided into lobules.

FIG. 3.—Syringoma (?). Drawing of the same section under a high power, which shows the tubule-like arrangement of the cell-growth and suggests the appearance of an adenoma of the sweat-glands, *i.e.* syringadenoma or syringoma, rather than that of an adenoma derived from pilosebaceous structures, *i.e.* epithelioma of Brooke or tricho-epithelioma.

TO ILLUSTRATE DR. S. NORMAN PAUL AND DR. H. G. ADAMSON'S
ARTICLE ON A CASE OF SYRINGOMA.

FIG. 4.—Section from a nodule on the forehead in a typical case of Epi-
thelioma adenoides cysticum of Brooke. Contrasted with Figs. 2 and 3 it
shows a lobulated tricho-epitheliomatous structure rather than a tubuli-
ferous syringomatous character.

FIG. 5.—Section from a benign lower eyelid tumour for comparison with
Figs. 2 and 3. It shows again a basal-cell growth of the tricho-epithelioma
or embryonic pilosebaceous type. The resemblance to rodent ulcer is
striking.

The growths may have the structures of fully developed gland tissue, in which case we call the tumour an adenoma: as Adenoma sebaceum and sudoriparous adenoma or sweat-gland tumours. They may be growths composed of more embryonic, less differentiated cells, as in Brooke's Epithelioma adenoides cysticum and in Lymphangioma tuberosum multiplex of Kaposi (*i. e.* Syringoma). They may be growths made up of basal cells which have lost all power of differentiation into gland-tissue cells and which have retained only that of proliferation, as in certain basal-cell epitheliomata, which we call rodent ulcer.

It seems probable that all these growths are derived from embryonic "cell-rests" as conceived by Cohnheim. Whether the cell-rest develops into an adenoma, into an adenoid epithelioma or into rodent ulcer, may perhaps depend upon the stage of development of the elements when the cell-rests were isolated, or perhaps upon the age of the patient at the moment the cell-rests become aroused into activity. It is conceivable that a glandular "cell-rest" which begins to grow during early life is more likely to develop into gland tissue than one which begins to proliferate only late in life. This question of the possible relationship between benign glandular tumours and rodent ulcer is one of considerable interest.

Since the case with which we are now concerned shows growths with no tendency to invade and destroy normal tissues or to ulcerate, we must place it among the benign forms, and, in order to know in which class to place it, it will be well to recall the already recognised types of adenomata and of adenoid epitheliomata.

The acknowledged types of these growths of glandular origin are:

(1) *Adenoma sebaceum*: in which there occur on the cheeks and sides of the nose and on the forehead little rounded tumours made up of sebaceous-gland tissue.

(2) *Epithelioma adenoides cysticum* of Brooke: in which the tumours are multiple and situate mainly on the face and especially about the forehead, temples, eyes and nose, but also on the scalp, neck and chest; and which is sometimes a family disease.

The structure of these growths is that of embryonic pilosebaceous follicles, so that they have been described as tricho-epitheliomata. Histologically the growths closely resemble rodent ulcer (Fig. 4) and it seems probable that rodent ulcer is also a tricho-epithelioma or

embryonic pilosebaceous tumour, but made up of cells less stable than those of *Epithelioma adenoides* of Brooke and with, therefore, a greater tendency to break down and ulcerate.

(3) The multiple benign tumours of the scalp, the so-called *Endotheliomata of the scalp* of Spiegler, and Mulert, have been shown by Dubreuilh and Auché to be benign epitheliomata derived from the basal-cell layer of the epidermis and histologically to resemble rodent ulcer. In one of my cases of *Epithelioma adenoides cysticum* of Brooke, and in a case published by Freibos, these multiple scalp-tumours were associated with the face and body tumours. Freibos considers them to be syringomata, but I should agree with Dubreuilh in regarding them as pilosebaceous or tricho-epitheliomata.

(4) *Benign tumours of the lower orbital region* and of the naso-facial groove which have in many cases the structure of basal-cell epithelioma of the embryonic pilosebaceous or tricho-epithelioma type as in the illustration here given (Fig. 5). Jarisch's so-called "*Hæmangioendothelioma*" was probably of this nature. Other of these growths are probably of embryonic sweat-gland origin, as in Darier's "*Hidroadénomes des paupières inférieures*."

It is becoming now a widely accepted view that rodent ulcer (which occurs so frequently in this situation) is a growth of embryonic pilosebaceous gland origin, or in other words a malignant tricho-epithelioma. It is an interesting question whether rodent ulcer may be sometimes of embryonic sweat-gland origin, *i. e.* a malignant syringoma; but there is certainly less resemblance between the histological structure of a rodent ulcer and that of a syringoma than there is between that of rodent ulcer and benign tricho-epithelioma.

(5) An important type, probably of embryonic sweat gland origin, is that originally described by Kaposi as "*Lymphangioma tuberosum multiplex*," and later by Jacquet and Darier as "*Hydradénome eruptif*," by Török as "*Syringo-cystadenoma*," by Gassmann as "*Nævi cyst-epitheliomatosi disseminata*" and now generally known by the name given to it by Unna, of *Syringoma* or *Syringadenoma*.

Cases of this type have been frequently confused with Brooke's "*Epithelioma adenoides cysticum*," which, as we have seen, is now regarded as an embryonic pilosebaceous gland tumour. But they differ from Brooke's type in that the tumours are situated on the chest below the clavicle rather than on the face and scalp, in that

they do not occur as a family affection, and in that their histological structure has a more tubular arrangement which recalls the tubules of an embryonic sweat-gland.

(6) Single tumours in various situations have been described as *sweat-gland tumours*, *sudoriparous adenoma*, or *Syringoma*. Unna quotes several cases by older writers as being genuine examples of sweat-gland adenoma. Darier and Fordyce have collected and reported cases (Darier, a case of multiple tumours), and Wolfheim, Torok, Blaschko, Rolleston, and others have described benign or malignant tumours of sweat-gland origin.

It is not always clear whether these writers are dealing with tumours which show a frankly tubular arrangement so that they could be recognised as sweat adenoma, just as we recognise sebaceous adenoma, or whether they are tumours which merely suggest a sweat-gland origin as in the type of multiple tumour which Kaposi described as *Lymphangioma tuberosum multiplex*, and which we now generally call syringoma.

It is interesting to note that some of these observers have recorded cases of sweat-gland tumours becoming malignant, but whether the malignancy is of the type of rodent ulcer is not quite apparent.

(7) Lastly, there are a few recorded examples of linear nævus made up of sweat-glands or of sebaceous glands.

Dr. Paul's case may, I think, be excluded from the group of pilo-sebaceous or tricho-epitheliomata. Although it has not a pronounced gland structure it may, perhaps, be placed among the sweat-gland tumours of class 6. The massing of the growths in one area is suggestive of a segmental or linear nævus, and unlike that of the typical case of "*Lymphangioma tuberosum multiplex*" of Kaposi. It is closely similar in appearance and in its histological structure to a case of "*Hypertrophic u. multiple Cystadenoma der Schweissdrüsen*" recorded by Thimm.*

* *Arch. f. Derm. u. Syph.*, 1904, lxi, Hft. 1 and 2, p. 1. (The photograph and coloured drawing of Thimm's case reproduce almost exactly the appearances of Dr. Paul's case, and the microscopical sections seem to show very similar features.)