What we wished to convey is that the men who returned to their companies, free of all lice immediately became infested with lice anew; and being still infectious, infected a new generation of lice, with the result that, in due course, a recrudescence of the epidemic occurred. In fact, the contention we wished to make is that the analogy between the spread of typhus and the spread of malaria is a complete one. We recognize that we have offered no proof of this, and indeed it may be that the louse—such a predominating feature of the picture—has warped our judgement in this, our sole experience of the disease.

While on the subject may I elaborate the point we made in regard to the class of case we call "high tension type"? In this it is quite common for the pulse-rate never to go above 90 during the entire pyrexial period and to remain between 70 and 80 while the temperature remains on a plateau of 103° to 104° for several days. It is a feature so peculiar as to impress itself on the notice. "The exigencies of the service" prevent me from communicating with my late collaborator before writing this, but I feel sure he will concur with me.—I am, etc.,

P. C. T. DAVY,

London, W., Dec. 7th.

Major R.A.M.C.

BROMIDES IN EPILEPSY.

SIR,—Whether epilepsy is one disease or several diseases cannot be decided, cannot even be discussed, until we have some clear understanding of what is meant by a disease; and there is no such understanding. It is the business of our lives to treat diseases, but we none of us know what we mean by diseases; at least I know of no satisfactory definition of the term. Some months ago I offered to open a discussion on the matter at the Royal Society of Medicine, but that society was at the moment engaged in discussing something much more important and more practical—I think it was, What is good for a cough?—and my offer was courteously but firmly declined. I know what I mean by a disease, but I cannot explain it within the limits of a letter, and therefore it is from want of space, not from lack of courtesy, that I am unable to satisfy Dr. Humphrey Davy, and I trust he will forgive me.

Dr. Hume Griffith's statistics are scarcely convincing. His experience is that in 1900 a set of patients, whose number and the peculiarities of whose cases are not specified, were not treated with bromide, and had fits averaging 13.3 per month. In 1907 a set of patients, whose number is not specified and the peculiarities of whose cases are not specified, and who may or may not have been, but probably were not, the same set of patients as were observed in 1900, were treated by heavy doses of bromide, and the fits averaged 1.99 per month. Again, after seven more years, another batch of patients, some or all of whom may have been different from those in the previous batches, were treated with moderate doses of bromides, *plus* individual treatment, and the fits averaged 1.5 per month. I am afraid that from these data no trustworthy conclusion whatever can be drawn. The frequency of the fits varies in different patients from scores in one day to one or two in a year. It is clear that the presence in the first batch of a single case of unusual frequency and its absence from the later batches would account for the whole of the average difference, supposing the bromides to have had no influence whatever on the fits, or even supposing that it increased their frequency.

Dr. Stephens finds bromides of use in three ways: First, as a means of controlling the increasing frequency and intensity of the attacks; this, of course, assumes that the bromides do exercise this control. My case is that there is good evidence of the exercise of such control in some cases, but no evidence of any such control in other cases. Second, he uses it as an adjuvant in status epilepticus. Thus used, it certainly does no harm, but I could never satisfy myself that it increased appreciably the effect of the chloral. Third, he finds that epileptics, when temporarily suffering from other diseases, are often benefited by the addition of bromides to the mixtures prescribed for those diseases. But, in the first place, Dr. Stephens does not say that they are always benefited, and thus does not contravene my assertion; and, in the second place, it is, if not impossible, extremely difficult to disentangle the several effects produced by several drugs prescribed in a single mixture.

I do not deny, on the contrary, I strongly assert, that in some cases of recurring convulsion, which is what we mean by epilepsy, the discreet administration of bromides is followed by great diminution of the frequency of the fits, without any discernible ill effect; but I assert also that in the majority of cases either the fits are not diminished in frequency; or if they diminish in frequency they are increased in severity; or the diminished frequency of the fits is accompanied by a deterioration of mind which is too high a price to give for the lessened convulsions. My own practice, therefore, is to begin the treatment of most cases of epilepsy by administering bromides, but to watch their effect narrowly; to leave them off from time to time, and watch the effect of the omission; and unless I can satisfy myself that there is a clear gain from their administration, to leave them off altogether, and trust to careful regulation of the diet, attention to the bowels, regulation of exercise, and so forth.

I am surprised to hear from Dr. Hume Griffith that bromide acne is easily avoided by the administration of arsenic. My experience is that the combination of arsenic with bromides delays the appearance of acne, and lessens its severity, but does not prevent it; and it has seemed to me that in proportion as the acne is delayed and lessened, the other effects of the bromides are delayed and lessened.

In estimating the improvement or otherwise of a case of epilepsy, it is not enough merely to count the number of the fits. We must take into consideration also their severity, the increase or decrease of physical energy and of mental capability, the feeling of well-being or the reverse, and other factors; and it is only when all these factors are taken into account that we can strike a balance and say whether the life of the patient is really improved, is better worth living, is more advantageous upon the whole to himself, his family, and the society in which he lives. It is not easy to strike such a balance, but it is to this that our efforts should be directed. Diminution in the frequency of the fits, important as it is in itself, is not the only thing to be considered, and even if we can effect it, it may be bought too dear.—I am, etc.,

Parkstone, Dorset, Dec. 4th.

CHAS. A. MERCIER.

SIR,—Our knowledge of the action of bromine and the bromides upon man can hardly be regarded as adequate and satisfactory. On the one hand a distinguished authority like Dr. Mercier regards the war shortage of bromides as possibly a "blessing in disguise." On the other hand, as testified by general practice and tradition, and more particularly by the correspondents in your issue of December 4th, bromide undoubtedly exercises a considerable control over epilepsy. The main point at issue, I take it, is whether on the whole the bromides may not do more harm than good.

Apart from any special experience in epilepsy, most medical men meet with more or less severe cases of bromism—that is to say, bromism as described in the textbooks and familiar to our observation. Can we be sure, however, that we recognize at all adequately the remote evil effects of bromine compounds upon the human body? Bromine is a powerful drug, and it may be that by its use we are actually setting up various tissue degenerations and other results of unrecognized chronic poisoning. Dr. Hume Griffith mentions the fact that patients treated in the Lingfield Colony with heavy doses of bromide were like "logs" in school. Do the textbooks warn us against such a phenomenon as a result of bromide administration? It would be interesting to ascertain, were it possible, what is the terminal state of patients who have been dosed over long periods with bromides. It has always seemed to me that the last state of many epileptics may have been in reality attributable—in some degree at any rate—to the bromides which form the classical remedy in that malady. In some cases of unconsciousness or what may perhaps be described as cataleptic conditions under my own observation I have been able to trace a history of prolonged bromide administration.

At any rate, in the bromides we are using a powerful drug of whose remote action after prolonged dosage we cannot be said to have an adequate knowledge. Comparative pathology has long ago shown its destructive effect in small doses on internal organs.—I am, etc.,

London, W., Dec. 6th.

DAVID WALSH.