ANOPHELES BREEDING AMONG WATER LETTUCE—A NEW HABITAT.

By JAMES ZETEK, B.A.,

Entomologist, Ancon, Panama Canal Zone.

With the filling up of Gatun and Mira Flores Lakes, water lettuce (*Pistia stratiotes*, Linné) and water hyacinth (*Eichornia crassipes*, Solms.) rapidly increased in numbers until large areas became completely covered by them. Masses of these plants would become detached and float about in these lakes. It was found necessary to destroy these "floating islands," not because they were serious mosquito habitats, but because they interfered with navigation and the operation of the spillways. The water lettuce is the habitat of the very specialised larvae of *Mansonia*, of which *titillans* is the commonest member. Knab (1913) and Busck (comments appended to Mr. Knab's paper) refer to the changes in the mosquito fauna brought about by the rapid increase in *Pistia*. Both *Mansonia titillans*, Walker, and *Aëdomyia squamipennis* (Lynch-Arrib.) Theo., have been collected by me in very large numbers since 1912, and to-day *Mansonia* is the dominant species caught in the army barracks on the west side of the canal. Knab expressed the opinion that perhaps measures would have to be taken to destroy the *Pistia* habitats; now that we have found *Anopheles* larvae in them, this prophesy comes closer home.

The only question involved is how much of a menace are these floating islands to our towns. It is my belief they are a menace to the towns on the west side of the canal, and if allowed to float toward Paraiso and Pedro Miguel, or close to Gatun, would be a source of danger there. It has been shown conclusively that our common malarial *Anopheles* do fly more than a mile (Zetek, 1915); in fact, it appears that they will fly as far as they must in order to get food. Should Las Cascades become a negro settlement, the dangers from malaria would be increased. I doubt if we can depend upon screening and mosquito-catching indoors to control the *Anopheles* and malaria at such a settlement. At any rate, the idea seems to be getting firmly established that the sanitation of the Panama Canal Zone is so efficient that there is virtually little danger from the *Anopheles*, and as a result our people are becoming somewhat negligent. It will depend mainly upon the cumulative evidence gathered by the District Sanitary Inspectors, whether or not these floating islands will need to be controlled.

That this evidence may be available and accurate, it is very necessary that the sanitary inspectors make it their practise to take ample field notes. These should contain as complete a statement of the particular habitat, date and locality added thereto, and specimens of the larvae should be sent to the laboratory for the accurate determination of the species. At the close of each month a general summary of these notes should be made, a copy of which should be sent to the entomologist. It is a regrettable fact that much valuable information is lost or buried in the heads of observers.

In 1918 I found *Anopheles* larvae among water lettuce at Juan Mina, a citrus plantation about five miles up the old Chagres River. On 21st November 1919, Messrs. Picket and Tolar, sanitary inspectors at Pedro Miguel, Dr. D. P. Curry,

Assistant Chief Health Officer, and I, made an inspection of floating islands at Gamboa, C. Z., both in the canal prism and across in the bayous; we went as far as Empire. Mr. Pickett previously reported the presence of *Anopheles* larvae in these islands at this place, and that they were going southward toward his station. Our joint inspection revealed larvae of *Anopheles albimanus*, Wied., and *A. tarsimaculatus*, Goeldi, to be plentiful among the leaves of the water lettuce. An unidentified *Culex* was also present.

This particular habitat is very favourable to the *Anopheles*. The larvae are afforded much protection from the schools of young fish that usually feed upon insect larvae. They are also protected to some extent from predaceous insect larvae, such as those of dragon-flies. Shade is another important factor in this habitat, and this means a more even temperature, and an almost total absence of the direct hot rays of the sun. Wave action does not disturb these larvae very much; but the most important advantage to them is the exceptionally favourable presence of oxygen, and this certainly must be effective in prolonging the life of the larvae and pupae. This factor is very easily explained. The *Pistia*, being a green plant, gives off oxygen. Its leaves are at or just above or below the surface of the water. Therefore this oxygen enriches the local atmosphere available to the larvae, and some of it is undoubtedly taken up by the water and made use of by the larvae through their cutaneous respiration.

During the summer of 1911, while engaged by Dr. S. A. Forbes to study the *Anopheles* at Havana, Illinois, a similar condition was found to exist, excepting that in place of *Pistia* the plant in question was the tiny *Lemna*. This plant covered the surface of the water in the bayous, ponds and quiet recesses of the rivers. The Chatagua grounds at Havana were invaded each year by *Anopheles*, but their source was not known until we found them in and among the floating islands of this *Lemna*. These islands broke away from the almost continuous mass across the river from Havana and by current and wind were swept across this deep body of water and came to rest along the shores of the Chatagua grounds; here the larvae developed, pupae formed and adults emerged in rapid succession.

There is one important aspect of this sort of habitat which must not be overlooked. It is that wind and current detach portions of this floating vegetation and sweep it across deep bodies of water to new localities, where were it not for this exceptionally good vehicle these larvae would be unable to cross the barrier of deep water.

Searching through the literature available at our laboratory, I find that Ingram and Macfie (1917) found larvae of *Anopheles costalis*, Lw., and *A. marshalli*, Theo., in exactly the same sort of habitat, composed of the same species of *Pistia*, at Christiansborg, near Accra, West Africa. Associated with them were: *Aëdomyia africana*, Nev.-Lem., *Mimomyia splendens*, Theo., *Culex quasigelidus*, Theo., and *Mansonioides africanus*, Theo.

In a previous paper, in which he discusses the limitations of kerosene as a larvicide, Macfie (1917) states (p. 278) that *Mansonioides africanus* flies over one mile and that its breeding places can be located with fair accuracy, inasmuch as they are composed of *Pistia*. On page 278 he states it is useless to hope for a total abolition of such mosquitos by ordinary oiling, and that a layer of kerosene (p. 294), no matter

how thick, applied to the surface of the water in which larvae of *M. africanus* live is absolutely innocuous. He is right, and this is corroborated by our observations upon *Mansonia titillans*, which is our counterpart to his *M. africanus*. But nowhere in this article, nor in the preceding one, does he touch upon the control of the *Anopheles* which live in this same habitat, nor does he refer to the use of emulsions which mix with the water and kill by contact, or the destruction of the *Pistia* by means of sodium arsenite or their removal by boats.

The logical control is to destroy the habitat. This may be costly and prolonged, and perhaps prohibitive where large numbers of bayous, ponds, cut-offs, etc., exist. If *Anopheles* alone are to be controlled, spraying with a phenol-resin soap emulsion every six days will probably be all that is necessary. It may even be effective against the *Mansonia*.

Macfie (1917) goes into great detail in his discussion of the favourable oxygen factor in this *Pistia* habitat and relates experiments made with certain mosquito larvae which corroborate his contentions. That this exchange of gases is a favourable one is quite certain. Our common *Anopheles* have been known to stay on the bottom for five minutes; if through the presence of green plants they have a more favourable habitat, then it follows that not only may cutaneous respiration be much more prolonged, but the general death-rate among the larvae may be greatly reduced. This death-rate is reduced also because the leaves of the plants protect the larvae from the numerous swimming enemies in the water, and from the hot, direct rays of the tropical sun.

References.

INGRAM A. & MACFIE J. W. S. (1917). The Early Stages of Certain West African Mosquitos.—Bull. Ent. Research, viii, pt. 2.

KNAB, Frederic (1913). Changes in the Mosquito Fauna of Panama.—Proc. Ent. Soc. Washington, xv, pp. 41–43.

MACFIE J. W. S. (1917). The Limitations of Kerosene as a Larvicide, with some Observations on the Cutaneous Respiration of Mosquito Larvae.—Bull. Ent. Research, vii, pt. 3.

ZETEK, J. (1915). Behavior of *Anopheles albimanus*, Wied., and *tarsimaculata*, Goeldi.—Ann. Ent. Soc. America, viii, pp. 221–271.