1910	α	δ	$\log r$	$\log A$
Dez. 19	4 ^h 34 ^m 25 ^s	+0° 56' 5		
21	32 43	0 57.4	0.3162	0.0650
23	31 8	0 59.5		
25	29 37	1 2.8	0.3148	0.0695
27	28 14	1 7.3		
29	4 26 57	+1 13.0	0.3134	0.0751
Gr. 9 ^m 6. Präzession bis 1855.0 — 2 ^m 51 ^s — 5'7.				

(97) Klotho.

Nov. 19	5 18 34	+0 54.1	0.2979	0.0300
21	17 22	0 43.3		
23	16 5	0 33.6	0.2980	0.0253
25	14 42	0 24.9		
27	13 15	0 17.4	0.2982	0.0218
29	11 44	0 11.0		
Dez. 1	10 9	0 5.9	0.2985	0.0197
3	8 31	+0 2.1		
5	6 52	—0 0.4	0.2988	0.0191
7	5 11	0 1.5		
9	3 30	—0 1.2	0.2992	0.0199
11	1 50	+0 0.4		
13	5 0 11	0 3.4	0.2996	0.0222
15	4 58 34	0 7.7		
17	56 59	0 13.3	0.3002	0.0260
19	55 27	0 20.3		
21	54 0	0 28.4	0.3008	0.0311
23	52 37	0 37.9		
25	51 19	0 48.5	0.3014	0.0375
27	50 6	1 0.2		
29	4 49 0	+1 13.0	0.3021	0.0451

Jurjew-Dorpat, 1910 Sept. 8.

1910-11	α	δ	$\log r$	$\log A$
Dez. 31	4 ^h 47 ^m 59 ^s	+ 1° 26' 8		
Jan. 2	4 47 7	+ 1 41.6	0.3029	0.0537
Gr. 9 ^m 0. Präzession bis 1855.0 — 2 ^m 49 ^s — 4'5.				

(356) Liguria.

Nov. 21	5 29 55	+37 45.5	0.3194	0.0666
23	28 23	37 53.4		
25	26 43	38 0.8	0.3195	0.0608
27	24 56	38 7.2		
29	23 3	38 12.6	0.3196	0.0564
Dez. 1	21 4	38 17.1		
3	19 1	38 20.7	0.3198	0.0533
5	16 54	38 23.1		
7	14 44	38 24.6	0.3201	0.0516
9	12 33	38 25.0		
11	10 21	38 24.4	0.3204	0.0514
13	8 10	38 22.7		
15	6 0	38 20.0	0.3208	0.0527
17	3 54	38 16.4		
19	5 1 52	38 11.7	0.3212	0.0555
21	4 59 51	38 6.2		
23	57 54	37 59.9	0.3216	0.0597
25	56 5	37 52.7		
27	54 23	37 44.9	0.3222	0.0653
29	52 49	37 36.4		
31	51 22	37 27.3	0.3227	0.0721
Jan. 2	50 4	37 17.8		
4	4 48 55	+37 8.0	0.3233	0.0801
Gr. 9 ^m 4. Präzession bis 1855.0 — 3 ^m 46 ^s — 4'0.				

Eugen Büss, stud. math.

The periods of some variables.

By J. T. Hedrick.

RR Persei.

A maximum was predicted for RR Persei (2^h 21^m +50°) for Febr. 3 of this year (V. J. S. 44.325). But on Febr. 7 the magnitude was only 13^m. The period used for the prediction, 372.6 days, was derived by dividing by 47 the interval between 1904 Dec. 26, J. D. 2416841, the well determined maximum by Dr. Graff, and 1857 Jan. 16, J. D. 2399331, when RR was estimated as 8^m5 at Bonn (Hbg. Mitt. 8.22). The divisor 45 gives 389 days as the length of the period. This period fits the observations in A. N. 164 p. 191, 207 and 215, those published by Prof. Whitney in A. N. 169.279 and 176.65 and observations by the same and Miss Furness in 1909 and this year, kindly communicated by letter. These last observations seem to indicate a period possibly one day less, or 388 days. A good determination of this year's maximum should give the length of the period more exactly, but daylight prevented it in this latitude.

RT Geminorum.

The accessible observations of RT Geminorum (6^h 40^m +18°) are those in A. N. 165.31 and 166.373 and some

made here. On 1908 Jan. 6, the magnitude was 9^m and a maximum must have been near. On 1909 April 26 the magnitude was 13^m. Five observations from 1909 Dec. 20, to 1910 April 9, show a decrease of about half a magnitude a month, with a maximum either in the middle or at the end of December. A period of 355 days fits the observations fairly well but one twice as long is not excluded. Supposing the variation nearly regular, the following tentative elements may be assigned.

$$\text{Max.} = \text{J. D. } 2417947 + 355^d (710^d?) E.$$

Hence the next maximum can be expected in December of this, or next, year.

The variable is 2'8 North of BD +18°1340 and is a little North of the centre of a triangle formed by BD +18°1340, +18°1343 and an 11^m which precedes the first 15^s, 3'4 North. A double star, 12¹/₂^m and 13¹/₂^m, follows RT 3^s, 0'8 South.

RV Geminorum.

Besides the observations in A. N. 166.45, RV Geminorum (7^h 11^m +24°) was observed here 1908 Jan. 3, as

$0^m.2$ brighter than BD +24°1596. In March and April, 1909, it was certainly faint and may have been invisible, but the power used then would not have separated it well from its companions. It was invisible ($<13^m.5$) in the middle of May, 1909, and in January, February and March of this year, though it was suspected on Febr. 10. On April 8 it was 13^m . Observations about every ten days showed a rise by May 15 to $0^m.1$ fainter than BD +24°1596 and by May 28 to equality with +24°1613 and +23°1658, or 10^m . The only observation possible since then, June 7, was made before daylight had altogether disappeared and in an unfavorable sky. It gave RV as probably fainter than the two stars last mentioned. Hence 1910 June 1, may be assigned as a probable date for this single known maximum. The period seems to be fairly well determined from the dates of equal brightness. The elements are

$$\text{Max.} = \text{J. D. } 2418824 + 288^d E.$$

The variable forms the following vertex of a nearly equilateral triangle with its base facing South and a length of side of about $25''$. The companions are 13^m and are s. pr. and n. f. with regard to each other. BD +24°1598, $8^m.5$, precedes RV 17^s , $4'.4$ North and +24°1600, $8^m.5$, follows it 4^s , $9'.3$ North.

RR Monocerotis.

$$\begin{array}{llll} 1900.0 & 7^h 12^m 22^s & (+3^s.10) & +1^o 16'.4 \quad (-0^o.10) \\ 1855.0 & 7 \ 10 \ 2 & & +1 \ 21.0. \end{array}$$

At most only a slight variation has been found for the star in the place heretofore given for RR Monocerotis, which should be 5^s and $0'.2$ less. The true variable was $13^m.5$. 1906 Nov. 24. It was not noticed 1908 Jan. 2 and 1909 Mar. 28 and was not seen, though looked for, 1908 April 23 and

Georgetown College Observatory, 1910 June 10.

May 1 and 1909 Mar. 21, April 22 and May 2 and 11. It was recognized as the variable when first seen again, 1910 Jan. 3. It was then $1\frac{1}{2}^m$ fainter than the following star namely 11^m , or brighter. By Mar. 7 it had sunk to $13\frac{1}{2}^m$.

The accompanying map gives all the stars visible in the 12-inch telescope here except a few of the very faintest. Three such were suspected within $2'$ of RR but the situation of the latter with regard to the following star and to the two stars South and a couple of seconds preceding prevents their being confused with it. The field can be recognized easily by the group of BD stars in the northern part. The star $1'$ North of the close pair of these may possibly be slightly variable, for it often failed to be noticed.

The recent maximum can not be determined from the observations here. It must have occurred near 1910 Jan. 3. In Harv. Ann. 55.141 are given several maxima determined at Harvard, compared with the elements

$$\text{Max.} = \text{J. D. } 2413143 + 392^d E.$$

The observations given in the announcement of discovery, A. N. 161.427 and the absence of RR from the Algiers photograph of 1902 Febr. 8 are not inconsistent with the Harvard elements. But J. D. 2417880 + $400^d E$ will reduce the residuals considerably, except for the Harvard maximum of 1894 Nov. 17. In Harv. Ann. 57.237 a list of comparison stars is given with their photometric magnitudes.

John T. Hedrick, S. J.

Safaríks Beobachtungen der veränderlichen Sterne.

Von Lad. Praška.

Der erste Band meiner Bearbeitung der großen Beobachtungsreihe von Prof. A. Šafařík ist im Verlage der Königl. böhm. Gesellschaft der Wissenschaften in Prag unter dem Titel: »Untersuchungen über den Lichtwechsel älterer veränderlicher Sterne« erschienen. Um über den Inhalt dieses Bandes die Interessenten zu informieren, lasse ich hier eine kurze Zusammenstellung der abgeleiteten Epochen des größten bzw. kleinsten Lichtes folgen. Die Beobachtungsreihen selbst sind meistens sehr homogen, so daß sie einen wertvollen Beitrag für Spezialarbeiten bilden. Über die in dieser Übersicht fehlenden Sterne 190.1904 Cassiopeiae und T Persei wurde in A. N. 4354 und 4434 bereits Bericht erstattet.

Variabilis	Größe	Phase	J. D.	Bemerkungen
T Cassiopeiae	$7^m.8$	M	2408962	flach
	10.6	m	9205	unsicher
	7.7	M	9420	flach
	7.5	M	2410780	
	7.6	M	1218	
	10.6	m	1400	

Variabilis	Größe	Phase	J. D.	Bemerkungen
T Cassiopeiae	$7^m.2$	M	2411634	
R Andromedae	6.5	M	2409155	
	6.0	M	9560	
	6.8	M	9950	unsicher
	6.6	M	2410802	
	6.5	M	1240	
	5.2	M	1630	
	7.3	M	2067	
	5.7	M	2463	
	8.3	M	2408300	unsicher
	7.5	M	8887	spitz
S Cassiopeiae	8.5	M	9510	flach
	8.3	M	2410120	flach
	8.6	M	0750	unsicher
	7.9	M	1328	Wellen
	7.8	M	1931	unsicher
	8.2	M	2552	flach
R Arietis	7.6	M	2410882	vergl. Bem.