not closely punctate. Elytra conjointly oval, one-fourth wider at base than thorax, umbone prominent; disk evenly, closely and rather finely punctate, the umbone and three or four elongate calloused spaces near middle of each elytron smooth. Under surface very finely and sparsely punctate; prosternum forming a large flat quadrate plate between the front coxae. Length 2.7-3 mm.

Described from three specimens from the Isle of Pines received, among other material for naming, from Prof. J. R. Watson, of the Florida Agricultural Experiment Station. They were labelled "on citrus," and Prof. Watson writes that he has considerable correspondence with citrus growers on the Isle of Pines, who are unable to get satisfaction from the Cuban station. Four species of *Colaspidea* have hitherto been described from North America, all from Southern California or adjacent islands. According to Horn,²³ others are known "from Southern Europe and the circum-Mediterranean region generally." Since Leng and Mutchler mention no species of the genus in their list of West India Coleoptera, *insularis* is apparently the first one to be known from the Atlantic region of North America.

Haltica schwarzi Blatch.—Described²⁴ from the east shore of Lake Okeecho-bee. Two specimens were secured on March 4 while sweeping low huckleberry near the edge of a brackish water marsh just east of Ft. Myers.

TWO FAMILIES OF INSECTS NEW TO BRITISH TERTIARY STRATA

BY T. D. A. COCKERELL,

University of Colorado.

When recently examining some hitherto overlooked fossil insects in the British Museum, I found two specimens collected by Mr. a'Court Smith in the Isle of Wight, representing families not yet recorded from the British Tertiary rocks. They come from the Bembridge series, at the famous Gurnet Bay locality.

Fig. 1, wing of *Vectevania vetula*; Fig. 2, wing of *Nemoura priscula*.

EVANIIDAE.

Vectevania n. gen.

Related to *Protofoenus* Ckll., from Burmese amber, but differing especially in the complete third discoidal cell and the deep triangular stigma. The second transverse-cubital nervure is so faint as to be practically obsolete, and there is perhaps a slight indication of the third, too vague to be positively recognized.

²³Trans. Amer. Ent. Soc., XIX, 1892, 204.

²⁴Can. Ent., XLVI, 1914, 141.

The position of the second is as in the living *Neuraulacinus braunsi* Kieffer. The small cuneiform first discoidal cell is characteristic, and the comparatively distinct venation of the hind wings is a primitive feature, lost in the modern *Aulacinus*, but preserved in *Neuraulacinus*.

Vectevania vetula n. sp.

Head narrower than thorax; thorax elongate-oval, 2 mm. long; abdomen claviform, nearly 3 mm. long, obtuse at apex; anterior wing about 4.5 mm. long, clear hyaline, with ferruginous nervures; stigma margined with brown, apparently originally all brown.

British Museum, In. 20535.

PERLIDÆ.

Nemoura (sens. lat.) **priscula** n. sp

Anterior wing, length about 5.4 mm., width nearly 2 mm.; colorless. About 1.7 mm. of the base is lost. The venation is very simple, but similar in principle to that of *Nemoura*. From the fork of the radial sector to the apex is nearly 3 mm. Comparing the venation with recent species, I found nothing very close, and it is probable that a distinct genus is indicated. Some of the species from Prussian amber are interesting for comparison. Thus *Taeniopteryx elongata* Berendt has the cross-nervure at the end of the subcosta, but it is oblique. *Leuctra gracilis* Berendt and *Leuctra linearis* Berendt have this cross-nervure well before the end of the subcosta, in *gracilis* producing an angulation of the subcosta, but not in *linearis*. On the other hand, *L. gracilis* agrees with the English fossil in having the cross-vein connecting R with R₃ beyond the fork of R₃; in the other two amber species it is at the fork. Our fossil nearly agrees with *L. gracilis* in the position of the cross-vein between R₃ and M.

British Museum, In. 17498.

Thirteen species of Perlidæ have been found in Baltic amber, and one in the Oligocene of Rott, but strangely enough none in the American Tertiaries.

UNDESCRIBED LEPIDOPTERA IN THE CANADIAN NATIONAL
COLLECTION.

BY J. MCDUNNOUGH,

Entomological Branch, Ottawa.*

For some time there has been a strong feeling among Canadian entomologists that the types of as many as possible of our Canadian insects should be deposited in the National Collection at Ottawa. During the past half century, owing to the lack of taxonomists in Canada, a large proportion of Canadian insect material has been sent to the United States or Europe for identification and as a result the types of numerous undescribed species have been retained in these countries. With the growth of taxonomy and the discovery of numerous closely allied forms which heretofore had been grouped

*Contribution from the Entomological Branch, Department of Agr., Ottawa