THE NATURE OF SECONDARY VALENCE

BY HOMER W. SMITH

Part I	Preliminary Communication,	Jour. Phys. Chem., 25, 160 (1921).
Part II	Systems water: xylene and water: chloroform,	Ibid., 25, 204 (1921).
Part III	System water: ether, Supplementary note on extrapolation method of correction,	Ibid., 25, 605 (1921). Ibid., 25, 616 (1921).
Part IV	System glycerine: acetone,	Ibid., 25, 721 (1921).
Part V	Systems containing water and paraffin oil; petroleum; tri-chlorobenzene; <i>iso</i> -amyl phenyl ether; toluene; benzene; bromo-benzene; carbon tetrachloride; dichlorobenzene; <i>n</i> - butyl bromide; <i>n</i> -butyl ether; ethyl- ene chloride; <i>sec</i> -octyl alcohol; <i>n</i> - butyl alcohol; <i>iso</i> -butyl alcohol; amyl alcohol and bromoform,	Ibid., 26, 256 (1922).

VI. Summary and Discussion

I. Summary

This investigation was prompted by the desire to ascertain something of the nature of the forces involved in liquid solutions. To this end the distribution of various organic solutes between immiscible solvents was chosen as the most promising and direct experimental procedure. In all, about twelve hundred final determinations have been presented, involving twenty-one systems each consisting of two immiscible liquids.

These results as they stand are not comparable with one another because of the complications introduced by dissociation, association and the mutual solubility of solvents. It has been shown, however, that these complications can be eliminated in part by an empirical method of correction, and the partition coefficients so obtained put on a mathematically comparable basis, even though their physical-chemical significance is still obscure. This method of correction consists

in determining by interpolation the partition coefficient when the concentration in one solvent has a fixed arbitrary value. The interpretation of the results obtained by means of this method of correction is independent of the arbitrary value chosen for interpolation.

The principal points established by a consideration of the experimental data are as follows:

When the partition coefficients of various substances for any one system are compared on a basis of molecular volume, it is found that they may be covered by the equation

$$\text{Log } 100 \mathfrak{K} = \frac{V_m}{60.00} \pm a$$

where V_m is the molecular volume at the boiling point.

A limited number of values of a are required to cover the behavior of all substances in any one system. (All substances covered by the same value of a in any one system have been called a "series," and the corresponding values of a the "series constants.")

The various values of a (or series constants) in any one system have been found to be related to each other by simple proportion.

It has also been found that the series constants for any one series in twenty systems containing a common solvent (water) are related to each other by simple proportion.

The above facts are interpreted as indicating that solubility is a function of the molecular volume of the solute. It is believed that the stepwise phenomena of series, as shown by the integrally related values of a in the above equation, are attributable to discrete or abrupt variations in the intensity of the intermolecular forces associated with various molecular species.

2. Homogeneity and Three-Dimensional Symmetry

Before discussing the fundamental significance of these results, it will be necessary to consider again certain points,

already discussed in Part I, disclosed by molecular volume relations. It has been said that the experimental facts relating to the molecular volumes of liquid chemical compounds indicate that every atom contributes a definite volume to the molecular volume. Some atoms vary in their atomic volumes as a result of constitutive influences (that is, interactions attributable to the chemical arrangement of the atoms in the molecule) but these constitutive variations in atomic volume are well defined and are related in no way to what we may conceive as spatial configuration. A consideration of molecular volume relations shows that the configurations suggested in structural formulae indicate only the chemical properties of the constituent atoms in the molecule, and in no way define the actual spatial relationships of the various parts or of the whole. The molecular volume is the simple sum of the volumes of the constituent atoms, and there is no additional space which can be attributed to interatomic or intermolecular crevices resulting from fixed spatial configurations as suggested by these formulae. To repeat the example previously used, there is no hole inside the benzene molecule or cracks or crevices between the atoms, as there would surely be if the molecule consisted of rigid spheres of carbon with smaller rigid spheres of hydrogen tacked on at various places. The principle of the perfect sum implies that the six carbons and their attendant hydrogens make up a domain which has the same shape, in the electro-dynamic sense, as has each of the components, much as though we had put several little spheres together to make one big sphere.

We have seen in the data presented in these papers that molecular volume appears to be one of the factors which determine the behavior of various molecular species in homogeneous liquid systems. This determinative rôle of molecular volume is independent of the chemical nature of the molecular species involved and constitutes new evidence of a most conclusive kind in regard to the shapes of molecules. We may summarize this evidence as follows: substances in the homogeneous liquid phase behave toward their environment as

though every molecule homogeneously filled a space which is symmetrical in three dimensions.¹

An assumption somewhat of this nature has been made from time to time by mathematicians when considering the attraction of one molecule for another, but the chemist, in the absence of direct evidence to the contrary, has persistently clung to notions of chemical configuration and spatial relationships.² The mathematician speaks of point-to-point action, and avoids the difficulties arising from chemical considerations by arguing that this alleged point-to-point action is only a statistical result. As to how far this is so, it is hard to say because of course we cannot avoid statistical observation and we do not know the exact extent to which the real molecular domain is increased by the translational motion of heat. But the specific nature of the optical properties, the compressibilities and more especially the radiation of heat from liquids and solids indicate that only a small fraction of the apparent molecular domain is to be attributed to translational motion. It seems instead that when there is an expansion of the molecular domain on the absorption of energy, it is to be attributed to an actual expansion of the size of the molecule.³

But even if we agree to subtract a certain amount from the observed molecular volume to allow for space taken up by translational motion, we must still apply the principle of homogeneity and three-dimensional symmetry to the remaining real volume, for otherwise molecular volume could not

¹ Attention is called to the phrase "in the homogeneous liquid phase." The work of Langmuir (*Jour. Am. Chem. Soc.*, **39**, 1848 (1917)) and Harkins and his collaborators (*Ibid.*, **39**, 541 (1917)) shows that molecules at the surfaces of liquids are oriented and possess more or less definite shapes, but neither Langmuir's nor Harkins' work applies to molecules situated within a homogeneous liquid where the surrounding field of force is symmetrical.

² Among others who have argued from this premise are S. C. Bradford: *Phil. Mag.*, **38**, 696 (1919); J. J. Thomson: **27**, 757 (1914); W. C. McC. Lewis: **28**, 104 (1914); W. D. Harkins and H. H. King: *Jour. Am. Chem. Soc.*, **41**, 970 (1919).

³ See Gervaise LeBas: "The Molecular Volumes of Liquid Chemical Compounds," p. 254 (1915).

function as a determinative factor in solubility in the precise manner in which it does. So taking into account the additive nature of molecular volume and the rôle of molecular volume in homogeneous liquid solutions, we must look upon this homogeneity and three-dimensional symmetry of liquid molecular domains as an actual fact and not an apparent one. We must picture the electrons of the atoms and molecules which make up a mass of liquid as filling, by their electrodynamic activity, the entire liquid domain, being at equilibrium at all points, and leaving no crevices or dead-spaces wherein other electrons or atoms or molecules can be inserted without disturbing this equilibrium.

3. The Nature of the Intermolecular Forces

Besides the size of the molecule, there is yet another factor which influences its physical relations with other molecules; this is the intensity factor in the intermolecular forces. It has already been pointed out that slight variations in the arrangement of atoms result in the most profound variations in molecular behavior as regards nearly all physical properties and we see this fact illustrated especially well in relative solubility. The striking thing about these variations in this case is that they consist of abrupt or stepwise variations from one condition to another—the various conditions being related to each other by simple proportion.

As the evidence from partition coefficients now stands it is apparent we are dealing not only with the properties of undissociated molecules, but with associated complexes and ions and hence with the chemical forces underlying association and dissociation, as well. But despite the fact that this complication precludes more direct physical-chemical consideration, it gives the results greater physical significance because it brings all these phenomena under one category without in any way invalidating the physical interpretations based upon the results.

This simple stepwise variation in behavior mentioned above is shown by a large variety of substances: compounds

containing carbon, hydrogen, oxygen, nitrogen, chlorine, bromine, iodine and sulphur in a variety of chemical combinations. Hence this phenomenon cannot be attributed to peculiarities in molecular or atomic structure because it appears to be wholly independent of the kind of atoms or molecules involved. It must instead be attributed to something common to all atomic species and inherent in atomic structure itself. Undoubtedly it is to be associated with the now generally accepted idea that atoms absorb and emit energy discontinuously, in whole multiples of a unit of energy $h\nu$, or Planck's quantum. This unit has a different value for every frequency of radiation, ν , but it involves a universal constant, h . It is to be noted that this quantum is a unit of action, since it has the dimensions of energy multiplied by time, whereas the element of time does not enter into our present consideration. We are comparing various physical systems in their characteristic and essentially fixed equilibrium states.

Bohr has assumed that the electron in a hydrogen atom could revolve about the nucleus only in certain orbits which represent conditions of stable equilibrium. On the absorption or emission of a quantum of energy the electron would jump from one orbit to another and thus enter another stable state.¹

No one has, as yet, satisfactorily explained the existence of stable states, though this conception underlies Bohr's explanation of the quantum and the theories of atomic structure as well. Thus Langmuir² has suggested that each electron occupied a more or less permanent position in the atom which he called a cell, the various cells being arranged in symmetrical positions around the nucleus of the atom. In discussing the properties of the electron he has said, "It seems that the electron must be regarded as a complex structure

¹ For a general discussion of the quantum theory see "The Quantum Theory," Edwin Plimpton Adams: Bull. of the National Research Council, Vol. I, Part 5, No. 5; Wm. C. McC. Lewis: "A System of Physical Chemistry," Vol. III, "Quantum Theory" (1919). Also Max Planck: Phil. Mag., **28**, 60 (1914). On the application of the quantum theory to radiation from liquids and solids see H. Stanley Allen: *Ibid.*, **35**, 338, 404, 445 (1918).

² Jour. Am. Chem. Soc., **41**, 868 (1919).

which undergoes a series of discontinuous changes while it is being bound by the nucleus or kernel of the atom."¹

Thomson² has considered the behavior of the electron on the hypothesis that the field of force in the immediate vicinity of the positive charge changes alternately from attraction to repulsion. At great distances these forces would reduce to the inverse square law but for atomic distances the change in sign would result in a series of positions of zero force which would be positions of stable equilibrium. Other assumptions of a more or less arbitrary nature have been made by Nicolson,³ Brillouin,⁴ Allen,⁵ Lamar⁶ and others to account for the discontinuous absorption and emission of energy as postulated under the quantum theory.

4. The Conservation of Energy and Inter-electronic Equilibration

The intermolecular forces are commonly supposed to have their origin in the so-called "chemical" or valence electrons. No one has advanced any very definite suggestion as to how these electrons function while holding molecules together in the liquid state. It is usually assumed that the cohesive forces are due to stray fields of force, residual affinity, etc.⁷

It does not seem possible that an electron may be so situated that its electrostatic or electro-magnetic affinities are satisfied, and that it may still be able to permanently influence, without dynamic expense, other electrons by virtue of "stray fields of force." It seems much more consistent

¹ Phys. Rev., **8**, 300 (1919).

² J. J. Thomson: Phil. Mag., **37**, 419 (1919).

³ J. W. Nicolson: Ibid., **28**, 90 (1914).

⁴ Marce Brillouin: Comptes rendus, **168**, 1318 (1920).

⁵ H. Stanley Allen: Phil. Mag., **41**, 113 (1921).

⁶ Joseph Lamar, Ibid., **42**, 592 (1921).

⁷ Albert P. Mathews: Jour. Phys. Chem., **20**, 554 (1916); Herbert Chatley: Proc. London Phys. Soc., **29**, 206 (1916); Irving Langmuir: Jour. Am. Chem. Soc., **39**, 1848 (1917); S. C. Bradford: Phil. Mag., **38**, 696 (1919); W. C. McC. Lewis: **28**, 104 (1914); Wm. D. Harkins and H. H. King: Jour. Am. Chem. Soc., **41**, 970 (1919).

to believe that a principle of conservation applies to all inter-electronic relations, including the intermolecular forces; that the intermolecular forces are expendable by the molecule to a definite and limited extent dependent on the electronic fabrication of the molecule, just as it has always been assumed in the case of the interatomic forces. Since the intermolecular forces have their genesis in the electrons of the constituent atoms, we can arrive at such a position simply by assuming that a state of conservative equilibration exists between all the electrons which are involved in the interatomic and intermolecular forces. That is, in the atom or molecule, or in any homogeneous system of atoms or molecules, all the electrons exert forces on all the other electrons, holding each other in certain positions of stable equilibrium. If any of the physical conditions of the system are altered by the absorption or emission of energy, or by admixture with another system, certain electrons are displaced and take up new positions of equilibrium. On the assumption that energy is added to or taken from a system only by a discontinuous process, it would necessarily follow that the alteration of any system from one physical state to another would likewise be discontinuous. Various systems (or various molecular species) could be treated as a single system in various equilibrium states, and the various systems would bear definite relations to each other which would result in simple mathematical relationships between the partition coefficients, the frequencies of spectral lines and all other physical properties.

It should be reiterated that the experimental data from partition coefficients deal with various systems in their characteristic equilibrium states, and that the element of time, a necessary consideration in the process of change from one equilibrium state to another, is not involved. In this respect, these data are unique, for they demonstrate that the existence of these discrete differences in the equilibrium states of a given system, or of various systems, is independent of the time element. The evidence warrants the assumption that these discrete differences are attributable to some property in-

herent in the fundamental charges themselves. Some such assumption as that made by Thomson, namely that the field of force around the fundamental charges changes alternately from attraction to repulsion, appears to offer the only explanation of the facts.

Johns Hopkins University
School of Hygiene and Public Health
Department of Physiology
Baltimore
and
The Lilly Research Laboratories
Eli Lilly and Company
Indianapolis