ously. But where the gouge has to be used the joint movements are less free, and there is a tendency to ankylosis.

Notwithstanding, this operation is a great gain when the dislocation is single or, after good motion has been established on one side, when dislocation is double. When consolidation has taken place, and a new articulation has been formed, as in Fournier's case I never interfere; but at any period before consolidation is complete, the head of the femur may be drawn down after the trochanteric muscles have been divided, and with or without the gouge, the head of the femur may be safely and permanently restored to the acetabulum.

AN ANTISEPTIC TABLET FOR NASAL WASH.

BY JOHN EDWIN RHODES, M.D.

CHICAGO.

The necessity of a cleansing and an antiseptic wash for the nasal cavities is indicated in a variety of conditions. In cases of simple chronic catarrhal inflammation in which there is an excess of mucus which gathers and forms crusts in the cavities, often due to a slight nasal obstruction and accompanied with an unpleasant "catarrhal odor," the indications for treatment are often chiefly fulfilled by some antiseptic wash which will effectually dislodge the secretions and counteract the odor. In another class of cases instead of obstruction, we find an enlarged passage, atrophy of the membrane, and sometimes of the turbinated bones. Here even an exaggerated expiratory blast of air is inefficient in dislodging the secretion. This rapidly dries upon the surface of fine membrane, decomposes and causes pathologic changes in the tissues, and the patient gives as one of the most distressing features of the condition the bad odor that accompanies it. Here there is an indication for an agent that will both cleanse the nasal passages, disinfect them and stimulate the mucous glands. After operations of various kinds, as the removal of exostoses, the cauterization of the membrane, the removal of polypi, tumors, etc., an agent is needed that will render the parts thoroughly aseptic until healing has taken place. In operations about the fauces, the naso-pharynx, the pharynx, the tonsils, it is always advisable to use some strongly antiseptic solution as a spray or gargle for a few days succeeding the operation.

To meet such indications as have been briefly enumerated above the following formula has been found to possess the requisite agents in suitable proportions. The tablets can be employed in all conditions, and are unirritating if used in the strength of one tablet in from one-half to two-thirds of a glass of lukewarm water. In cleansing the nares it is much to be preferred that the patient should draw the water gently through the passages from the hand than to use any kind of a douche, as the latter in very many cases is improperly used, and the liquid is carried through the nasal passages with such force as to do local damage and perhaps reach the middle ear through the Eustachian tubes and set up a middle ear inflammation. This formula has been thoroughly tested by use in a large number of cases in private practice in the last two years, and is more satisfactory than any agents I have ever used for like purposes.

It has been put up in the form of compressed tab-

lets, which, as prepared by Gale & Blocki, are readily soluble, and in every way satisfactory. Each tablet contains the amount indicated in the formula.

DR. RHODES' NASAL TABLETS.

Potass. chlorate	2 1/2 grs.	15
Soda bicarbonate	10 grs.	60
Soda chlorid, C. P	10 grs.	60
Soda salicylate	5 grs.	30
Soda bichlorate	5 grs.	30
Thymol	1/8 gr.	007
Eucalyptol	1/4 min.	015

MYRINGITIS ACUTA—MYRINGITIS BULLOSA.

BY FRANK ALLPORT, M.D.

PROFESSOR OF CLINICAL OPHTHALMOLOGY AND OTOTOLOGY IN THE MINNESOTA STATE UNIVERSITY, PRESIDENT OF THE MINNESOTA STATE MEDICAL SOCIETY, ETC.

MINNEAPOLIS, MINN.

Miss M. C., age 20; left ear; general health not good; has had naso-pharyngeal colds for two or three weeks, accompanied by a hard cough. Nov. 11, 1895, at about 11 P.M., experienced suddenly a severe earache, which continued with increasing severity till the morning of November 12, when I was consulted. I found the view of the membrana tympanum obstructed by three large bullæ situated as follows: One at the upper anterior quadrant, extending down nearly to the inferior extremity of the handle of the malleus. This was large and deeply congested, the redness passing on to the meatal wall, as it joins the drum-head.

Another was situated at the junction of the drum-head with the posterior meatal wall, and was more on the latter than the former. This was also much inflamed and large. Another blister occurred exactly at the inferior extremity of the malleus, and was smaller than the others—quite white and not inflamed. A cluster of prominent blood-vessels extended from this to the posterior blister. The bullæ were separate and distinct, for I punctured all three, as each one required a separate incision to liberate its individual serous fluid. I was careful not to penetrate beneath the dermic layer of the membrane, and a thorough inflation of the middle ear with the catheter and air bag after the puncture was made evidenced an intact condition of the underlying membranous coats. The hearing in this ear was somewhat but not markedly reduced. The case was seen by Dr. W. N. Porteus, of this city, at my request.

November 13, I found the blisters all thoroughly collapsed; some membranous inflammation, no pain, slept well, hearing almost completely restored. November 19, drum-head fully restored to a normal appearance and hearing perfect.

THE DEGENERATE EAR.

Read in the Section on Neurology and Medical Jurisprudence, at the Forty-sixth Annual Meeting of the American Medical Association, at Baltimore, Md., May 7-10, 1895.

BY EUGENE S. TALBOT, M.D.

FELLOW OF THE CHICAGO ACADEMY OF MEDICINE.
CHICAGO.

(Concluded from page 58.)

Figures 10, 11 and 12 illustrate the different stages of excessive and arrested development of the helix. For the purpose of giving a better description of the stigmata of the ear I have divided the ear into upper, middle and lower third. Figure 10 shows arrest of the helix as far up as the upper third, while the upper part is well developed. Figure 11 shows the helix normally developed. Figure 12 illustrates the lower

third normally developed, while the upper middle and upper third are excessively developed.

When the whole or a large part of the helix is arrested the ear is usually (not always) larger and the antihelix is excessively developed. On the other hand, when the whole helix or larger part is excessively developed the ear is usually (not always) small and thick, with arrested antihelix. When the ear is at or near an angle of 90 degrees the helix is usually quite or entirely arrested or very thin. When it is small and at an angle of from 10 to 45 degrees it is usually well developed and thick. The root of the helix is sometimes arrested. When this is the case

Fig. 10.

Fig. 11.

it is lost in the floor of the concha. When normally developed it produces a distinct ridge extending across the floor and when excessively developed it divides the concha into unequal parts. In 465 criminals at Pontiac, there were 236 arrested, 55 medium, 174 excessive; 181 folded, 48 medium, 236 open. Root of helix, normal 157, abnormal 308. In 1,041 criminals at Elmira, there were arrested 771, medium 52, excessive 206; 294 folded, 747 open. Root of helix, 907 normal, 132 abnormal.

The most singular deformity of the helix is the tubercle of Darwin (Fig. 13), which is a little blunt point projecting from the inwardly folded margin or

Fig. 12.

Fig. 13.

helix. When present it is developed at birth and according to Prof. Ludwig Meyer, more frequent in man than in woman. Figure 13 shows ear and tubercle taken from Darwin's "Descent of Man."³ These points not only project in toward the center of the ear, but are often a little outwards from its plane, so as to be visible when the head is viewed from directly in front and behind. They are variable in size and somewhat in position, standing either a little higher or lower; and they sometimes occur on one ear and not on the other. They are not confined to mankind, for Darwin observed a case in one of the spider

³ The Descent of Man, page 15.

monkeys (*ateles bulzebuth*) in the Zoological Gardens. Dr. E. Ray Lankester tells of another case in a chimpanzee in the garden at Hamburg. The helix obviously consists of the extreme margin of the ear folded inward, and this folding appears to be in some manner connected with the whole external ear being permanently pressed backward. In many monkeys which do not stand high in the order, as baboons and some species of macacus, the upper portion of the ear is slightly pointed and the margin is not at all folded inward; but if the margins were to be thus folded, a slight point would necessarily project inward toward the center and probably a little

Fig. 14.

Fig. 15.

outward from the plane of the ear, and this Darwin believes to be their origin in many cases.

The hundreds of these cases observed by me have always been found among the neurotic and degenerate. As according to Darwin, "when present it is developed at birth," it therefore must have formed when the ear was developing before the fourth month of fetal life. At birth, however, it is not as prominent as it will become when the ear has obtained its full development, although prominent enough to be recognized.

Prof. Meyer finds these tubercles more common among men than women. This is in accordance with the general rule in evolution that women are truer to the type than men. According to Meyer, it is due to

Fig. 16.

Fig. 17.

arrested development of the helix. In my experience, it is due to both excessive and arrest of development of tissues. I have observed not infrequently both conditions upon the helix of one ear. Tubercles may be located upon both ears or upon one. They also vary in number, one only, or sometimes three, even four have been noticed upon one ear.

The position of these tubercles depends entirely upon the shape of the helix. If the helix is turned toward the center of the ear, the tubercles will point in that direction; if at right angles, they will stand at right angles. If the ear lies perfectly flat without a helix, the tubercles will stand in direct line with

the flat ear and point backward or upward; whether they can be observed standing in front, at the side or back, will depend upon the position of the ear to the head. Some ears stand close to the head, others at an angle of ninety degrees. The location of the tubercle upon the helix varies in different individuals. The most frequent localities, however, are at the upper border of the middle third, as illustrated by Darwin; also at any point on the helix from its origin to its insertion; again, just above and below where the root of the antihelix crosses the root of the helix in the floor of the concha. I have also observed upon the left ear of a physician, which measured 3 inches in length, 1.50 in width, and stood almost at right angles, five tubercles in the upper third of the helix, due to arrest of development. Again, one may be located on the outer border looking outward, and the other on the inner border, looking inward. (Fig. 14.)

In one family coming under observation, the grandmother has three tubercles on left ear, one at the lower border of the middle third, one above and below on the root of the helix where the root of the antihelix crosses it in the concha. A daughter has one on the root of each helix, in the concha below the point where the root of the antihelix crosses it.

Fig. 18.

Fig. 19.

The oldest grandchild, eighteen years of age, has the same tubercles in same places as mother, and also half developed tubercles on the root of the helix above the point where the root of the antihelix crosses the helix. The second grandchild, a boy ten years of age, has a fully developed tubercle upon each root of helix below the point where the root of the antihelix crosses it, and a semi-developed tubercle on each helix just above that point. The third child (Fig. 15), a daughter one year of age, has four tubercles upon the left ear, two on the root of the helix, like grandma, one at lower border of middle third and one at lower border of lower third; on right ear two on root of helix like other side, and a semi-developed tubercle at the lower border of middle third.

It is impossible to illustrate the tubercles on the root of the helix, owing to the fact that they are situated inside the concha. Figures 13, 14 and 15 form an excellent illustration in the regicide, J. Wilkes Booth; one at the top, the other just below the middle of the helix. In Cesare Giovanni Santo, one at the top, the other at the lower third. In Patrick Eugene Joseph Prendergast, one at the locality pointed out by Darwin, the other at the lower third, near the lobule.

A young professional man, 29 years of age, is a marked neurotic; nervous temperament, suffers con-

tinually with indigestion; has insomnia; possesses a good mind, is a great reader; has full forehead, wears glasses for astigmatism; stigmata show arrest of development of the bones of the face and upper jaw; asymmetrical head; V-shaped arch; high vault; arrest of development of the lower jaw; possesses a long, narrow, handle-shaped ear, with arrest of the upper third and outer border of the helix, showing two distinct tubercles, one near the cranium, the other at the outer border as illustrated by Darwin.

A remarkable case is that of a degenerate whose left ear is 2.75 inches in length, 1.12 inches in width, possesses a tubercle at the lower border of the lower third, one at the middle of the upper third upon the border which looks downward toward the concha, and another just on the opposite side of the helix looking upward; the one looking downward is due to arrest of the edges of the helix, the other to excessive development of the outer border of the helix. From what has already been said it will be observed that these tubercles are in all stages of development from a minute point to the size of a large pea.

As stated at the outset, these tubercles are always observed among the neurotics and degenerate classes; they are due to arrest and excessive development of the helix. Sometimes there is a total arrest of the

Fig. 20.

Fig. 21.

helix, giving the ear the appearance of having been cut or burned away. When these tubercles are observed upon one or both ears, marked stigmata are also observed in other parts of the head, face, jaws and body. This is nicely demonstrated in the case of Prendergast.

In the ear illustrated by Darwin, are other points which demonstrate more clearly developmental neurosis, or degeneracy, than his tubercles. Since this ear was modeled by the celebrated sculptor, Woolner, it certainly must be very accurate. The first thing to be noticed is a general arrest of development. This ear is only about two-thirds the normal size and is unusually thick. (I have already observed that when the ear is unusually large, it is also very thin; when unusually small, it is very thick.) The lobule is arrested to such an extent that it is almost obliterated. The tragus and antitragus are almost entirely undeveloped, leaving the concha unprotected. The helix is rolled upon itself, the result of general arrest. The arrest of development of the rim of the helix at two different places and its development in a small sphere, produces the tubercle. From such a degenerate ear, one accustomed to the study of degeneracy could describe the deformities of the face of the individual possessing the ear, without difficulty.

Sometimes these tubercles grow to an immense size; thus, a business man, 35 years of age, possesses

an ear at right angles, with tubercles equal in size on each ear, as large as the end of the little finger; these are located at the upper middle third. The helix is entirely arrested.

Another singular case is that in which the helix of the left ear is excessively developed. At the extreme upper border there is a large tubercle at the inner edge, looking downward; and opposite, on the upper border looking upward, is another of the same size. Both of these are due to excessive development. Another case is that of a baby 14 months old. There are four tubercles on the left ear and none upon the right. One at the upper middle third, one at the ex-

Fig. 22.

Fig. 23.

treme top of the helix, both due to arrest of helix; and two on root of helix, in the concha, one anterior, the other posterior to the lower root of the antihelix, making a groove through which it is to pass.

The same exists in a brother and sister, mother and grandmother, and not in the father of the children. They are not, however, exactly alike in all generations, or in the same generation. This is, no doubt, due to malnutrition.

According to Frigerio, the exaggerated tubercle of Darwin is seen exceptionally in normal individuals, rarely in non-hereditary insane; with an average of 36.95 per cent. in hereditary cases and of 27 per cent.

Fig. 24.

Fig. 25.

in neuropathic degenerates, and not so marked in criminals.

I have taken statistics of tubercles of 1,000 so-called normal persons, with the following results: Right 414, left 446. In 100 noted men, right 33, left 29.

In an examination of 1,506 criminals, making 3,012 ears, these were found upon upper third, right 152, left 127; middle third, right 504, left 452; lower third, right 2, left 3; none, 1,772 ears.

ANTIHILIX.

Figures 16, 17 and 18 show arrested, medium and excessive development of the antihelix. My experience has been that where there is arrest of devel-

opment of the helix, the antihelix is excessively developed, and *vice versa*.

This would seem to be a natural compensation for the purpose of collecting the waves of sound. In apes the antihelix is rudimentary, thus showing that the helix is most usually developed. It is claimed by some scientists that the antihelix is also wanting in the most pronounced forms of insanity and in criminals. The following are the results of a most thorough investigation of criminals:

At Pontiac, arrested, 195; excessive, 271. At Elmira, arrested, 317; medium, 13; excessive, 722.

Fig. 26.

Fig. 27.

TRAGUS.

Figures 19, 20 and 21 show arrested, medium and excessive development of the tragus. The tragus and antitragus are most frequently malformed in the degenerate, though the helix and antihelix are more noticeable. One of the peculiar forms of degeneracy noticed in the tragus is a fissure dividing it into two distinct lobes. Sometimes it is very slight, making a long, double tragus with a slight depression in the center. Again, a little deeper, showing two tubercles, and again, two distinct organs.

In an examination of criminals at Pontiac: Arrested, 117; medium, 244; excessive, 124. At Elmira: Arrested, 534; medium, 70; excessive, 427.

Fig. 28.

Fig. 29.

ANTITRAGUS.

Figures 22, 23 and 24 show arrested, medium and excessive development of the antitragus. This organ is very frequently absent among the degenerate classes.

Examination of criminals at Pontiac shows: Arrested, 313; medium, 59; excessive, 76. At Elmira: Arrested, 693; medium, 21; excessive, 216.

LOBULE.

Figures 25, 26 and 27 illustrate long, medium and short lobule. Perhaps no part of the ear has received so much attention by scientists as the lobule. The long, handle-shaped ear (meaning, of course, the

long lobule) is a marked sign of degeneracy; the short ear, without lobule, also denoting arrest of development. Very little, therefore, is left for a normal lobule. So frequent do we find these conditions among normal individuals that, without other stigmata, either excessive or arrested lobule would be without significance. In extreme cases the lobule

will develop forward and frequently double upon itself, not unlike the ears in the lower forms of animal life.

Figures 28, 29 and 30 illustrate attached, medium and separated from the face. These conditions have also received considerable weight by scientists and neurologists, as demonstrating stigmata of insanity

STATISTICS OF DEGENERACY OF THE EAR.

												Auricle.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
												Length.						Width.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
												1.75		2.00		2.12		2.37		2.25		2.50		2.75		3.00		.84		.75		1.1		1.12		1.37		1.25		1.50		1.75		2.25																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
												Number.		Large.		Medium.		Small.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
Normal persons, right ears...												1,000	595	68	258																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Per cent...													59.5	6.8	25.3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Normal persons, left ears...												1,000	561	68	372																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Per cent...													56.1	6.8	37.2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Noted men, right ears...												100	62	5	33																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Per cent...													62	5	33																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Noted men, left ears...												100	56	5	44																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Per cent...													56		44																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Criminals...												1,506				1	33	9	9	295	930	196	33	2	2	100	275	172	824	115	3	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						</

Helix.									Antihelix.						Tubercles of Darwin.									Lobule.											
Development.			Form.			Root.			Development.			Upper 3d.		Middle 3d.		Lower 3d.		Attachment.			Shape.			Development.											
Arrest.	Medium.	Excessive.	Folded.	Medium.	Open.	Abnormal.	Normal.	Arrested.	Medium.	Excessive.	Right.	Left.	Right.	Left.	Right.	Left.	None.	Close.	Medium.	Separate.	Broad.	Narrow.	Long.	Arrested.	Medium.	Excessive.									
667	13	312	325		667	325	667	627	13	352	33		378		3		605	920	74	8	554	444	18	447	33	520									
66.7	1.3	31.2	32.5		66.7	32.5	66.7	62.7	1.3	35.2	3.3		37.8		3		60.5	92	7.4	8	55.4	44.4	1.3	44.7	3.3	52									
532	14	135	449		532	389	582	532	14	469							582	865	73	17	580	421	7	429	7	565									
53.2	1.4	43.5	44.9		53.2	33.9	58.2	53.2	1.4	46.9							59.2	86.5	9.2	1.7	58	42.1	7	42.9	7	56.5									
59		41	41		59	28	72	41		59			33				67	100			59	41		41	1	58									
59		41	41		59	28	72	41		59			33				67	100			59	41		41	1	58									
57		43	43		57	17	83	43		57							79	94	5	1	55	45		46		54									
57		43	43		57	17	83	43		57							79	94	5	1	55	45		46		54									
1007	107	380	475	48	983	440	1024	512	13	993	152	127	504	452	2	3	1008	1140	354	19	1065	465	68	736	37	711									
33.43	3.55	12.62	15.77	1.59	32.64	14.61	35.33	17.00	.43	32.97	5.05	4.22	16.73	15.01	3	.06	3	33.47	37.85	11.75	.63	35.36	15.44	2.26	24.44	1.23	22.61								
296		147	147		296	119	324	149		290	77	77	174	172	3		222	418	25		255	188	1	237		206									
33.41		16.59	16.59		33.41	13.43	36.57	16.82		32.73	8.69	8.69	19.64	19.41	3	34	25.06	47.18	2.82		28.78	21.22	.11	26.75		23.25									

and criminality. Like the length of the ear, no more stress can be placed upon these conditions than upon the length, without other stigmata.

In an examination of criminals, the following results are obtained:

Attachment.			LOBULE.		Shape.			Development.		
Close.	Medium.	Separated.	Broad.	Narrow.	Long.	Ar't.	Med.	Exc'v.		
PONTIAC.										
309	163	11	334	104	33	255	12	198		
ELMIRA.										
881	191	19	671	361	35	481	25	513

Fig. 30.

Fig. 31.

Fig. 32.

Figures 31, 32 and 33 illustrate ears of illegitimate children at birth. Owing to their age their mentality is not fully determinable, but numbers 32 and 33 have every appearance of idiocy.

Figures 34 and 35 show the congenital elephantine ear in monstrosities. For the purposes of comparison, the results of my investigations are given in full.

The tables show that the various parts of the ear are often not in harmony with each other.

In the study of aural relations to degeneracy, it is customary to speak of large, medium, small or handle-shaped ears without setting a definite standard for

comparison. Having ascertained the average size length 2.50 and width 1.25 inches, of a given number of ears of normal persons, it may be stated that an ear which measures below 2 inches can be considered a very short ear, between 2 and 2.25 inches short, between 2.25 and 2.75 inches medium, between 2.75 and 3 inches long, and above 3 inches very long. In regard to width an ear below 1 inch is narrow, between 1 and 1.50 medium, and above 1.50 wide.

Fig. 33.

Fig. 34.

Fig. 35.

There is no means at hand of knowing what changes occur in the length of the ear at different ages over 50 years in normal persons. There seems, however, to be a gradual increase in length and width. I have noticed, casually, the ears of a number of persons on the street, and should judge that a large percentage of people possessed ears from 3 to 3.36 inches in length and from 1.50 upward in width. Comparison of the length and width of the ears of paupers, criminals and the insane, indicates that degenerates possessed much the largest and widest ears.

The statistics show that the ear is not in harmony

in its relation to its various parts, and that so-called normal individuals have just as deformed ears as criminals and the insane.

On comparison of the figures of the table it will be seen that the percentage of deformities of criminals and insane is nearly the same, and taken together is about one-half as great as the normal.

There is a somewhat prevalent theory that persons whose lobule is attached its entire length have insane tendencies. The foregoing data do not corroborate this, since in an examination of 1,000 normal persons 92 per cent. are found to have attached lobules, while among the insane only 47 per cent. was found. It follows that an attached lobule is more common among the normal than among the insane. Among criminals is found 37 per cent., and in noted men 9 per cent. for left ears, 10 per cent. for right ears.

The table shows that the tubercle of Darwin is not always found at the upper part of the middle third of the helix (Fig. 13). Tubercles, practically identical, may be found at any point on its border. These tubercles are obviously similar in appearance and due to the same causes that produce the tubercle of Darwin (arrest and excessive development of the helix). Hence, the theory advanced by Darwin that this tubercle is a rudimentary point on the ear is not sustained. In normal persons 3 per cent. of the tubercles of Darwin occur on the upper third, 37 per cent. middle third, 3 per cent. lower third, in right ears, while in criminals 5 per cent. occur on the upper third, 16 per cent. middle third, 3 per cent. lower third, right ears, and in the insane 8 per cent. upper third, 19 per cent. middle third, and 34 per cent. lower third, right ears. They are obviously more numerous upon the middle third of the helix. There is a larger percentage among normal individuals than with the degenerate class.

The left ear, however, seems to be least affected in either the normal or the degenerate. That these tubercles denote degeneracy may be admitted, but it does not seem probable that it comes under the type of degeneracy to which Darwin ascribes it.

The ear at 90 degrees is said to be a marked sign of degeneracy. The figures, however, do not bear out this claim. In normal persons this occurred in 3 per cent. right and left ears, 4 per cent. and 2 per cent. right and left ears of noted men, 10 per cent. right and left in criminals, and 3 per cent. in the insane.

In the relation of the ears to each other there is found a larger percentage of right ears higher than left ears; in criminals 17 per cent. right were higher than left and only 3 per cent. left higher than right, while in the insane 2 per cent. right were higher than left and 4 per cent. left higher than right. In the 1,000 normal individuals there were more deformities upon the right ear than upon the left. This is also true of the 100 noted men. At the time of the examination of criminals and insane these were not classified, as in the normal persons and noted men, except as to the tubercles of Darwin and the interrelations of the ears.

That the external ear as an organ is tending to disappear is shown by the feeble power of its muscles which, except in cases of reversion or special training, are inactive. Changes in the ear by themselves are merely suggestive and not demonstrative of degeneracy, except in accordance with the general rule that accumulation of defect constitutes strong evidence.

SOCIETY PROCEEDINGS.

Chicago Medico-Legal Society.

Regular Meeting, held in Handel Hall, Jan. 11, 1896.

THE PRESIDENT, DR. E. J. DOERING, in the Chair.

The minutes of the previous meeting were read and approved.

DR. F. E. DANIEL, of Austin, Texas, contributed a paper entitled

A PLEA FOR REFORM IN CRIMINAL JURISPRUDENCE, which was read by the Secretary, DR. JOHN RIDLON, in the absence of the author.

He referred, at the outset, to a paper which he presented on "Castration of Sexual Perverts," read at the International Medico-Legal Congress, held in Chicago in 1893, in which he advocated castration of sexual pervers, and especially as applied to rapists, as being more humane, more rational, and more effective as a deterrent than hanging, and more in accord with science, progress and an enlightened civilization; not as a penalty, but as a therapeutic measure in some cases, and prophylactic in all, for the present protection of society, and as a guarantee against a like progeny to curse future generations. Incidentally the author advocated the castration of certain criminals also in this paper, and for the same reasons. In the light of the rapid and alarming increase of crime and criminals in this country, it is evident not only that there is something radically wrong with our system in dealing with crime and criminals; that our penal methods are a dismal failure, but that reform has become an imperative, pressing and immediate necessity. According to the United States census for 1890, there were in 1850, 6,737 prisoners in the United States, or one to every 3,442 of population. In 1890 there were 82,329 prisoners, or one to every 757 of population. The end and object of all penal enactments are supposed to be the protection of society against crime and criminals, and it is unfortunate that no measures are provided as a protection against this increase, principally by heredity. Nothing could testify more emphatically to the inadequacy of our system than the figures just given, nor appeal more forcibly for reform. The thinking members of the medical profession are being rapidly converted to the belief that crime is a disease; that criminals are sick persons, and that their condition calls for a more enlightened method of management. They know that many, if not most of them, are subjects of hereditary transmissions of vicious temperament, and all of them the victims of vicious environment.

The classification of criminals can only be done by medical men. When the character of the crime has been determined by the court, it would seem to be in accord with the requirements of the case and the dictates of an enlightened humanity, that there should be a commission of medical men to diagnose the criminal and prescribe the course which is best calculated to meet the demands. If it be one of the curable class, the treatment, consisting of restraint, discipline, hygiene, education, environment and healthy labor should be such as to induce a determination never to offend again. Pride and self-respect should be fostered, for they are the highest incentives in life to good behavior. The term of imprisonment should depend upon the progress made in reformation, and on good behavior; the culprit made to realize that when he gives evidence of fitness to be trusted with his liberty it will be restored to him. The incurable—the born criminal of Lombroso—should be dealt with as a permanent enemy to society, and the first aim in his case, after sequestration, should be precautions against a progeny. When a man has been diagnosed as a natural criminal, twice convicted of any felony, along with the forfeiture of liberty for life and all other rights, he certainly should be deprived of the right to inflict upon the next generation a progeny sure to be like him. Why not? Can any one give a single reason why this right should be respected when all others are taken away? The author thinks not. The criminal insane should also be embraced in this category. Were it understood that being insane at the time of the crime, or becoming so after, makes no difference, and that castration is to follow, sane or insane, there would be fewer pleas of insanity, and thus would be stopped one loophole of escape by which