a knowledge of the constitution of the ether which we do not possess. We now assume that the ether is not set in motion by an electro-magnetic wave. If we do not make this assumption, we must introduce into our equation quantities representing the components of the velocity of the ether, and unless we know the constitution of the ether, so as to be able to deduce these velocities from the forces acting on it, there will be in the equations of the electro-magnetic field more unknown quantities than we have equations to determine. It is, therefore, a very essential point in electro-magnetic theory to investigate whether or not there is any motion of the ether in a varying electro-magnetic field. We have at the Cavendish Laboratory, using Prof. Lodge's arrangement of interference fringes, made some experiments to see if we could detect any movement of the ether in the neighborhood of an electric vibrator, using the spark which starts the vibrations as the source of light. The movement of the ether, if it exists, will be oscillatory, and with an undamped vibrator the average velocity would be zero; we used, therefore, a heavily damped vibrator, with which the average velocity might be expected to be finite. The experiments are not complete, but so far the results are entirely negative. We also tried by the same method to see if we could detect any movement of the ether in the neighborhood of a vacuum tube emitting Roentgen rays, but could not find any trace of such a movement. Prof. Threlfall, who independently tried the same experiment, has, I believe, arrived at the same conclusion.

Unless the ether is immovable under the mechanical forces in a varying electro-magnetic field, there are a multitude of phenomena awaiting discovery. If the ether does move, then the velocity of transmission of electrical vibrations, and therefore of light, will be affected by a steady magnetic field. Such a field, even if containing nothing but ether, will behave toward light like a crystal, and the velocity of propagation will depend upon the direction of the rays. A similar result would also hold in a steady electric field. We may hope that experiments on these and similar points may throw some light on the properties of that medium which is universal, which plays so large a part in our explanation of physical phenomena, and of which we know so little. J. J. THOMSON.

THE ACTION OF LIGHT UPON PERFUMES.

In a preceding article we described a process devised by Mr. E. Mesnard for comparing the intensity of perfumes. The same investigator has endeavored to produce an apparatus that should be less delicate and more easily transportable, and that should give just as accurate results. The model that he has decided upon is very ingenious. We shall give a brief description of it, as seen by us in operation at M. Gaston Bonnier's botanical laboratory.

In an oblong box provided with a hinged cover (Fig.

FIG. 1.—APPARATUS FOR COMPARING ODORS.

1) there are arranged two parallel cylinders of fine wire gauze, each mounted upon a horizontal axis. Each cylinder is maneuvered from the exterior by means of a winch placed at one of the extremities of the axis. The other extremity of this axis carries a graduated disk which moves along a fixed rule that is likewise graduated. It is possible in this way to estimate the number of revolutions and fractions thereof communicated to each of the cylinders.

It is in the oblong box that the odors are mixed. To this effect there is placed externally a frame divided into two compartments, each of which contains a small grooved pulley over which passes a thread. This thread, thoroughly saturated with the essential oil, is dried by its passage through a wad of cotton, enters the box through a very small orifice and afterward winds around one of the cylinders.

By this arrangement, it is therefore possible, at one's will, to cause a certain length of thread impregnated with essential oil to wind up in the box, or else to cause the thread that was first inclosed in the frame to re-enter the latter in order to become saturated anew. It will be readily understood that upon winding a greater or less length of thread around the cylinder, there is introduced into the box a quantity of essential oil that is proportional to the length of the thread wound.

The cover is provided with an orifice surmounted by a cone that has exactly the form of the nose and permits of obtaining a smell of the interior when a valve is opened through a pressure upon a small button properly arranged. If it is a question of comparing two essential oils, one of the threads is impregnated with the unknown oil and the other with spirit of turpentine taken as a standard, and an endeavor is made to ascertain what length of this latter thread must be introduced into the box in order to produce upon the olfactory membrane an odoriferous intensity equivalent to that which is produced by a given length of thread impregnated with the unknown essential oil.

This therefore amounts, upon the whole, to the same thing as measuring the intensity of a perfume with a meter measure. The model just described is especially arranged for the comparison of the intensity of liquid perfumes. If it is desired to experiment upon a

bouquet of flowers, the latter is introduced into a cylinder closed by rubber cloth and connected with the apparatus by properly arranged tubes.

This arrangement is no longer applicable when it is a question of but a single flower, whose odor is necessarily too feeble to allow of its being led to a distance by tubes. The apparatus is then modified (Fig. 2).

The box, which is carried by a small support and stands vertically, is open at the bottom, in order to allow the flowering top of the plant to enter it; but it may afterward be inclosed in rubber cloth tied around the stalk with a cord. The top of the box is arranged as before. It contains an orifice provided with a valve and a piece to which the nose is applied when it is desired to obtain a smell of the interior. In this way the nose is very near the flower.

At the side of the box there is a frame that carries the mechanism designed to measure the quantity of spirit of turpentine necessary to neutralize the intensity of the flower's perfume.

In both models of the apparatus the mixing of the odors is effected by means of rubber bulbs. The cleaning of the tubes is effected by means of a current of hot air. Thanks to these apparatus, Mr. Mesnard has been able to perform a large number of experiments, of which we shall give only the results.

It is light, and not oxygen, as is commonly thought, that is the principal cause of the destruction of odoriferous substances; but these two agents seem sometimes to combine their effects in such a way as to produce a maximum action.

The action of light manifests itself in two different ways. On the one hand, it acts as a chemical power capable of furnishing energy to all the transformations through which odoriferous substances pass from their elaboration to their total resinification. On another hand, it exerts a mechanical action that permits of explaining the method of periodical disengagement of the perfume of flowers.

The intensity of the perfume disengaged by a plant depends upon the state of equilibrium that is established at every hour of the day between the pressure of the water in the cells (which tends to expel to the exterior the essential oils contained in the epidermis) and the action of light, which combats such turgescence. Just as a simple watering suffices to increase the turgescence

FIG. 2.—APPARATUS FOR MEASURING THE INTENSITY OF PERFUMES.

of the cells, so the interposition of a simple glass screen, the presence of a cloud or the advent of cloudy or rainy weather suffice to greatly attenuate the effect produced by light. In both cases the intensity of the perfume increases.

In reality, it is the irritability of the protoplasm that is the primordial cause of the variation in intensity of the perfume of flowers, and experiment proves that a sudden insolation or simple contact suffices to provoke such irritability, and consequently to bring about a notable variation in the state of equilibrium of the plant and an almost sudden increase of the intensity of the perfume.

The alternations of day and night determine maximum and minimum values of the intensity of perfumes, which are often modified, it is true, by the changeableness of the weather, but which nevertheless bring about a true periodicity in the disengagement of odors. There are some plants that are more sensitive than others, certain orchids, for example, in which such periodicity occurs in a much more marked manner.—*La Nature*.

THE CAUSES OF COLOR.

By J. J. STEWART, B.A. Cantab., B.Sc. Lond., in Knowledge.

THE beauty of the objects of nature around us depends so much on the varieties of color which they exhibit that it becomes an interesting question, What is the cause of these differences? We have only to think of the blue sky, the green foliage, and the various splendors of flowers and fruits to realize to how great a degree the pleasure we experience in viewing a country scene is due to the many-hued surfaces of natural objects. The pleasurable feelings aroused by the sight of the deep blue of the sea contrasting with the bright red color of the sandstone cliffs, or the effect of the sunshine when it lights up the splendid purple of the heather on the slopes of the hills, can be forgotten by none who have experienced them; and the striking effect of Eastern scenes is owing largely to the brilliance of the tints which meet the eye. It is needless

to enlarge on the part played by color in this world of ours; the difference between a world of color and one in which surrounding objects were only distinguished by different shades of gray can be realized by all.

When we come to inquire how the red color of a rose is produced, and why it differs in appearance from a blue flower, we must consider what happens to the light which falls upon the petals of the rose. We see the flower because the light from the sun is reflected from it, but something has happened to the light before it reaches our eyes: the light we receive differs from that which fell upon the flower. The rays from the sun penetrate to a certain extent into the substance of the flower, and most of them are reflected from particles beneath the surface. Now the cells making up the petals of a rose contain a fluid which has the power of absorbing certain of the rays of light, and the light entering the eye after penetrating a short distance below the surface of the petals and coming back has passed through this fluid, and in its course some of the rays of the sunlight have been abstracted from it. Thus the light reaching us is lacking in certain of the constituents of white light—that is, it is colored. White light may be considered as made up of the three fundamental colors, red, green and violet, blended together. The light which has passed through a certain extent of the substance of the rose petals has been deprived of its green and violet portions, and thus the red rays alone reach our eyes. All substances which possess color exercise this power of sifting the rays of light. Light falling on the leaves of the rose bush passes through their superficial layers and is reflected from below the surface; thus it has to traverse certain particles which take from it the red and violet rays and allow the green to pass. The green rays which escape absorption are the only ones which reach our eyes, and we therefore call the leaves green.

When white light passes through a prism it is found that in the band of color or spectrum produced, the blue and violet rays are the most bent out of their original course; they are the most refrangible. The red rays are the least bent. It is found, also, that the vibrations of the ether filling space which produce waves of light are quickest in violet light and slowest in red light. Some substances absorb the quick vibrations more readily and thus appear reddish in color. Others absorb the slower vibrations, allowing the others to pass through, and therefore have a green or blue color.

When a substance is heated its particles are thrown into a state of rapid motion, and soon set up a motion in the ether which, when the vibrations are of a certain rapidity, produces light. If a ball of iron is heated, it first of all gives out radiation consisting of dark rays, which have the effect of producing heat. As its particles become hotter and hotter, and thus vibrate more rapidly, the radiation begins to affect our eyes and the ball glows with a dull red heat. As vibrations of greater and greater rapidity take place owing to the further heating, the ball appears bright yellow, and finally white, when vibrations of all the different rapidities which affect our eyes are given out. In this case it is the motion of the particles or molecules of the heated body which is imparted to the ether, and so produces the effect we call light. Now those molecules which are able to execute certain vibrations and give them to the ether absorb these same vibrations from the ether—that is, absorb certain rays of light. This effect is best seen in the case of gases, and the phenomenon is analogous to that which occurs with sound. A tuning fork or stretched wire which can give a certain note when it is struck is able to take up and absorb the note from the vibrating air around it when that note is sounded in its neighborhood.

It may be noticed that the light from the electric arc when seen near at hand has a distinctly bluish color; but this same light when viewed from a distance appears yellowish, as certain of its rays have been absorbed by the water vapor in the air on its passage to the eye. For a similar reason the sun is now considered to be a blue star; his light, which would appear intensely white, and rich in blue rays especially, if it could be seen from beyond our atmosphere, appears yellowish after it has passed through that atmosphere and has lost some of its most refrangible constituent rays.

The effect of absorption in producing color is seen from the fact that powdered bodies generally appear white. This is accounted for when we consider that a powder consists of particles arranged at all angles, so that the light falling upon it meets various surfaces and is mostly reflected before it has passed below the surface. Thus the white light reaching it is not deprived of some of its constituents by selective absorption, as it would be if it penetrated the substance and was then reflected. In this way powdered red glass appears white. For a similar reason the froth of colored liquids, such as brown ale, appears pure white. The light is reflected from the surface of numerous small bubbles, and does not pass much through the liquid itself. Thus also a cloud is very opaque to light, the light falling on it being reflected at the surface of the numerous globules of water. To this is due the brilliantly white appearance of large fleecy clouds in bright sunshine.

Some substances absorb equally all the rays of light. Such substances, of which soot is an example, appear black. The reason why a flower like a white lily appears white is that the fluid contained in its cells does not absorb one sort of rays more than another, but allows all to pass with comparative freedom. White light then reflected from its surface, or from a little below, is not deprived of any of its constituents, but remains white.

The effect of reflection from internal surfaces, accompanied by absorption in producing color, can be seen by pouring a colored liquid, carefully freed from floating particles, into a white porcelain basin. Light is reflected from the sides of the basin, passes through the liquid, and its color is seen. If now the sides of the basin be covered with some black substance, no light will be reflected from them and the liquid will appear black; no light comes to the eye from the interior, and the surface of the liquid reflects all the rays equally. If next we place in the black looking liquid a white powder like chalk, its color is at once restored, light being now reflected from the interior at the surfaces of the chalk particles.

From the above considerations we can understand

to what causes the color produced on mixing pigments is due. A mixture of blue and yellow paints has a green color because that is the only color transmitted by both pigments. The blue paint absorbs the red, orange and yellow rays, allowing the others to pass through it; the yellow paint absorbs the blue, indigo and violet. Thus green rays alone are permitted to pass through both, and the result is that the mixture appears of that color.

Some substances appear of one color when viewed by reflected light, and another when seen by transmitted light. Thus the light reflected by gold is yellow; but a leaf of gold made so thin that light can pass through it appears of a green color. This appearance of different colors on reflection and transmission is also seen with many of the aniline dyes. The color of the light due to reflection is then made up of those rays which are not admitted at all, but sent back at the surface, together with that light which has been reflected from a certain depth below the surface, and has thus lost some of its constituent rays by internal absorption. The light to which the color is due when the substance is viewed by light which has passed through it, is that which has been deprived of some rays by reflection at the first surface, and again of others by absorption in passing through. Hence the difference of color when viewed in the two different ways.

Variations in color perception no doubt depend on varying sensations in our own eyes, as well as on changes in the light itself. Some curious experiments have been made with a view to testing our different sensations as to color. It has often been noticed that a bright scarlet uniform will appear perfectly white in a good photographic dark room with ruby glass windows. With regard to such effects, Herr H. W. Vogel described recently in Berlin some experiments he had made. He used oil lamps and fitted on to them pure red, green and blue color screens. It was found that when the white light was entirely shut out, no sense of color was perceptible to the observers, and objects in the room appeared of various shades of black and white. He found that when a set of colors was lit up by red light, the red pigments appeared white or gray, and this changed at once into yellow, not into red, when blue was added to the light under which they were viewed. Thus a color was perceived which did not exist in either of the sources of light used. The color sensation produced by a source of light also depends partly on the intensity of the illumination. From these and similar experiments, Herr Vogel comes to the conclusion that our opinion as to the color of a pigment depends upon our perception of the absence of certain constituents from the light reflected from it. Thus a surface which has a red color is only perceived as red by us when light of other colors shines upon it, and we observe its incapacity for reflecting these colors.

When a solution of quinine is viewed in sunlight, a remarkable blue shimmer is noticed extending for a short distance beyond the surface at which the light enters. A similar effect is noticed with many other substances, the color being different in different cases. The phenomenon is known as fluorescence, as it is well observed in the mineral fluor spar, and is due to the fact that light is absorbed by the substance and is again given out as light of a different color. For instance, rays of high refrangibility toward the blue end of the spectrum may be taken in and given out as yellow rays of lower refrangibility. In the case of quinine, invisible rays beyond the violet are absorbed and blue or violet rays are emitted by the solution. In all cases of fluorescence a degradation of the rays takes place; those given out are of lower refrangibility than those which disappear on absorption. By painting a screen with a solution of sulphate of quinine, the spectrum beyond the violet can be made visible, as those vibrations which are too rapid to affect our eyes are changed into others of lower refrangibility, which can be perceived when they fall on the retina. The curious blue color of the solution of quinine extends only a short distance into the liquid, because those rays which are capable of producing it are soon absorbed, and the light which passes onward through the fluid is destitute of such rays. The color produced in cases of fluorescence has a different origin from that of the ordinary surface color of substances, for the rays absorbed do not disappear as light, but their place is taken by other rays of a different sort.

The nature of the vibrations which constitute light still remains mysterious, but great advances have been made in recent years in our acquaintance with these and allied phenomena, and we may hope for still greater accessions to our knowledge in the not distant future.

SOME INTERESTING SPIDERS.

By Prof. FRANZ MÜLLER.

WHO does not know the common shepherd spider that moves about in such ghostly fashion on its long, slender, stiltlike legs? Even as children we were entertained by the grotesque movements of the perfectly harmless creature. This phalangium, although related to spiders and scorpions, has no poisonous weapons, but in place of the poison fangs of real spiders, it is provided with downwardly bent scissorlike palpi, by means of which it catches little insects and carries them to its mouth. It is so well known that it needs no description, but an enumeration of the characteristics which distinguish it from other spiders may be of interest: These are the filiform maxillary palpi terminating in simple hooks, the segmentation of its pea-shaped abdomen, and the small number of its eyes—for it has only two simple eyes. These spiders prefer to spend the day grouped together in dark corners and nooks, and if one of them is touched lightly, it begins to rock or swing its body. This motion is imitated by those near it and soon the whole company are making one another comical bows. Gradually they become quiet again. If touched more roughly or caught, they try to escape, even if obliged to leave behind one or more of their legs. As every one knows, these legs that have been torn off keep up a fluttering motion for hours. This is the result of the decentralization of the nervous system found in all articulated animals, which causes single dismembered portions of the body to show signs of life for a long time; but, in all probability, there is no conscious feeling. One may, for instance, see wasps, May bugs, and other insects running around

for days after having lost the whole abdomen, with the viscera, separate legs, wings or other parts of the body of which they have been deprived by being stepped on by a careless pedestrian. Our long-legged phalangium seems not to experience the least pain upon the amputation of one or more legs, and those lost members grow again after the next shedding of the skin, as those of the crab do under similar circumstances. The whitish eggs are laid by the female in damp places, where they can easily be hidden in hollows or cracks.

The swinging or swaying motion adopted by the phalangium when slightly disturbed is not peculiar to it, but is also characteristic of certain other spiders. If a cross spider is approached in warm weather, when in the center of its beautiful wheellike web, or if the web is moved slightly, it begins to swing so energetically that it seems to be a shadow in a rapidly vibrating web. This occurrence can easily be observed in summer in a sunny grove of young trees, where hundreds of these beautiful webs are spread. The object the creature has in making these strange movements it is not difficult to guess, for it must be the same as that of silkworms and other creatures which, when a sudden danger threatens, draw in their heads or take a defensive attitude, which is indicated by the forms and colors they assume. When a cat crouches her back upon the approach of a dog, bristling up her fur and showing her teeth and sharp claws, she does it for the same reason, viz., to frighten the assailant.

A certain spider has been called the "trembling" spider because of this strange motion which it imparts to its dwelling. Several specimens are shown on a somewhat enlarged scale in the accompanying engraving,

after day, and often at night also. Rooms and entries of country dwellings, cellars, garrets, and stables are favorite places of abode for these spiders. Those I watched were in a room which had been used for the storage of furniture, but was temporarily arranged for a sleeping room. They could always be found in the angle formed by the wall and the ceiling, or on the latter, but never far from the wall. Like the shepherd spider, they remain quiet and motionless, apparently sleeping, during the day, and at night they cannot wander very far, for in the morning they are always to be found in the same place. The reason for this is that their nets are there, stretched between rough places in the wall. They are scarcely visible, but upon looking closely, their presence is betrayed by the little flies and particles of dust which have been caught in them. If one of these spiders is touched slightly, it immediately begins its droll swaying motion, and then it can easily be seen that it is holding onto the threads by its long, slender legs. Larger flies are never seen in their webs, and if a live struggling fly is held in the web, the spider is frightened by the violent motion and takes flight. From all that I have been able to observe, the pholcus seems to live entirely on small bipennates which it, like the shepherd spider, catches by spreading its long legs and running over it. One night I saw many tiny flies run along the wall from the open window until some of them reached the domain of my spider. By making a sudden movement to that side of the web where the motion of the threads betrayed the presence of a fly, the spider caught his victim as quick as lightning and sucked its life out.

The swaying motion of one of these spiders is accom-

PHALANGIUM AND PHOLCUS.

Above at the left is a pholcus carrying its cocoon; at the right the cocoon is suspended. Below at the right a phalangium, at the left a white field showing the position of the eyes of the pholcus.

and as their habits of life are interesting, which perhaps some of our readers may have had as favorable opportunities of observing as I have, we will now study them more closely. This "trembling" spider is also called the "squinting" spider (pholcus, from the Greek pholcos, or squinting), which name was given it on account of the peculiar shape and position of its eight eyes (see lower part of the engraving). It has three large eyes grouped on each side, which differ from those of other spiders in being oval and milky white, with black edges, which gives the appearance of "squinting." The two median eyes are smaller and solid black. The creature is slender and the cephalo-thorax, which carries its long, delicate, hairy legs, seems almost round when looked down upon; its abdomen is rather long and almost cylinder shaped, and its front legs are nearly six times as long as its entire body. Its color is generally a grayish brown.

In regard to the geographical distribution of these spiders, they are found in Switzerland, the Netherlands, France, Italy, Sweden, Hungary, and in some localities in upper and lower Austria. Strangely enough, it and all other spiders of the same family (Phocidae) are very numerous in East India, and many students have come to the conclusion that this spider came to us from that country, not, it is true, over the land, but across the sea.

The habits of animals of all kinds can best be learned by watching them when free, for in captivity they generally change their ways and do not live as long as if they had been left in their natural state. This seems to be particularly true in regard to spiders. I once had an opportunity, during a stay of eight weeks at Mondsee, of watching nine of these interesting little creatures day

plished by placing its long legs on separate threads of the web or on rough places in the wall and moving its body in a circular fashion, and on account of the elasticity of its feet and of the threads of the web, the motion finally becomes so rapid that the body of the spider can scarcely be seen. This performance is most interesting, and a spider never tires of repeating it, whenever disturbed by a slight movement; it takes flight only when roughly disturbed.

Both of the spiders which we have been describing practice another ruse, which seems to be quite common to many other creatures higher and lower—that is, feigning death. Once I came upon a pholcus in a vestibule; upon being touched it began to sway, but stopped suddenly as if it had thought of something better, stretched out its legs and lay as if dead. When I wanted to take it up carefully, it freed itself by a sudden jerk and hurried off, leaving three legs behind it. The next day I found it in the same place, and, strange to say, it swung gayly in a circle with its five feet, as if it had suffered no inconvenience. Unfortunately, I could never know whether the lost legs grew again, for, greatly to my regret, it soon vanished forever.

Another characteristic which adds to the interest of watching these spiders is the special care manifested by the female for her offspring. It is well known that spiders place their eggs in a cocoon. In winter one often sees the egg sacs of the cross spider, about as large as a hazel nut, hanging on the wall, in the windows, and in similar places, and in summer wolf spiders can be seen in the fields, in woods, everywhere, running about with egg sacs fastened under the thorax by slender threads. If relieved of their precious burdens, they