he wished to say a word in favor of Bier's hyperemic treatment. He recalled the case of a young man of nineteen who about ten years ago had one testis extirpated for tuberculosis; the second organ subsequently became involved, and the patient begged to retain it, if possible. Bier's hyperemic method of treatment was instituted and faithfully carried out for a long time, and now he was cured so far as the tuberculosis was concerned, although the epididymis was still slightly enlarged. Quite a number of such cases were on record. This case was shown before this society about six to eight years ago. The treatment, which could be carried out by the patients themselves, was indicated in those cases where the one remaining testicle became affected by tuberculosis.

KIDNEY BOARD AND ARM GALLOWES.

DR. F. TILDEN BROWN demonstrated a later model in thin steel of a huge hinge, which was shown a year ago when made as a folding wooden board. The purpose of the apparatus is that of making a good surgical exposure of the ileo-costal space in kidney operations, when placed under the opposite region, where its degree of elevation is controlled by crank. In operations on gall-ducts and bladder also, its easy introduction under the dorsal spine makes it useful.

Dr. Brown also showed an adjustable gallows for slinging the superimposed arm when the patient is on the side. Relieving the chest of the shoulder and arm weight, besides favoring comfortable anesthetization, it serves as a prop to prevent the patient's thorax and trunk twisting forward.