mained, but there was no constipation, the bowels being opened forty-eight hours before operation. On admission, and for several days before, the patient suffered intense constitutional depression, and became feverish and very exhausted, but there was no vomiting, nor did he complain of much pain. When taken into the hospital, he was in a state of extreme prostration and collapse. The right scrotum was enlarged, swollen, tender, and of a dusky red hue.

Chloroform was administered and the sac cut down upon and opened. The tissues of the scrotum were found thickened and infiltrated; the hernial sac was distended with about half a pint of turbid, fetid serum, and large masses of recent lymph. The neck of the sac was quite free, and no constriction could be felt. At the upper part of the sac was a knuckle of intestine, which contained a mass of fæces of almost stony hardness, but which could be indented by firm pressure with the finger-nail. At the lower part of the knuckle was a small sloughy patch of about the size of a sixpenny piece, and from this point the intestine was incised for about an inch and a half, and then the hard fæcal mass (of about the size of two walnuts end to end) was turned out. After this about a pint of soft liquid fæces escaped. The bowel was then stitched to the margins of the scrotal wound, and an artificial anus thus established, all the fæces escaping through it.

On the day after the operation the patient complained of slight pain in the abdomen, but this was soon relieved by the application of poultices to the abdominal walls, and by the administration of opium in suppositories. He was ordered liquid diet and some stimulants. On the second night after the operation his temperature was 101.5° Fahr.; pulse about 100. Fæces passed through the artificial anus without pain, only flatus passing by the rectum.

Since then he has gradually improved, never having had a bad symptom, and is now convalescent, the greater part of the fæces passing through the natural anus.

KING'S COLLEGE HOSPITAL.

TUMOUR OF TIBIA; AMPUTATION THROUGH THE KNEE-JOINT.

(Under the care of Sir WILLIAM FERGUSSON.)

A. B—, a delicate-looking man, aged thirty, had been under observation with a tumour of the right tibia for the last seven months. The growth commenced between eight and nine months ago as a swelling from the front of the tibia; it gradually increased in size for several months, but then took on more rapid growth, the limb increasing an inch and a half in circumference during the last three months, until it formed a large mass of irregular consistence, extending downwards from about two inches below the head of the tibia, and taking in about seven inches of that bone. It appeared also to embrace the fibula, which at first was quite free. The greatest circumferential measurement was twenty-two inches. The skin was distended, but of normal colour, the veins being only slightly increased in number and size. The operation was performed in the following manner:—

Chloroform being administered, the artery was compressed against the pubis by means of the thumb. An oblique incision was then made through the skin from above downwards and backwards for about two inches on each side of the limb opposite the upper part of the tumour; the lower ends of the incisions were then united by a transverse cut above and below. The anterior skin flap thus made was then dissected back, the limb flexed, and the joint opened, the ligaments divided, and the knife carried behind the joint, cutting outwards and downwards to form the short posterior muscular flap. A few vessels were ligatured with whipcord, and the flaps brought together by silk sutures, and supported by means of pads of wet lint covered with oiled silk and a bandage.

Sir William remarked that when the tumour was first seen it was so small that no opinion could be given as to its nature, except the opinion founded on the experience that tumours of the tibia, not venereal, will probably increase in size. It would be impossible to say whether it is benign or malignant; the first impression is that the tumour is malignant. From an early period it was evident that nothing short of amputation would be of any value. If the

tumour were in the fibula, the whole or a portion of that bone might be removed; but not so in the tibia, and especially in a case like this, where the growth affected the whole thickness of the shaft, and probably began from its interior. Amputation through the knee-joint seemed to be a justifiable operation in this case. This operation, which was formerly rare, has now become more frequent, owing largely to the advocacy of Mr. Carden, of Worcester. Whether it is a good operation remains to be seen; but Sir William thinks it is. Further, this would seem to be a good case in which to employ it; for the disease is entirely limited to the tibia, and if the growth be not positively malignant it will not recur. But it does not necessarily follow that if there be a tumour of a bone, the whole of the bone must be removed, as some surgeons think, and Sir William has again and again protested against such extreme measures. The anterior flap was sufficiently long to give a good covering to the end of the femur, and did not necessitate the removal of the patella, which remained *in situ*, so that the part of the joint opposite and above that bone was not exposed. It is a moot point whether the patella should remain or not. Should the cartilage be removed? In this case it was not, and this is perhaps the best practice when it is healthy; for then there is little fear of absorption of septic matters from the wound, as may happen when the cancellous tissue is exposed. The stump, as far as could be judged, was a good one. But the operating-table is not the place from which to judge a stump; it is when the limb is used, three, four, or six months after, that we must form our opinion.

WESTMINSTER HOSPITAL.

A CASE OF CONGENITAL DISLOCATION OF THE PATELLA OUTWARDS.

(Under the care of Mr. HOLTHOUSE.)

DISPLACEMENT of the patella outwards is the most common of those to which this bone is liable, and may result from blows, or from twists of the limb, or from muscular action. Some persons seem to be particularly prone to this accident, and it has been observed to occur most frequently in those who are knock-kneed. The late Mr. Skey had seen two cases which occurred spontaneously during sleep, but, as far as we know, a case of congenital dislocation has not hitherto been recorded.

The subject in this case was a little boy, aged seven, who was admitted into the hospital on the 7th of May, with fracture of the left thigh at the junction of the middle and lower third, caused by getting the limb between the spokes of a cart-wheel while in motion. Next day, after the fracture had been put up, Mr. Holthouse observed while measuring the limb that the patella of the uninjured limb occupied an abnormal position, being partly dislocated outwards, so that its inner articular facet rested on the outer condyle of the femur, and on flexing the knee-joint the displacement became complete. In the injured limb the patella was in its proper position, and on neither side was there any inclination of the knee inwards.

According to the statement of the boy's mother this abnormal position of the patella had existed from birth, and originally both bones were displaced, but the left patella had been reduced by the mother and kept in place by compresses and a bandage. It is further remarkable, if this was really the case, that the boy had always been able to run about and play as other boys, both he and his mother declaring that he was not more liable to fall or less firm on his feet than other children of similar age.

Two casts have been taken of the knee, one with the limb flexed, the other with the limb extended, and are now in the museum of the hospital.

ROYAL FREE HOSPITAL.

The following operations were performed by Mr. GANT on the 10th instant.

TWO CASES OF HIP-JOINT DISEASE; EXCISION.

Henry W—, aged sixteen, a delicate-looking lad, had suffered from disease in his left hip-joint for over nine months. The left thigh was flexed, and only slightly movable at the joint. At the inner and fore part of the thigh