This article was downloaded by: [University of Cambridge]
On: 08 October 2014, At: 17:37
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 6

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

VII.—On a new chalcosiid moth obtained in Formosa by Mr. H. E. Hobson

Arthur G. Butler F.L.S.

Published online: 12 Oct 2009.

To cite this article: Arthur G. Butler F.L.S. (1889) VII.—On a new chalcosiid moth obtained in Formosa by Mr. H. E. Hobson , *Annals and Magazine of Natural History: Series 6*, 4:19, 53-53, DOI: [10.1080/00222938909460471](https://doi.org/10.1080/00222938909460471)

To link to this article: <http://dx.doi.org/10.1080/00222938909460471>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

VII.—On a new *Chalcosiid* Moth obtained in Formosa by
Mr. H. E. Hobson. By ARTHUR G. BUTLER, F.L.S. &c.

THE following new species was in a series of moths sent to us by Mr. Hobson about six years ago, and of which I have on several occasions commenced drawing up a list, but have been prevented from doing so by more pressing work.

Chalcosiidæ.

Erasmia Hobsoni, sp. n.

♀. Allied to *E. pulchella* of India, but smaller and less brilliantly coloured; the basal metallic green markings on the primaries noticeably smaller; the irregular oblique band beyond them broader and of a deep ochreous (instead of reddish clay-colour); the central metallic green markings narrower, the broad macular white belt less broken up, slightly broader, and with scarcely perceptible greenish edging; all the green streaks on the apical area and external border replaced by grey: secondaries whiter than in *E. pulchella*, with the metallic bluish-green colour confined to the basal sixth, not extending forwards into the cell, the black external border only narrowly edged internally with green, which colour does not extend along the veins or across the border, as in *E. pulchella*. On the under surface the differences are similar, the reddish clay-coloured markings being replaced by clear ochreous, and the green stripes on the external areas of the wings almost wholly obliterated. Expanse of wings 73 millim.

N. Formosa (*Hobson*).

The female of *E. pulchella* expands about 90 millim.

VIII.—On *Isometrus americanus* (Linn.), with a Description of a new Species of the Genus. By R. I. POCKOCK, of the British (Natural-History) Museum.

Isometrus americanus (Linn.).

This species was described by Linnæus (Mus. Adolph. Frid. p. 84, 1754); subsequently (in Syst. Nat. ed. 10, p. 625, 1758) its name was altered to *europæus*, and as *europæus* it