THE ETIOLOGY OF EPIZOOTIC ABORTION.¹

By Professor Dr B. BANG, Copenhagen.

IN most civilised countries epizootic abortion of the cow appears to play a highly important rôle; indeed, in many regions it appears to be the disease which more than any other detracts from the profit of cattle breeding. It is therefore matter for surprise that the disease has, as yet, been to such a slight extent accurately investigated.

Although it must be admitted that abortion may arise from the most various causes (sporadic abortion), and even assume enzootic proportions when such causal circumstances operate simultaneously on a large number of pregnant animals, yet, at the present day, it can hardly be denied that the disease by far most frequently occurs in circumstances that compel us to assume the existence of an infectious agent.

As early as the beginning of the present century this view found expression in England. For example, in the widely read book "The Complete Farmer," 5th edition, 1807, the following occurs: "It is considered certainly contagious, and when it happens the abortion should be immediately buried, and the cow kept as widely apart as possible from the herd, and not receive the bull that goes with them."² With respect to this point it may be remarked that there is a resemblance between the disease now under consideration and tuberculosis. As the result of simple observation the layman had very early recognised the infectious nature of the disease, but the representative of science hesitated to adopt this view; at least this was the case towards the middle of the present century,³ and it was not till the beginning of the bacteriological era that the contagious nature of abortion again gradually came to be generally recognised. This theory was very strongly advanced by Zundel⁴ and Styrl,⁵ and very important contributions to the elucidation of the question were made by Franck, Roloff, Brauer and Johnne.

¹ The copyright of this article is reserved.

² Penberthy. "Enzootic Abortion." *Journal of Comparative Pathology and Therapeutics*, Vol. VIII., page 100, 1895.

³ Hurtrel d'Arboval writes in 1826: "One is not content with the belief that it is epizootic, for it is also said to be contagious, but in that another erroneous view has been put forward." (*Dictionnaire I.*, page 122, *Avortement*.)

Youatt: "There is no contagion, but the result is as fatal as the direct contagion would have made it." ("Cattle," 1834, page 529.)

Rychner: "As regards the causes of abortion, a predisposition is often present, and is very frequently inherited, and hence it comes that epizootic influences tend to produce abortion among many animals." Infection is not even mentioned. (Bujatrik, 1841, page 179.)

Hering writes: "The ordinary causes are not accurately known; there are years in which abortion occur so frequently and with such a wide distribution in one or several species of animals that it cannot be explained on the ground of accidental and local causes." And he adds thereto: "Tögl and Tessier write regarding epizootic abortion, and the latter indeed names it contagious." (*Pathologie*, 3rd edition, 1858, page 690.)

As late as in 1878, Stockfleth warns against the absolute acceptance of the idea of contagion. He says, "Such a theory is, no doubt, quite convenient, as it makes us overlook other more essential causes, which it may be very difficult to detect and to remedy, but it may thus do much harm." The apparent contagion he ascribes to a pollution of the air in the byre. (*D. Zeitschrift für Thiermed.*, Bd. IV., p. 177.)

⁴ Recueil, 1871. In speaking of the causes, however, he distinguishes those which may originate enzootic abortion and those which spread it, and says: "Enzootic abortion is in fact propagated by contagion."

⁵ *Traité d'obstétrique vétérinaire*, 1875. Under the head of "De l'avortement epizootique," one finds a very clear critical exposition of the etiological hypotheses, and several valuable observations by French veterinary surgeons which make the infectious character of the disease probable.

The first, and as yet the only one, to submit the question to accurate investigation was Nocard, who, in December 1885, was commissioned by the Minister for Agriculture to study the epizootic abortion which had for years prevailed among the excellent breeds of cattle in Nièvre.

In the general consideration of the question with which Nocard introduces his report,¹ he mentions that up till that time there had been very few detailed descriptions of the condition of the foetal membranes and fluids, and none at all regarding the uterine mucous membrane; and he makes the very weighty observation that this admits of very easy explanation, since abortion never directly causes death, but is fatal only through the secondary inflammation of the uterus which is set up by the retention of the placenta, and then the alterations which were perhaps the cause of the abortion are difficult to distinguish from those which have set in as the result of sepsis of the foetal membranes. Strictly speaking, all that was known was that "as a rule the after-birth is much altered, macerated, and covered with pus, and that the cow that has aborted furnishes for a long time afterwards a foul, purulent discharge."

After having established the infectious nature of the disease on the basis of admirable clinical observations, he discusses the various hypotheses which might be brought forward in explanation of the anatomical changes causing the abortion. The first, namely, that the epizootic abortion might result from a general affection of the mother, may, he thinks, be put aside, first, because the cow both immediately before and immediately after the act of abortion exhibits no symptoms of disease, and second, because there is no elevation of temperature, and it is impossible to recognise either by macro- or microscopic examination, or by bacteriological investigation, any changes in the urine, blood, or milk, or in the various tissues. It therefore only remains to examine whether abortion is the consequence of a *local affection of the genital organs of the mother*, or whether it is a disease of the foetus or its envelopes.

Although at the *post-mortem* examination of cows which had quite recently aborted, he found between the uterus and the foetal envelopes a considerable quantity of yellowish flocculent purulent material, he nevertheless came to the conclusion that it was not a case of disease of the uterine mucous membrane, because, in sections made through the uterine walls, he could recognise only an intense epithelial desquamation. On the contrary, he thought that he was able to make out from his investigations that epizootic abortion is brought about by a bacterial disease of the foetus and its envelopes.

His bacteriological investigations are naturally of most interest to us. In the puriform substance deposited between the uterus and the foetal envelopes, Nocard found two different microbes abundantly present, namely (1) micrococci, isolated, in pairs, or in short chains of from three to five members; and (2) short, thick bacilli, isolated or united together in pairs. Both could be stained with methylene blue. The micrococci were most abundant in the fluid lying at the base of the cotyledons and round about them, while the bacilli were by far the most abundant in the juice expressed from the follicles of the cotyledons.

In order to cultivate these bacteria he unfortunately did not use a

¹ Recueil de Médecine Vétérinaire, 1886, p. 669.

separation method, but only simple stab cultures in peptone gelatine or bouillon. After forty-eight hours the latter medium had become turbid. When the inoculation was made from cotyledon juice the bouillon contained, it was said, a pure culture of the short bacilli, while when the inoculation was made from the yellow flocculent material the cultures contained both the bacilli and the micrococci. A growth showed itself in gelatine also. After inoculation with cotyledon juice fine parallel streaks grew outwards from the needle track towards the sides, and when the inoculation was made from the yellowish material small whitish, round, granular colonies formed along the line of perforation.

In one case he had the opportunity to examine a foetus that was still contained within the uterus. He made inoculations from the amniotic fluid in bouillon, and thus obtained a culture of micrococci, which he held to be analogous to the micrococci above mentioned.

His examination of aborted foetuses (which had died immediately) did not reveal any microbes in the blood, bile, serous fluids, spleen, liver, kidneys, thymus gland, or lymphatic glands. On the contrary, he found in the intestinal contents in all cases (fourteen) a quantity of different microbes (*microbes variés*). Furthermore, when he inoculated bouillon and peptone gelatine from the medulla oblongata he obtained a growth of micrococci, which, it was said, were very similar to those above mentioned. The idea of making this peculiar examination had been suggested to him by Roux, owing to the analogy between the peculiar continuous bellowing of aborted calves and the altered voice of a rabid dog.

Nocard also makes the observation that aborted foetuses putrefy with striking rapidity, and, in his opinion, this stood in connection with the constant presence of microbes in the intestine.

The recognition of two different sorts of bacteria is by itself calculated to raise a certain amount of doubt regarding the causal importance of these, and, as a matter of fact, he did not succeed in leading decisive proof on this point—that is to say, cause pregnant cows to abort by inoculating them with one of the microbes which he had discovered.

The cause of Nocard's non-success is ascribable to the following: The main cause of it undoubtedly was that he was not in possession of an appropriate nutritive medium, since he did not add blood serum to the substances which he employed. Besides that, the material which he used for examination was not quite free from objection. Most of the cows had aborted some time before the examination of the uterus; the best of them was slaughtered eight and a half hours after the expulsion of the foetus, within which time foreign bacteria might easily have penetrated, and the majority of the foetuses had been expelled one to two days before they were examined. In only one case had he the opportunity to examine the yet unopened uterus of a pregnant cow immediately after slaughter (the only material absolutely free from objection), and unfortunately this case appears to have been not at all an appropriate one. He alleges, it is true, that the signs of impending abortion were in this cow unmistakable, but it is certainly remarkable that it was just in this case, in contradistinction to all the others, that he was unable to recognise any flocculent, purulent secretion between the

uterus and the foetal envelopes. It is therefore in my opinion extremely doubtful whether it was in reality a case of epizootic abortion.¹

Although it thus appears that Nocard's investigations did not lead to a conclusive result, there still remains to him the great credit of having collected good clinical observations in support of the infectious nature of abortion, and also of having pointed out the right way in which to find the cause of the disease.

The investigations which I shall now describe were carried out in my laboratory by myself and my assistant, Mr V. Stribolt, veterinary surgeon. The latter has had a very essential share in them, and all the experiments bearing on the peculiar behaviour of the abortion bacillus with regard to oxygen were carried out by him.

As soon as we had placed before ourselves the task of discovering the cause of epizootic abortion, it became evident that we must obtain a pregnant cow from a herd suffering from the disease, and indeed we required such a cow already showing the signs of impending abortion. In this we easily succeeded, thanks to the friendly assistance of Veterinary-Surgeon Fasting, in Frederiksborg, who was able to indicate in his practice a large farm on which many cows had aborted. From this herd we purchased on the 19th of December, 1895, a five-year-old cow, which had been served with the bull on the 21st of May, and which since the 15th of December had shown the well-known premonitory symptoms of abortion. The cow was there and then slaughtered under Mr Stribolt's directions, and after the genital organs had been removed from the body he bound an elastic ligature round the vagina and brought the preparation to the laboratory, where it was examined six hours after slaughter.

The external surface of the uterus was normal. The os uteri was firmly closed and the cervical canal was filled with the normal thick mucus. After disinfection of the serous covering of the uterus by burning, I made a section through the uterine wall; when the mucous membrane was divided we saw *between that and the foetal envelopes an abundant odourless exudate*—a dirty yellow, somewhat thin, pultaceous material, of a slimy, somewhat lumpy character. At some places where the fluid constituents had run out the exudate was of a semi-solid nature; its reaction was alkaline. When it was allowed to stand in a glass it separated into two strata, namely, superiorly a reddish-yellow cloudy serum, and at the bottom a thick greyish-yellow precipitate.

On cutting through the chorion, we saw under that a thin, clear, apparently gelatinous substance, with very fine membranes running through it; closer examination showed that this was the fine connective tissue lying between the chorion and the allantois, saturated with an œdematous exudate. This was present over the entire extent of the foetal envelopes, and formed a layer $1\frac{1}{2}$ cm. thick. The allantoic fluid was natural in appearance, thin, yellowish, and containing only fine flocculi. Nothing abnormal was observable in connection with the amniotic fluid. The umbilical cord was œdematous. The size of the foetus and the degree of development of the hair on it indicated

¹ From the case here mentioned he obtained a culture of his short bacillus after inoculation from cotyledon juice, and of the micrococci after inoculation from liquor amnii. In the intestine of the foetus he found a quantity of microbes, and in cultures made from the medulla oblongata micrococci made their appearance.

an age of seven months. It was quite fresh, and on section it showed no striking alteration. The pericardium contained a little reddish fluid; the intestinal mucous membrane was, perhaps, rather redder than ordinary; the spleen was in very slight degree swollen, and the blood was fluid.

The examination of a cover-glass preparation made from the yellowish exudate and stained with Loeffler's methylene-blue immediately showed the presence of a very small bacterium, apparently in pure culture. This organism was present in very considerable numbers; many examples lay free, but most striking were the large dense clumps of bacteria. Closer examination showed that these heaps were included within cells, whose bodies were often in this way greatly distended. Sometimes the body of the cell was very indistinct, but as a rule one could still recognise external to the heap a part of the cell body, and often also the cell nucleus. Not seldom the cell body had assumed a peculiar homogeneous appearance.

In the dense heaps the bacteria mostly had the appearance of cocci, but some of the free-lying individuals were of a longer shape, and these were at first regarded as short oval structures; closer examination, however, under very high magnification showed that we had in fact to deal with a small bacillus,¹ whose body contained one, two, or more rarely three, roundish or elongated granules. These granules most readily took up the stain. The length of the bacilli is very variable; the largest examples are about as long as tubercle bacilli.

The before-mentioned granules frequently lie at the ends of the bacillus, but they may occur a little distance from the extremities. They stain with the ordinary anilin dyes, but not by the method of Gram. The bacilli are non-motile. In the subchorial œdema I found no bacteria. In the heart blood of the foetus there were a few, and in the intestinal contents there were many staining granules, but it was not possible to say with certainty whether these were bacteria or not.

This discovery indicates that the *epizootic abortion ought to be regarded as a specific uterine catarrh, determined by a definite species of bacterium*. It is true that the uterine mucous membrane was not strikingly altered, but a chronic catarrh is not necessarily associated with striking anatomical alteration. In my opinion the very abundant exudate, which contained a quantity of shed epithelial cells, pus cells, and detritus, must necessarily have been furnished by the uterine mucous membrane, and not by the thin chorion, and consequently the disease must be regarded as a uterine catarrh.

In the foetal envelopes the œdematous condition of the fine subchorial connective tissue represented the most striking change. This œdema ought probably to be ascribed to a state of stagnation in the foetal circulation. Curiously enough, this condition is not distinctly mentioned in the literature with which I am acquainted, although, according to my observations, it is constantly present. In all the foetal membranes that I have seen from cases of epizootic abortion, it was present to a more or less pronounced degree, and it is to practical

¹ This is most easily recognised in preparations from a bouillon-serum culture. With fuchsin staining the whole body of the bacillus often takes on the colour (see Fig. 4).

veterinary surgeons a well-known fact that the after-birth of a cow that has aborted is uncommonly thick, slimy, and gelatinous, and that the removal of it is difficult, owing to the ease with which the membranes tear.¹

The presence of the uterine exudate is also a well-known fact, but one has not deduced from it with sufficient distinctness the seat and nature of the disease. Most practical veterinary surgeons know that during, or immediately after, the act of abortion, or sometimes even before it, a more or less abundant, dirty, reddish-yellow, slimy, flocculent, pus-like fluid escapes from the vagina, and that this is easily distinguished by its odourless character and its appearance *immediately after the birth* from the discharge which results from an ordinary uterine inflammation. When the after-birth is retained the discharge, of course, even after abortion, gradually becomes offensive in smell in consequence of the penetration of putrefactive bacteria.

After this short digression I return to our first case. The next object in our investigation of the disease, namely, to cultivate the bacterium which we had found in a state of purity, and to furnish evidence that it constituted one definite species, was very easily accomplished by making cultures in test tubes containing serum-gelatine-agar² and placing them in the incubator. At the end of two days in the case of the first tube, and two days after that in the case of the third (the second tube was accidentally broken), there appeared a great number of *very small colonies, which developed only in a definite zone of the tubes*. This zone lay about $\frac{1}{2}$ cm. under the surface of the nutritive medium, and it had a thickness of from 1 to $1\frac{1}{2}$ cm.; colonies were not present either above or below this. We had thus not to do with an aerobic bacterium, which would have pushed its growth as far as the surface of the nutritive medium, and still less had we to do with an anaerobic form, which would have grown as far as the bottom of the tube. The under limit of the zone of growth lay exactly where the upper limit of the growth of a strictly anaerobic bacterium shows itself (the necrosis bacillus, for example). This highly peculiar behaviour of the abortion bacillus towards oxygen made it at once apparent that we had to do with a distinct species, and therefore rendered it in the highest degree probable that there was a causal connection between the bacterium and the disease.

Obviously, the peculiar mode of growth which has just been mentioned greatly facilitated our further studies, as we were thereby enabled to recognise the specific bacillus even when it was present in very impure cultures.

The colonies are, as has been mentioned, very small, often punctiform, the largest scarcely attaining to the size of a pin's head. Their form is roundish, and when slightly magnified their edge is seen to be

¹ Not seldom one finds the foetal membranes somewhat oedematous after normal parturition, but scarcely to such a degree as after abortion.

² This nutritive medium (originally introduced by Stribolt) is very useful, especially for the investigation of anaerobic bacteria. After having liquefied the agar-gelatine (that is to say, $\frac{3}{4}$ per cent. agar and 5 per cent. gelatine) the liquid is cooled to 45° C, and it is then mixed with about half its volume of liquid sterile serum. One now sows some of the material in the first test tube, and in the ordinary way makes several degrees of dilution, pouring a few drops from the first into the second tube, and from that into the third, and so on. The test tubes are then immediately cooled and solidified in a stream of water. This mixture of serum and gelatine-agar remains beautifully transparent, and permits accurate observation of the isolated developing colonies, as well as the starting of fresh cultures from them. The solid medium also serves very well for the preparation of stab cultures by means of a glass needle. Hereafter, for the sake of brevity, I call this nutritive medium simply agar-serum.

finely dentated. They are composed of small bacilli similar in appearance to those found in the uterine exudate.¹

We tried, of course, to obtain cultures from the uterine exudate in other nutrient media, and in other conditions. No growth took place in gelatine-agar, either by the method of plate culture or when the medium was left in the test tubes. In strict correspondence with the above-mentioned sensibility of the abortion bacillus to the oxygen present in the atmospheric air, we found that no growth took place at the surface of solid serum or when the agar-serum was poured out on plates. In glycerine bouillon (5 per cent), however, we succeeded in obtaining a very sparing growth, that is to say, after about fourteen days we saw a very scanty fine precipitate, which contained a few small whitish granules. These granules contained bacilli similar in appearance to the abortion bacilli, and when reinoculated in agar-serum they grew in the typical manner. In pure liquid serum the bacillus also grows only sparingly, but it thrives somewhat better in a mixture of serum (one part) and glycerine bouillon (two parts).

I shall now mention some investigations which still further explain the *peculiar behaviour of the abortion bacillus towards oxygen*.

As was to be expected, one does not succeed in cultivating the bacillus when the oxygen is removed by alkaline pyrogallol solution. The before-mentioned mode of growth of the bacillus in a definite zone under the surface of the agar-serum appears to show distinctly that the organism requires oxygen, but does not agree with such a large quantity of it as is present in atmospheric air, for otherwise it would grow in the very uppermost strata, and even at the free surface of the medium itself. It is therefore all the more striking that Stribolt succeeded in altering in an important degree the conditions of growth of the bacillus, by replacing the atmospheric air above the agar-serum as far as possible by oxygen. He sowed the bacilli in fluid agar-serum, which he then poured into small quadrangular flasks (Nielsen's culture flasks), in which the medium formed a thin stratum on one of the walls. He then conducted oxygen (90 per cent., as it occurs in commerce) into the flasks, and afterwards closed the neck of the flask by means of melted paraffin. *The bacilli now grew very actively in the thin layer of agar-serum as well as at its surface.* He also succeeded in rendering the growth of the bacilli in glycerine-bouillon-serum distinctly more luxuriant by bubbling a large quantity of oxygen through the fluid, and then closing the neck of the flask with paraffin. In these conditions the medium became diffusely turbid, and a somewhat rich, finely granular precipitate of abortion bacilli was formed.²

From this one might suppose that it is perhaps the carbonic acid present in the atmospheric air which interferes with the growth of the abortion bacillus at the surface, and in the uppermost strata of the agar-serum. This hypothesis, however, is contradicted by the fact that the addition of from 4 to 5 per cent. of carbonic acid to the

¹ It ought to be added that small air bubbles not infrequently make their appearance in the nutritive medium, and not exclusively in the zone of growth. They are especially numerous in the first tube when a large amount of material has been made used to inoculate it, and they may then occur throughout the whole of the medium.

² The growth in bouillon-serum, however, was somewhat variable, and sometimes it was very slight or did not take place at all. No doubt this depends upon the greater or lesser tension of the oxygen (*see below*).

oxygen did not affect the result in the above-mentioned experiment. Moreover, the removal of the carbonic acid from the atmospheric air by inserting a cotton wool plug saturated with caustic soda solution in the upper part of the test tubes, had no influence on the growth of the bacillus (the possibility that the nitrogen here played a part is considered below and contradicted).

Strange as it may appear, we must in the meanwhile conclude from the above-mentioned experiment that the abortion bacillus cannot bear the presence of oxygen in the concentration of 21 per cent. (as it occurs in atmospheric air), but that, on the other hand, the presence of an atmosphere very rich in oxygen exerts a very stimulating influence on the organism.

The following experiments¹ appear to show beyond any doubt that *for the abortion bacillus in its behaviour towards oxygen there are two optima*, namely: first, a degree of oxygen tension in the nutritive medium less than that of the atmospheric air; and second, the presence in the nutritive medium of a very high oxygen tension, which, however, lies somewhat under 100 per cent. Between these two optima there is an intermediate zone, in which the abortion bacillus grows badly or not at all.

When one displaces the atmospheric air above the agar-serum in a test tube by oxygen (as may be done by conducting the oxygen through a fine glass tube inserted through the cotton wool plug, and then closing with melted paraffin, or by placing an agar-serum culture tube in a litre flask and filling the latter with oxygen through the rubber stopper), the zone of growth assumes the following appearance: Under the surface there lies a stratum in which, with the naked eye, one discovers no colonies or extremely few (with a slight magnification one sometimes finds several examples).² Then follows with a sharp upper boundary a stratum which has taken a dull whitish-grey colour owing to the fact that it is filled with very numerous and relatively large colonies. In the downward direction this stratum (the upper optimum) becomes gradually clearer because the colonies are less compressed and also somewhat smaller. Hereon follows a rather broad stratum in which, with the naked eye, one discovers only a few colonies; after that one comes suddenly to a stratum which is sharply defined and closely beset with medium-sized colonies (the lower optimum). In this stratum the colonies do not attain to the same size as in the upper layer. This zone of growth is gradually lost in the downward direction, until all growth ceases. That the intermediate zone is never quite free from colonies, may, we think, be explained by the fact that the oxygen can only penetrate gradually into the nutritive medium, and hence at every point of the intermediate zone there must at a certain moment exist just that tension of oxygen which corresponds to the conditions in the lower optimum; at that moment the bacilli there present find the conditions favourable for their development, but subsequently the oxygen tension becomes too great, and then the growth stops. That the optima zones are less sharply defined below

¹ These experiments were carried out mainly at the instigation of the physiologist, Dr Henriquez, and we are indebted to him for many valuable hints regarding the method of conducting them.

² The absence of growth at the surface is to some extent opposed to what was found in the previously mentioned experiments in which plate cultures were made in a flask filled with oxygen, since in these the colonies occurred even at the surface. To explain this one must suppose that in these experiments the oxygen tension was less.

than above no doubt depends upon the gradual penetration of the oxygen.

The following experiment was made in order to show that the above-mentioned conditions of growth are not influenced by the nitrogen present in the atmospheric air. Test tubes or glasses of agar-serum were inoculated and placed in an iron vessel, which was then filled with atmospheric air under a tension of five atmospheres. In these conditions the partial pressure of the oxygen should be almost exactly the same as that of pure oxygen, and if the nitrogen has no influence on the growth of the abortion bacillus one would expect to find that the above-described mode of growth would be repeated; such we found to be the case. The photographic picture reproduced in Fig. 2, illustrates the conditions very distinctly.

It must, however, be mentioned that one does not always succeed in bringing out the before-described conditions of growth so distinctly. When, for example, the pressure in the iron vessel is not kept continually at the same height (as happened on one occasion through a leak in the stopcock), then, of course, both the optima do not appear distinctly. A greater or smaller addition of serum to the gelatine-agar mixture also appears to have a certain influence. Lastly, it appears to be of importance whether one, in adding the serum, shakes the fluid much or little, because with strong shaking more air is absorbed. In one case, for example, the serum and the agar-gelatine were mixed in a flask, then sucked up with pipettes, and distributed in the tubes, instead of as usual making the mixture in each separate tube. Probably through this procedure more than an ordinary amount of air was taken up, and perhaps that is the explanation why in this case the cultures, which were placed in a flask filled with oxygen, showed an upper optimum zone, while under that zone the whole tube (for a height of over 3 inches) contained colonies, which, however, were very small.

Lastly, it ought to be mentioned that when the air over the nutritive medium is much rarefied, as happens when the culture tube is placed in a small flask out of which part of the air is pumped, the zone of growth reaches to the surface. When the rarefaction is continued until the pressure amounts to 2 inches of mercury all growth stops.

Through the kindness of the botanical physiologist, Mr W. Johannsen, our attention was called to the fact that for the higher plants also a condition of affairs analogous to that here recognised for a bacterium has been alleged to exist. For example, A Wieler¹ found that the growth of several plants was accelerated when they were put under bell jars which were filled with rarefied air. With a certain degree of rarefaction he found an optimum of growth; with further rarefaction the rate of growth diminished, until it ceased altogether with a complete absence of oxygen. When the plants were put into a gas mixture containing more oxygen than the atmospheric air he thought that he could recognise a retardation of growth as soon as the partial pressure of oxygen corresponded to an air compression of from two to two-and-a-half atmospheres. With a higher partial pressure an acceleration of growth again took place (second optimum), until ultimately with a sufficiently high oxygen pressure the growth stopped altogether. Subsequent investigations by S. Jentys,² however, have not confirmed Wieler's views regarding the conditions of growth under a high oxygen pressure. At the same time he was able to recognise in

¹ Pfeffer : Untersuchungen aus dem botan. Institut zu Tübingen, Band I., page 189.

² Pfeffer : Untersuchungen, Band II., page 419.

the case of some plants a decidedly favourable influence on growth through increase of the partial oxygen pressure up to one atmosphere.¹

In 1893, Dr M. W. Beyerinck,² in the case of various motile bacteria recognised conditions which have not a little resemblance to the peculiar mode of growth of the abortion bacillus. He describes his principal experiments in the following terms: "One lays a brown bean at the bottom of a test tube, fills the tube nearly quite full with distilled water, and allows it to stand at rest at room temperature. . . . The dry bean immediately becomes swollen up with water, and speedily absorbs the oxygen dissolved in the liquid, while at the same time there diffuse from it certain soluble bodies which serve as nutriment for bacteria. . . . During the swelling of the bean the dried bacteria on its surface spring into activity, and immediately begin to multiply at the cost of the before-mentioned nutritive materials. There thus develops at the surface of the bean a slight turbidity due to the presence of motile bacteria, but this turbidity soon recedes in consequence of the deficiency of oxygen which sets in there. One observes after twenty-four hours, at a temperature of 20° C., or somewhat later at a lower temperature, the following peculiar appearances. Whereas the water in the upper part of the tube remains perfectly clear, a space which is also quite clear gradually develops in the neighbourhood of the bean, since the bacteria accumulated there recede further and further. *At a definite place*, which depends upon the time of the experiment, *the bacteria collect together in an extremely thin stratum* which when regarded from the side appears as a *sharply defined white line*. This stratum I have named a "bacterial level" (Bacterienniveau). With the before-described plan of experiment it is from 2 to 3 cm. distant from the bean after forty-eight hours. He alleges that the bacterial stratum always contained one definite bacillus (*Bacillus perlibratus*) when he used a bean just taken out of its pod. "The 'level' also marks that place where the oxygen coming from above and the diffusion stream of nutrient matters ascending from the bean meet one another, *and here only do the bacteria find the conditions appropriate for their growth.*" The "level" may remain for several days (it remained apparent for weeks or months by using pure cultures of the bacillus grown on gelatine, of which a small quantity was poured out into the bottom of the tube). It gradually ascends until ultimately a position of equilibrium is reached. When he conducted hydrogen into the test tubes over water the 'level' rose to the surface in a few hours, and if he then readmitted the air it again sank to that point where the bacteria found the condition most favourable for their nourishment and respiration. If oxygen is then introduced the "level" sinks still deeper.

For various other bacteria he found that similar conditions existed, but that the height of the "level" varied, which fact he ascribes to the varying requirements of different bacteria for oxygen.

Even in a comparatively thin stratum of water included between a glass slide and a cover glass slightly tilted at one edge, he saw some similar figures (respiration figures), and these were different according as he used ærobic or anærobic bacteria, or such as required only a limited supply of oxygen.

After we had succeeded in one case of epizootic abortion in detecting as the probable cause of the disease a peculiar bacillus, we

¹ From the older investigations regarding the influence of compressed air on the growth of plants I may cite those made by P. Bert (*La pression barométrique*, 1878). He found that germination was delayed in air compressed to two atmospheres, or in gas mixtures with 40 per cent. of oxygen. With an air pressure of five atmospheres the interference with the process of germination was very marked (cited by Jentys, *loc. cit.*) In the case of anthrax bacilli, as is well known, Chauveau in 1884 observed an attenuation of the virulence as a result of the action of compressed oxygen on cultures, and in this way he prepared a vaccin (see Nocard and Leclainche: *Les maladies microbiennes des animaux*, page 169).

² *Über Athmungsfiguren beweglicher Bakterien*. *Centralblatt für Bakteriologie und Parasitenkunde*, Bd. XIV., page 827.

naturally wished to examine a number of other cases of the same sort. Through the kindness of several of our colleagues in the neighbourhood we succeeded in the course of the following months in obtaining suitable specimens from twenty-one cases of epizootic abortion, namely, part of the after-birth which had been manually removed immediately after parturition, and in many cases also some of the exudate that had escaped from the uterus. In a couple of cases we obtained at the same time the aborted foetus. During the past winter we have again obtained many such specimens from different colleagues. In all the cases we were able to recognise alterations which corresponded exactly with those observed in our first case. The uterine exudate had a quite similar appearance, save that it was often somewhat reddened owing to its being mixed with a little blood. In the foetal membranes one could recognise more or less distinctly (often *very* distinctly) the subchorial oedema. The small vessels of the chorion were always more or less injected, and as a rule flakes of pus were adherent to the outer surface of the chorion, while these were always present between the villi of the cotyledons. The appearance of the placental cotyledons was somewhat variable; most frequently they were strongly injected, not seldom hæmorrhagic. They were often of a dirty greyish-yellow colour, seldom, however, throughout their whole extent; more frequently, in part only, and then either in the centre or at the edge while the remaining parts of the same cotyledons were red in colour. All the cotyledons were seldom of a greyish-yellow colour.¹ In cover-glass preparations made from the uterine exudate (or from the pus adherent to the chorion) we could in almost all cases detect the abortion bacilli, partly lying free and partly collected together in intracellular heaps. The number of bacilli appears to be very variable; in some preparations they were immediately apparent, while in others they were very difficult to find. As a rule they are most easily found when the exudate contains very small greyish-white granules, which one may then pick up with a needle. It need hardly be said that such preparations contained not only the abortion bacillus, but always at the same time many other bacteria of various kinds which had penetrated after opening the uterus. Nevertheless, from the great majority of the specimens which reached us in the cold season of the year we were able to isolate the characteristic abortion bacilli by means of separation cultures in agar-serum, and also to obtain pure cultures of the same. In the cases in which the foetus was forwarded we obtained at once from the intestinal contents pure cultures of the abortion bacillus without a trace of foreign bacteria. It is thus proved that *the abortion bacillus may infect the foetus*. Probably this may sometimes occur through the medium of the amniotic fluid. In one case in which culture media were inoculated from the blood of the foetus we obtained no growth, and in the other we obtained immediately a pure culture of the abortion bacilli from the blood and also from the medulla oblongata, fourth stomach, and intestinal contents; this foetus was five months old.

We have also had the opportunity to recognise the abortion bacilli in the uterus in two cases of dead mummified foetus. On the 16th of March 1896, at the slaughter-house here, I made the *post-*

¹ In these cases the foetus had been dead before abortion.

mortem examination of a four-and-a-half years old cow which formed one of the reacting division of animals at Thurebylille (the farm at which I have carried on the main experiments with tuberculin). This cow had calved in October 1893 and then aborted in September 1894, about two months before the date of parturition. Subsequently the cow apparently would not conceive and she was therefore fattened. In the uterus of this cow I found a mummified fœtus about 9 inches long and closely invested by the fœtal membranes. This shrunken fœtus was surrounded by a large quantity of brownish, thick, extremely tough tenacious exudate. As I was aware that epizootic abortion had prevailed in this stock for years, it occurred to me to examine the before-described exudate for the presence of the abortion bacilli. In cover-glass preparations I was unable to detect the bacilli, but a pure culture of these developed in agar-serum; the colonies, however, were unusually slow in developing, since they did not appear distinctly until the ninth day. They were also rather few in number. *It is thus seen that the uterine catarrh which is set up by the abortion bacilli does not always entail the expulsion of the fœtus, but that the result is sometimes merely the death of the fœtus.* When the latter is contained in the uterus one can easily understand that a large quantity of exudate gradually forms around the shrunken envelopes which invest a mummified fœtus, and that this exudate itself in the course of time becomes inspissated and takes on the before-described tough, tenacious consistence.

Subsequently we had a quite similar case. On the 15th of February 1897, Mr Stribolt made a *post-mortem* examination of a heifer rather more than two years old, which had been sent in from the sound division at Thurebylille. This animal had been bulled on the 19th of March 1896; at the beginning of September while at grass it showed the premonitory symptoms of abortion, but these gradually disappeared when the animal was brought into the house. At the *post-mortem* of the animal, which was otherwise sound, Stribolt found in the uterus lesions exactly corresponding with those which I have described above, save that the mummified fœtus was somewhat smaller. In cover-glass preparations made from the yellowish-brown tough exudate bacilli could not be detected, but agar-serum tubes inoculated from the same yielded a pure culture of the abortion bacillus after six days. In this case also the number of colonies was very small. It is thus certainly permissible to draw the conclusion that in those cases in which one finds in the uterus a mummified calf surrounded by a large quantity of yellowish-brown, tough, tenacious exudate, the death of the fœtus has been caused by the specific inflammation of abortion; whereas the death of the fœtus must be ascribed to other circumstances (*e.g.*, torsion of the uterus) when a mummified fœtus is closely surrounded by the uterine wall.¹

The before-described cases are also interesting inasmuch as they illustrate the great vitality of the abortion bacillus. The first cow had

¹ At the *post-mortem* examination of a Thurebylille three-year-old cow, in the year 1893, I found what probably represents an early stage of this affection. The cow had aborted on the 29th of November 1892; when she was slaughtered on the 7th of July 1893 I found in the uterus a large somewhat macerated fœtus, surrounded by dirty reddish membranes, while between them and the uterine wall there was a large quantity of turbid fluid. The fact that in this case the fœtus was not mummified and the uterine exudate not inspissated, is probably explained by the shorter interval which had elapsed between the death of the fœtus and the slaughter of the cow.

probably been bulled in December 1894, or in January 1895, and the death of the fœtus had probably taken place five months later, that is to say, in May or June 1895. The bacilli had therefore remained for at least nine months longer in the uterus, and, nevertheless, they were alive. In the second case the death of the fœtus had occurred in September 1896, and the bacilli were still alive five months afterwards. We are also able to prove the vitality of the abortion bacillus in another way. At the *post-mortem* examination of our first case in December 1895, we collected some of the uterine exudate in sterile test tubes, which were placed in the ice chest. On the 8th March, 13th May, and 14th June 1896 we inoculated agar-serum from these tubes, and in each case obtained a perfectly pure culture. Hence, under these conditions the bacilli retain their vitality for at least seven months.

This property of the abortion bacillus is certainly of the very greatest importance in connection with the disease of which it is the cause. It makes it easy to understand how a cow that has once aborted has a great tendency to abort again during the following pregnancy, since the bacilli remain in the uterus unless a careful disinfection of the uterine cavity is carried out. The vitality of the bacillus also explains the difficulty which is encountered in getting rid of the disease from an infected building.

The above-described investigations made it highly probable that we had actually discovered the specific cause of epizootic abortion, but in order to make the proof complete it was necessary for us to bring about abortion by introducing abortion bacilli into the bodies of pregnant cows. As is well known, several veterinary surgeons have already reported that they have succeeded in doing this by introducing vaginal secretion or part of the after-birth of aborted cows into the vagina of pregnant cows. It is also extremely likely that the agent of infection is, as a rule, taken in in a similar way, the external parts of the genital organs of the cow being brought into contact with infected objects during her sojourn in the byre. When the bull conveys the disease one must assume that infection takes place during copulation. It therefore naturally occurred to us to carry out inoculation experiments by simply introducing pure cultures of the abortion bacillus into the vagina. Our first experiments in that direction, however, proved unfruitful, for the two cows, into the vagina of each of which pure cultures of the bacillus had been introduced, during the following nineteen and twenty-nine days respectively showed no sign of impending abortion, and when they were slaughtered after these intervals we could not detect the slightest pathological change in the uterus or in the fœtal envelopes. However, the result was the same in the case of two other cows into the vagina of each of which we introduced part of the after-birth from cows that had aborted. When these two cows were killed, thirty-three and thirty-five days respectively afterwards, we found no lesions. We were greatly surprised at these results, since experiments carried out in the same way by Brauer were said to have been followed by positive results in from nine to twenty-one days, those by Lehnert after twelve to twenty days, and those by Trinchera after nine to thirteen days. The previous history of the cows used for our experiments was not known, as they had been sold for slaughter in the

Copenhagen cattle market. It was therefore conceivable that they might have come from infected stocks, and have repeatedly aborted and thereby acquired an immunity. We thought at first that this hypothesis might explain the non-success of our experiments. I have, however, since observed that English experimenters in similar experiments have in some cases found a much longer period of incubation, viz., five-and-a-half to ten weeks.¹

It therefore appeared quite reasonable to suppose that our experimental cows would have aborted if we had not slaughtered them so soon, and that we did not find the slightest trace of uterine inflammation might perhaps be explained by supposing that the bacilli had not yet penetrated into the uterine cavity.

Experiments I. and II. In order to guard against the possibility of an immunity accidentally existing in the experimental animals, we now purchased two pregnant cows from a stock in which abortion was unknown. One of the cows was four years old, and the other seven years old, and they had been bulled on the 14th and 16th of January respectively. These cows were placed in one of the horse stables in my division of the clinique, and on the 14th April 1896 we injected into the vagina of each a rather large quantity of pure culture of the abortion bacillus, the culture being partly in glycerine-bouillon-serum, and partly in agar-serum. The seven-year-old cow received cultures that had been grown in an oxygen atmosphere, and the other one cultures which had been made in contact with ordinary atmospheric air. By the aid of a large glass pipette the cultures were introduced as far as the anterior part of the vagina. Inasmuch as the cows exhibited no pathological symptoms during the first five weeks, we feared that these experiments were also going to miscarry, and we therefore repeated the injections on the 23rd of May and 4th of June.

At length, on the 23rd of June we observed in the seven-year-old cow a relaxation of the pelvic ligaments, an œdematous swelling of the labia of the vulva, and a slight flow of mucus from the vagina. The four-year-old cow on the same day showed a slight œdema of the lower parts of the labia of the vulva, but the pelvic ligaments were still firm. Nevertheless the latter cow aborted at 4 A.M. on the 24th June. The foetus was five months old, and it had apparently been dead for some days, as the claws were loose and the cornea opaque, while an abundant exudation of reddish serum had taken place into the subcutaneous tissue and into the thoracic and abdominal cavities. The after-birth was retained, and when it was partially detached at 9 A.M. we observed in it a very abundant subchorial œdema. The chorion was much injected, the cotyledons of a dirty yellow colour, and between the villi of the latter there lay a large quantity of a mucopurulent flaky exudate, which was also discharged from the uterus in moderate quantity. The exudate contained the abortion bacilli, but other bacteria, namely micrococci, had already found their way into it, although the examination was made six hours after the act of

¹ Compare "Second report of a Committee appointed by the Highland and Agricultural Society of Scotland" (7th April 1886), consisting of Dr G. Sims Woodhead and Dr A. P. Aitken, to whom were added Professor M'Fadyean, and more recently Mr Campbell, F.R.C.V.S., Kirkcudbright, "Journal of Comparative Pathology and Therapeutics," Vol. II., page 97.

abortion.¹ The cultures were thus impure, but by means of separation methods of culture in agar-serum we were able to recognise the colonies of the abortion bacillus in the usual stratum.

We had the seven-year-old cow slaughtered on the forenoon of the 24th June, in order that we might have the opportunity to examine the still unopened uterus. From a small incision in the latter we obtained a little exudate, in which we again found the abortion bacilli, and from which we immediately obtained pure cultures of the same in agar-serum. The uterus was then laid on ice in order that I might have the opportunity to demonstrate the specimen five days later on the occasion of a lecture at a meeting of the Veterinary Union. On that occasion I was able to recognise quite the same alterations as I have already described in connection with our first observation (page 128), but the subchorial oedema was still more striking than in that case. The fœtus was quite fresh, and it had therefore beyond doubt been alive when the cow was slaughtered.

By both these experiments we have furnished the complete proof that the bacillus discovered by us is the cause of epizootic abortion. They also prove that the mere deposition of the bacilli in the vagina can produce the disease. The organism must thus be able to propagate through the mucus of the cervical canal. Unfortunately the experiments do not furnish any certain information regarding the period of incubation, owing to the fact that three separate attempts to infect were made. In my opinion, however, it is most probable that the first inoculation was the active one, and that the period of incubation in these cases was therefore ten weeks.

Since the publication of my lecture² we have had the opportunity to pursue our experiments regarding the production of abortion by inoculating with cultures of our abortion bacillus. I shall describe in the first place an experiment with a cow.

Experiment III. This was a six-year-old animal affected with chronic pyelo-nephritis, which I had purchased in October 1896 for the purpose of another experiment. Having accidentally discovered that the cow was pregnant, on the 19th January we introduced into her vagina, in the same way as in the two above-mentioned cases, a culture which had been grown in oxygen. On the 9th April (that is to say, after eighty days) the cow gave birth to a small but apparently full-term calf, which at first appeared healthy, but after about fourteen days was attacked with diarrhoea, and had to be killed. At the time of parturition there escaped from the uterus a not specially abundant, reddish, muco-purulent, flaky exudate, which contained the abortion bacilli. The after-birth, which was somewhat oedematous, was retained, and this induced a rather severe attack of metritis, which however ended in recovery. In this case the abortion bacilli had thus brought about the specific uterine catarrh, and it is also probable that in consequence of this parturition had set in a little too soon, but the pathological alterations developed so slowly that the duration of pregnancy was comparatively little influenced. Although this case speaks in favour of a long period of incubation, I must admit that even it does not justify us in drawing such a conclusion, inasmuch as

¹ Compare Nocard's experience, who found various species of bacteria when he made the examination eight and a half hours after abortion.

² Maanedskrift for Dyrleger, 8 Bind, p. 146 : August 1896.

we had practised a subcutaneous injection of abortion bacilli on the 6th of March. On that date we injected into the neck of the cow about 18 cc. of a bouillon-serum culture; in consequence of this injection the cow had a sharp attack of fever lasting for three days, but at the seat of injection there formed only a small and but slightly sensitive infiltration, which entirely disappeared after a few days. The injection was made on the 6th of March at 3 P.M., and the following is the temperature record:—

6th March, 3.30 P.M.	38°9'	7th March, 5 P.M.	41°3'
„ 5.30 „	39°4'	„ 7 „	41°1'
„ 7.30 „	39°9'	„ 12 „	40°6'
„ 10 „	40°4'	8th March, 4 A.M.	40°6'
„ 12 „	40°6'	„ 7 „	40°0'
7th March, 2 A.M.	40°8'	„ 12 „	40°0'
„ 4 „	41°2'	„ 7 P.M.	40°8'
„ 6 „	40°6'	9th March, 10 A.M.	40°3'
„ 8 „	40°2'	„ 7 „	40°7'
„ 11 „	41°1'	10th March, 10 A.M.	39°1'
„ 2 P.M.	40°7'		

In this experiment we had in view merely to quite provisionally obtain information as to whether by inoculation with cultures or with the toxin obtained from the bacillus one could obtain a diagnostic indication regarding the presence of the abortion inflammation in pregnant cows, and we, of course, intended in later experiments to inject sterile cultures. The marked increase of temperature, which was also accompanied by slight loss of appetite and the passage of soft fæces, appeared to indicate the possibility of a favourable result. However, on the 8th of March we practised a subcutaneous injection to a healthy bull calf, and this showed that the reaction fever can have no diagnostic importance, for in this animal a quite similar elevation of temperature took place and lasted for about four days.

So far as is known to me very little accurate information regarding epizootic abortion among sheep is to be found in the literature. It is generally said that these animals are less inclined to abort than cows; and the cause of abortion in them is usually ascribed to general injuries (sudden change of food, over-exertion, fright, etc.) According to Franck,¹ however, abortion on a large scale is not infrequently observed in flocks of sheep, and he is inclined to attribute the same to infection by straw impregnated with septic material. I have heard from Danish veterinary surgeons and farmers that abortion among sheep has been observed at places where epizootic abortion prevailed among cows, and the investigations of the already mentioned Scottish

¹ Geburtshülfe, 1876, page 276.

Commission showed that the introduction of vaginal discharge from an aborted cow into the vagina of a pregnant sheep could produce abortion. We therefore determined to make some experiments with sheep also. Unfortunately, our experimental animals were already well advanced in pregnancy.

Experiments IV. and V. On the 29th January 1897, we inoculated intravaginally (as in the cows) two sheep, namely, a three-year-old white ewe, which had been tupped on the 7th of November, and a two-year-old black ewe, which had been tupped on the 11th of November. The first received about 20 cc. of an agar-serum culture grown in oxygen, and the black ewe the same quantity of culture which had been grown in atmospheric air.

On the 3rd of April, that is to say 147 days after copulation, and hence probably a little too soon, the white ewe gave birth to two living lambs, which thrived quite well. The cotyledons of the after-birth, which soon came away, were very red, and on them, as well as on the surface of the much injected chorion, there was a not inconsiderable quantity of thick, reddish, turbid exudate, which contained an immense number of the abortion bacilli, partly lying free, and partly included within cells. In agar-serum they grew in the customary manner. This experiment therefore shows that the abortion bacilli, even in sheep, can penetrate from the vagina into the uterus, and there set up a specific catarrh. If the sheep had been inoculated at an earlier period of pregnancy it is very probable that a decidedly premature parturition would have set in. The incubation period in this case was sixty-four days.

The black ewe was killed on the 3rd of April. The result in this animal was negative. The uterus contained two large lambs, and on the outer surface of the envelopes there was present only a very small quantity of a whitish, slimy fluid, in which neither by microscopic examination nor by culture in agar-serum could we detect bacteria. This experiment certainly proves that abortion bacilli introduced into the vagina do not always penetrate into the uterus.

Experiment VI.—It now appeared to us to be of interest to determine experimentally whether it was possible to introduce abortion bacilli into the uterus by way of the blood stream. On the 6th of March 1897, we therefore injected into the jugular vein of a pregnant ewe 18 cc. of a bouillon-serum culture.¹ On the following day the sheep would not feed, respiration was very hurried, and it had a temperature of 40.9° C. On the 8th of March the respiration was still somewhat hurried, but the animal was otherwise better; temperature 39.5°, no swelling in the neck. As early as the 18th of March the ewe gave birth to two lambs, which were small but healthy, and developed quite well. The after-birth came away after an hour; the entire chorion and also the cotyledons were deep red, and covered with a rich red, flaky exudate. The membranes were a little œdematous. The exudate contained very large numbers of abortion bacilli, of which we obtained beautiful cultures.

Experiment VII.—On the 1st of April 1897, we injected 8 cc. of a bouillon-serum culture, which had been grown in oxygen, into the jugular vein of a small ewe, whose time for lambing was not known.

¹ In the opinion of the seller the ewe was due to lamb at the beginning of April.

The animal belonged to a flock which had been used by my colleague, C. O. Jensen, for braxy inoculations, and in consequence of this inoculation it had a large wound on one hind leg. It was somewhat sickly, and therefore the immediate consequences of the experiment were not so pronounced. The fever on the second day was not important, but on the third day the temperature was 41°C . On the 8th of April it gave birth to a small living lamb, weighing 1840 grammes. The after-birth was covered by a sparing quantity of turbid exudate, which contained abortion bacilli as well as a few other bacteria. The sheep died on the 11th of April, in consequence of a not recent, diffuse, sero-fibrinous, double pleurisy and peritonitis. In the exudate I found immense numbers of streptococci which in all probability had penetrated at the skin wound.

Although in this case it may possibly be doubted whether the termination of the pregnancy was accelerated by the injection, nevertheless both these cases prove that *the abortion bacilli can be introduced into the uterus through the blood stream also*. Experiment VI. also shows distinctly that the introduction of abortion bacilli into the blood excites a transient fever (just as after subcutaneous inoculation), but no essential lesions in the internal organs. Only in the mucous membrane of the pregnant uterus does the organism find the conditions favourable for its development, and there only does it set up a specific inflammation.

A knowledge of the possibility of the penetration of the abortion bacilli through the blood stream is also of interest, in so far as it raises the question whether or not the agents of infection may sometimes be taken in through the respiratory or digestive organs.¹ I certainly do not doubt that, as a rule, the sexual organs form the port of entrance. There are cases, however, in which it is difficult to see how infection can take place in this way. I refer especially to those cases where on large farms the heifers are kept in separate byres, isolated from the cow byres; where the heifers are served by a young bull that has never been in connection with cows that have aborted; and where, nevertheless, a not inconsiderable number of the heifers abort. In such cases it is easy to suppose that the heifers, for example, have been infected while grazing in fields where cows have aborted.¹

We have also taken the opportunity to investigate the influence of the abortion bacilli on mares.

Experiment VIII.—On the 1st of April 1897 we injected into the jugular vein of an eight-years-old pregnant mare about 25 cc. of a bouillon-serum culture which had been grown in oxygen. One hour afterwards the temperature was 37.7° , but in six hours it had risen to 40.2° . On the following day at noon the temperature was 39.4° , and in the evening 39.2° ; on the succeeding morning 38.1° . Notwithstanding this transient elevation of temperature the mare showed no symptom of disease, and the appetite was good. On the 29th of April she gave birth to a small living foal, which was strikingly weak, took little nourishment, and breathed quickly. It died on the 1st of May, and at the *post-mortem* examination I found the lungs very incompletely inflated (foetal atelectasis). The after-birth began to come away after three hours, and it was with some difficulty removed. The chorion was perhaps a little redder than usual, and on its outer

¹ Nocard has already called attention to this possibility.

surface there lay a considerable quantity of a reddish-grey, mucopurulent, flaky exudate. There was a little œdema in the subchorial tissue, and a larger quantity between the allantois and the amnion. The exudate contained masses of abortion bacilli, and by inoculating agar serum we obtained beautiful cultures with the typical appearance.

This case is in complete agreement with the sheep experiments, only here it was beyond doubt that the birth was premature, and although from a single experiment one ought not to draw the conclusion that epizootic abortion among mares¹ is always determined by the bacillus discovered by us, that is nevertheless in the highest degree probable; at any rate one ought not to entirely neglect the possibility of the infection taking place from cows and sheep to mares.

We have not yet had the opportunity to make any experiments with goats. I have, however, been informed by Mr Fasting that a goat which had been placed in a cow byre where abortion prevailed (in consequence of an old superstition that this would protect the cows against abortion) twice aborted about seven weeks before full term, and exhibited a discharge similar to that seen in cows. Still another colleague observed a case of abortion in a goat which was kept in a stock where the cows had formerly aborted. It is hence in the highest degree probable that our bacillus is infectious for goats.

It appears to be a well established fact that although a cow that has aborted shows a great tendency to abort again at the next pregnancy, and perhaps even the third year if then pregnant, nevertheless this tendency gradually diminishes, so that it is quite exceptional for the cow to continue aborting year after year.² In correspondence with this, it is the rule that outbreaks of abortion spontaneously disappear after some years, provided that new cows are not being constantly brought in. It has been alleged by several observers that in each successive case the period of pregnancy at which abortion takes place becomes later and later.³ Whether this actually ought to be regarded as a rule must in the meantime be left undecided. In the discussion of these questions (so far as I can see) we lack a solid basis of sufficiently numerous accurate observations, and are therefore exposed to the risk of drawing too far-reaching conclusions from single observations and insufficiently controlled estimates. In the hope that it may serve as an incentive to the collection and publication of

¹ In the year 1751, at the breeding stud at Frederiksborg, 51 out of 208 mares (24·5 per cent.) aborted; in the year 1762, 31 out of 163 (19 per cent.) aborted, whereas the percentage of abortion among mares was ordinarily only 3·9 (Prosch: "Hestens Avl og Pleje," 1870, S. 200). According to Penberthy, at the present day epizootic abortion occurs very frequently in England and North America, sometimes to an alarming extent, so that in large studs the most of the mares are affected (*loc. cit.*, pages 98 to 107).

² Compare Sand's views founded on the reports from Danish veterinary surgeons (*Deutsche Zeitschrift für Thiermedizin.* 21 Band., page 204).

³ Zundel. *Recueil*, 1871, page 167. Penberthy says that while some cows which have once aborted become sterile, and others usually abort again, there is a tendency for the period of gestation to be prolonged after each succeeding conception, and that after a while (three to five years) the disease in a herd into which no fresh cows are introduced tends to die out, as with specific contagious diseases, owing to acquired immunity of the cows themselves, or to attenuation of the virus (*loc. cit.*, page 104). Compare also Sand, who repeats the opinion of several veterinary surgeons, that when abortion has prevailed for two or three years in a herd only the heifers and the cows that have been bought in abort. Nocard (*Recueil*, 1886, page 675) mentions the retardation of the date of abortion in a byre as a common symptom, though not a constant one (*n'a rien de fixe ni de regulier*). He does not, however, expressly state that it is the same cows that abort later the next year. According to his account cows that have once aborted are, as a rule, prepared for the butcher. If the retardation thus applies to other individuals the cause of that would appear to be an *attenuation of the virulence* (see under).

accurate statistical data from as many affected herds as possible, I will here record the experience in connection with the farm of Thurebylille.

The stock of cattle consists of a little over 200 head, of which 130 are milch cows and thirty heifers. Since the spring of 1885 (in which year the management was taken over by the present company) an accurate account has been kept of the circumstances connected with pregnancy, etc. The number of aborting animals has been as follows: 16, 11, 13, 18, 13, 15, 26, 20, 23, 14, 22, 17. And here a peculiar fact shows itself, namely, that whereas during the first year no primiparæ aborted, the number of these that subsequently aborted increased in the following manner: 1, 4, 9, 8, 10, 16, 11, 13, 7, 11, and 9. Thus, out of the total number of 208 animals that aborted, ninety-nine, or about the half, were primiparæ, and in some years about two-thirds were primiparæ. This corresponds well with Sand's views on the subject, and I have also heard from several Danish veterinary surgeons and farmers that a similar state of affairs is of common occurrence on the larger farms, where it is chiefly the heifers and the in-bought cows that abort; the latter, however, as a rule aborting first on the year following purchase, inasmuch as they are usually bought in the pregnant condition, and the bacillus therefore has not the necessary time to penetrate into the uterus.¹

I have in the next place examined into the later history of the primiparæ which have aborted their first calf. In this inquiry I have of course to leave out of account the nine which have aborted for the first time during last winter. Of the others, five were slaughtered soon after abortion, and in the case of two no record has been preserved. *Of the remaining eighty-three only twenty have aborted during the following year*, and seven did not become pregnant. Five out of these seven were slaughtered, and two calved at full term during the third year. *Fifty-four animals that aborted as primiparæ carried their calves to full term during the following year*, and thirty-one of these have also calved at full term for many years in succession afterwards, while others have after one or several normal parturitions proved barren, in which case they were as a rule fattened and slaughtered. Only eight of the fifty-four animals have subsequently after one (five cases), two (two cases), or three (one case) normal parturitions again aborted, most of them only once, but one animal two years in succession. Of the twenty animals that aborted both their first calves, two carried their third calf for the full term, but aborted the fourth again. Four animals have also aborted the third calf; these were then fattened and slaughtered, and this was also done with the remaining fourteen.

In the older writings it is asserted that secundiparæ (cows carrying their second calf) are especially apt to abort, and a physiological explanation of this has been sought in the excessive strain which is put upon such young animals.² On Thurebylille, however, the number of young cows which have gone to full term during their first preg-

¹ Nocard has already called attention to this fact, and says: "On peut en enduire que la maladie doit avoir une assez longue période d'incubation." As an example he mentions a farmer who took over the stock on a farm where abortion had been very prevalent, and introduced into this stock thirty-two cows at from the sixth to the eighth month of pregnancy. All these cows calved at full time, but the next year eleven of them aborted at from the fourth to the eighth month (*loc. cit.*, p. 674).

² Stockfleth d. Zeitschrift für Theiirmedizin. 4 B., page 172.

nancy but have cast their second calf, is in no way great, namely, only sixteen. Of these cases two occurred last winter, and of one there is no record; of the remaining thirteen, six were slaughtered soon after, two were barren in the following year, three aborted during the following year (one of these also the year after that), while one subsequently went three times to full term, and another six times to full term.

Of the multiparous cows whose first abortion occurred later than their first pregnancy, I find that six aborted two years in succession, and one three years in succession; two again aborted once or twice, with intervening normal parturitions.

I thus find that out of all the animals at Thurebylille only thirty cows have aborted in two successive years, and of these only six have cast their calves three times in succession; twenty-two cows that had aborted would not conceive during the following year, but eighty-one cows carried their calves to full term. These figures are not in accordance with the commonly held view that a cow which has aborted is very apt to repeat the abortion during the following year or years.¹

I will not deny the correctness of that view, but I believe that on closer examination it will be found that it applies especially to outbreaks in previously uncontaminated herds. Under such circumstances it may be the rule that the cows first acquire a relative or absolute immunity after they have aborted two or three (or several) times. In large herds, such as are common on our farms, where the plague very frequently continues in existence for many years (probably often for over fifty years), the *virulence of the disease appears to become attenuated*, so that in the case of many animals one abortion serves to confer a temporary or permanent immunity. In this way one may, I think, explain the above-mentioned fact regarding the aborting heifers at Thurebylille. It may be that in such herds there is a congenital indisposition. Such a view is supported by a number of veterinary surgeons, who allege that on large farms the bought-in cows are more apt to abort than those which have been bred on the place.

The fact that at Thurebylille abortion in very many cases takes place at a very late period of pregnancy speaks in favour of an attenuation of the virulence. Among the cases in which the exact period of abortion is correctly recorded, that occurred in thirteen cases one month or less before the completion of full term; in thirty-seven cases between one and two months, in twenty-seven between two and three months, and in only twenty-six more than three months, before full term.

The experience at Thurebylille does not confirm the above-mentioned view regarding a retardation in the case of repeated abortion. Only in six cases did the second abortion set in at a later period of pregnancy than the first; in ten cases the cow cast the second calf (sometimes even the third) at the same or an earlier period than the first.

The account here given of the course of the plague in a large herd in which it was stationary for years is in my opinion of great interest, inasmuch as the facts are well calculated to divert one's thoughts from

¹ Nocard states: "Bien rares sont les vaches qui ayant une première fois avorté parcourent heureusement toutes les phases de la gestation et mettent bas à terme un veau bien conformé" (Recueil 1886, page 677).

the view of infection. When the abortion as a rule affects only the heifers carrying their first calf and in-bought cows, and when in the majority of cases the abortion is not repeated, one is inclined to seek the cause in entirely different conditions, such as conception at too early a period of life, change of food, chills on damp cold pastures, etc.—in fact, in the great arsenal of suggested causes of abortion which from time immemorial have been found in the text-books. I do not doubt that more accurate investigation will show that the majority of cases of “enzootic” abortion (Johne), or “plague-like” abortion (Harms), the cause of which has hitherto been sought in ordinary defects of feeding, general management, etc., will actually be found to belong to *bacterial abortion*. In this connection I may point out that when, for example, Stockfleth, in opposition to the theory of infection, cited the fact that cows often abort while at pasture, this argument at once falls to the ground when one remembers that the period of incubation is often, and indeed probably as a rule, a long one, so that such animals may very well have been infected while they were kept indoors.

In considering the way in which infection takes place too little importance has hitherto been attached to the bull. Nocard, for example, does not mention the possibility of the bull being the carrier of the contagion. Penberthy, on the contrary, speaks of this as *prima facie* probable, and also says that now and again proof of this method of infection is present. Trinchera and Reindl have also given expression to similar views, and we are indebted to Sand for having collected several confirmatory observations from Danish veterinary surgeons. It is indeed self evident that if the vagina is to be regarded as the chief port of entrance of the contagion, copulation must afford the most favourable opportunity of all for infection, as that is the only way in which a direct introduction of the virus into the uterus can be effected. The specific abortion catarrh affords a close analogy with the gonorrhœal inflammation of the mucous membrane, and on *a priori* grounds one is inclined to apply this analogy also to the mode of infection. That this has not hitherto been done as much as it deserves to be no doubt depends upon the fact that people have hitherto been inclined to assume that the period of incubation is a short one. When, however, we know that the abortion catarrh may be, and probably as a rule is, insidious in its course, we must recognise the great probability of infection being spread by means of the bull. It is in the meantime not possible to pronounce a certain opinion as to how often infection takes place by means of the bull, but I am inclined to think that it is of frequent occurrence. Besides those already mentioned by Sand, I know of several similar observations made in Denmark,¹ and the best of these I shall cite hereafter. Mr Winsnes, veterinary surgeon in Christiania, has also informed me that on several farms in an isolated position in Norway, and on which no bull was kept, he has seen abortion break out after cows had been taken to the bull at neighbouring farms where the plague prevailed. It is of very great practical importance to keep this circumstance in mind, and I would here call attention to the risk of spreading the disease which is associated with the custom, otherwise so useful for the improvement

¹ Compare the discussion after my lecture on the 29th of June 1896. *Maanedskrift for Dyr læger*, Bd. VIII., page 165.

of the breed (and very general in Denmark), of keeping a bull to serve the cows of all the members of a society (*Stiervereinen*).

In this connection I must still touch upon the question as to whether it is possible that a cow that has calved at full term may nevertheless furnish a vaginal discharge that is infective. In my opinion the possibility must certainly be admitted when one remembers that from several quarters it has been alleged that aborting cows have at each succeeding pregnancy a tendency to abort at a later period.¹ Furthermore, Mr Paulsen, veterinary surgeon, Ringsted, alleges that in herds in which abortion prevails it may often be observed that the after-birth of cows that have carried their calves to full term shows morbid alterations, being œdematous and covered with exudate, as is common in animals that have actually aborted.² Such animals must be capable of infecting others.

By way of conclusion I may make some observations regarding the prophylactic treatment of epizootic abortion, but in that I can be brief. It is evident that a recognition of the long-suspected agent of infection must in the highest degree inspire confidence in the now generally recommended treatment which is based on the principles of isolation and disinfection. It is precisely this increased confidence in rational prophylactic treatment that I regard as the most valuable acquisition for practice in the meantime. It will be generally admitted that such treatment is capable of yielding very good results, and, if nevertheless, the evil has scarcely diminished, that is no doubt due to the fact that this method of treatment has not as a rule been vigorously and consistently carried out.

So long as we did not actually know the contagium and its way of entrance into the body, and could merely form conjectures, we cannot wonder that the farmer has not generally recognised the necessity of taking immediate measures against infection, or that he has frequently at a later stage grudged the necessary amount of trouble, and the not inconsiderable expense. Now that we are able to tell him that abortion belongs to the category of infectious diseases, and are able to explain to him that the agent of infection is in the uterus and first gains the outer world when the foetus is expelled, that it afterwards may persist for months in the genital organs of the cow and probably for just as long in the floor of the byre and on objects soiled with the vaginal discharges, that it is actually capable of penetrating from the vagina into the pregnant uterus, that the bull may be an active agent in the dissemination of the disease, etc., he will be more inclined to follow the advice of the veterinary surgeon.

The discovery of the abortion bacillus will, it may be hoped, exert on the rational prophylactic treatment of abortion a stimulating influence similar to that which the discovery of the tubercle bacillus did in the case of tuberculosis. Before that discovery it was known that tuberculosis was an infectious disease, and methods of treatment

¹ Zundel says; "If the animals are well fed the period of gestation lasts longer, and the cow which has aborted at the sixth month will abort the second time at the seventh, and then a little before the ninth, so that after three years she will carry her calf to full term." (*Recueil*, 1871, page 467.)

² *Maanedskrift for Dyrlæger*. Band. VIII., page 169. A farmer among whose cows abortion has prevailed has also made a similar observation. He observed in several cows that calved at the normal period a discharge the same as that with which he was familiar in aborted cows.

grounded on that belief were recommended, but the discovery of the bacillus and an accurate knowledge of its conditions of existence first pointed a way to the successful struggle with tuberculosis. As an example of how much may be done by adopting a rational method of prophylactic treatment against epizootic abortion, I will cite the following observation, for which I am indebted to a young colleague, Mr M. Paulsen, veterinary surgeon, Ringsted. A farmer who owned sixteen cows, among which abortion had never occurred, nine years ago allowed seven of his cows to be served by the bull at a neighbouring farm, where abortion had prevailed for some years. When the first of these cows began to show symptoms of impending abortion she was immediately removed from the byre, and that was also done with the other six when they in succession exhibited the same symptoms. In the whole seven cows the after-birth was retained, and Mr Paulsen therefore had it removed by hand within twenty-four hours. The removal was complete in the case of six of the cows, but not in the seventh, which became emaciated and was soon afterwards sold. After removal of the after-birth the cows were for some time washed out daily with carbolised water, and they were not readmitted to the byre until some considerable time afterwards, when they appeared to be entirely free from discharge. During the following year the six cows again became pregnant and carried their calves to full term. No case of abortion has since then occurred in this herd.¹

This observation furnishes a good example of the importance of the bull as a carrier of the contagion, and an excellent model of prophylactic treatment. Since the cows were isolated as soon as they showed symptoms of impending abortion no abortion bacilli were discharged into the byre. The diseased animals were first readmitted after complete disinfection of their genital organs, and this disinfection was also successful in putting an end to the disease in them.

In cases in which aborting cows have not been removed from the byre before abortion it is obvious that one must carry out a careful disinfection of the byre (especially of the stalls of the cows, the gutter behind them, and all the utensils that may have in any way become infected). The foetus and its envelopes must be buried or burnt, and if possible the cow must be removed to another building, and attended by some person who is prohibited from entering the byre. When many cows abort, as is especially the case on large farms, it is unfortunately often impossible, at least in northern countries during the winter, to remove the aborting cows from the byre. One must then at least place them in a separate row, and take care that the discharge from their genital organs is not directly through the channel for the urine, or indirectly through the byre utensils, brought into contact with the vaginal orifice of healthy pregnant cows. In such circumstances it is also well to carry out prophylactic disinfection, such as washing of the external genital parts and their neighbourhood

¹ It is interesting that five of these cows aborted at ten weeks, one at three months, and one at four-and-a-half months, before the normal end of pregnancy. This shows that the abortion catarrh may have an insidious course even when the bacilli are introduced through the act of copulation. The observation is well calculated to strengthen belief in the importance of the bull as an intermediary bearer.

1

3

2

4

in the pregnant animals, and perhaps also to practise prophylactic vaginal injections.

Careful cleansing and disinfection of the floor of the byre are also to be recommended. Under the head of disinfecting agents I would call especial attention to the free use of powdered lime; that is sometimes used by the farmers in this country, and it has the advantage of not imparting any disagreeable smell to the milk, and of not diminishing the value of the dung.

The uterus of the cow that has aborted must be submitted to a careful disinfectant treatment, and the cow must not be put to the bull until she appears to be quite clean; even when this is strictly attended to the same bull ought not to be used for healthy cows. If this cannot be carried out, the preputial cavity of the bull must be washed out with a disinfectant solution every time after he has served a suspected cow.

As has been specially insisted upon by Sand, the system recommended by the older authors of selling the cows that have aborted, and replacing them by newly purchased animals, is not a good one. A careful disinfection of the uterus will, in many cases, succeed in freeing the cows from the specific uterine inflammation, so that they will carry to full term at next pregnancy; and even when this result is not obtained it is certainly better to bear the loss caused by repetition of the abortion, in expectation of immunity setting in, than to provide fresh fuel for the fire. One must certainly admit that this method may be associated with great loss, but here, as is often the case, the first loss is probably the least.

Inasmuch as epizootic abortion belongs to the diseases which leave behind them at least a relative immunity, the possibility of conferring such immunity by prophylactic injections of a vaccin or of an abortion serum naturally suggests itself. To the future must be left the task of accurately investigating this question as well as that of the possibility of a serum therapeutics. Perhaps nothing can be done in this way. Fortunately, however, it will be possible to successfully combat the plague by circumspect use of the old approved means— isolation and disinfection.

EXPLANATION OF PLATE II.

Fig. 1. Colonies of abortion bacilli in a tube of agar-serum. Natural size. (The spots above and below the zone of growth are due to cracks in the glass.)

Fig. 2. Colonies of abortion bacilli in a tube of agar-serum which was submitted to a pressure of five atmospheres. Both optima-zones are shown, and also a few colonies in the intermediate zone. Natural size (in this figure also there are some spots due to cracks in the glass).

Fig. 3. Uterine exudate with large cells completely filled with bacilli. Many free-lying bacilli can be seen (but very indistinctly) by the aid of a lens. The preparation is contaminated with some large bacilli. Magnified 500.

Fig. 4. Abortion bacilli from a bouillon-serum culture, stained with fuchsin. Magnification 1000.

For the photographs I am indebted to the kindness of Professor V. Storch.