I cannot leave this subject without some mention of that interesting form, *Phengodes*. A few years ago the writer found a larva of *Phengodes plumosa* under a stone near Mount Tom, Massachusetts. It emitted light of a brilliant blue colour at the sides of the body segments. Apparently both the larva and the larviform female of some of the tropical species of *Phengodes* and related genera emit light of two colours, as has been reported by a considerable number of observers. In these forms the head, or the region immediately behind it, glows in a brilliant red light, while points along the body segments shine in a white, yellow, green or blue light. Some of these forms are certainly the larviform females, but it seems that the larva also emits exactly the same kinds of light. The reader will find a most interesting account of these luminous forms, accompanied by many bibliographical references, in a paper by Dr. Erich Haase in *Deutsche Entomol. Zeitschrift*, vol. 32, p. 145-167. The forms found in North America appear to be rare, or at least very local, and our knowledge of them is still very incomplete. It is to be hoped that observers will take every opportunity to make observations on these remarkable forms. Gondot (*Revue Zool.*, 1843, p. 17) states that the male *Phengodes* also is brilliantly luminous, and Haase asserts that the male of his *Phengodes hieronymi* emits a greenish light from the under side of the abdomen. The only male of *Phengodes plumosa* which the writer has seen alive certainly gave forth no trace of light, though this does not prove that it may not be luminous at certain times. This specimen was captured at twilight, near Holyoke, Mass., as it flew rapidly across a road through the woods. The date of capture, May 30th, is at variance with the observation of Thomas Say, published in the *Boston Journal of Natural History*, vol. 1, p. 157. Under *Phengodes plumosa* he there remarks: "Not uncommon for a short period in the autumn. Attracted by the candle, they enter the house in the evening and fly repeatedly against the ceiling in their efforts to escape."

THE GENUS *VENUSIA* AND ITS INCLUDED SPECIES.*

BY GEORGE W. TAYLOR, WELLINGTON, B. C.

Probably everyone will agree with Dr. Pearsall's remarks in the April number of this journal as to the necessity for a revision of the North American Geometridæ. The whole group is in a sad state of disorder; the old species are not all well understood and many of the newer ones

---

*See Pearsall, *CANADIAN ENTOMOLOGIST*, XXXVII, 125; April, 1905.

(Dr. Hulst's and Dr. Strecker's), being unfigured and insufficiently described, are very difficult to identify. It will be some time yet, I fear, before anyone will be in a position to attempt a complete monograph, and probably it will be better to take up the work genus by genus, as Dr. Pearsall is doing.

With regard to the new genus *Nomenia*, I have carefully examined all the specimens in my collection from California, Nevada and British Columbia, which, according to the nomenclature of Dr. Dyar's Catalogue, would stand under the name *Venusia duodecimlineata*, Packard, but I cannot find among them any that possess the antennal structure described by Dr. Pearsall.

This does not, of course, prove that the genus *Nomenia* is not a good one, but it shows that *not all* the California so-called *12-lineata* belong to it, and so until Dr. Packard's original types (from California) can be examined it will be doubtful whether the name *12-lineata* should be attached to the *Nomenia* or to the ordinary form. The genus *Venusia* as Dr. Pearsall points out, is represented in North America by a single species, *V. cambrica*, Curtis. The other species listed by Hulst under *Venusia*, form with *Euchæca lucata*, Guenée, a distinct and natural group. But I am afraid we have not yet got these quite correctly named in our lists.

My good friend Mr. L. B. Prout, of London, is very carefully going over, on my behalf, Walker's type specimens in the British Museum; and he tells me that the type of *Tephrosia? comptaria*, Walker, (from Nova Scotia), is certainly not a specimen of *L. perlineata*, Packard, as Hulst supposed, but a *12-lineata*, Packard—eastern form.

I think, therefore, that the name *Euchæca comptaria*, Walker, must be used for the eastern, British Columbia and the bulk of the California specimens of *12-lineata*, Packard. The species now called *Comptaria* will retake its former name *E. perlineata*, Packard, and *Euchæca lucata*, Guenée, with *condensata*, Walker, as a synonym, will complete the group.

I would therefore list the species referred to as follows :

- Venusia cambrica*, Curtis.
- Nomenia 12-lineata?*, Packard.
- Euchæca comptaria*, Walker.
- = *12-lineata*, Packard (part).
- = *salienta*, Pearsall.
- Euchæca perlineata*, Packard.
- Euchæca lucata*, Guenée.
- = *condensata*, Walker.

I may add that the type of *inclinataria*, Walker, (= *inclinata* Hulst), placed by Hulst in the synonymy of *comptaria*, is a specimen of *Xanthorhoe ferrugata*.