centres on the cerebral cortex—so the difference can only be in the end organs. He refers to the experiments of Blix, Goldscheider, and Donaldson, and thinks the apparent results do not amount to a demonstration, and have been worked for more than they were worth. By means of electricity no sensations can be aroused in these so-called heat and cold spots. He is tempted to regard many of the seeming temperature impressions resulting from mechanical stimulus as illusions of sense, just as when one moves a pencil point over the skin, he can discover besides heat and cold spots, "points of cutting, pain, quivering, thrilling, whirling, tickling, scratching and acceleration." The *nerve* is excitable by means of pressure or electricity, but not excitable by means of temperature stimulus. The "*spots*" are excitable through temperature stimuli, but *not* by means of pressure or electricity. He makes a great many interesting experiments upon normal dogs, in which he extirpates certain regions with a view to discovering temperature centres. He finds that each centre is a temperature centre. The evidence from pathology is confusing and on the whole unsatisfactory. We would like to give the results of his exceedingly interesting investigations in detail, but lack of space precludes. Suffice it to say that Dessoir's treatment is a thorough-going success, and is a little gem of its kind. Its real worth commends it to every psychological student. Even his opponents cannot help but concede this.

ACROMEGALY.

By JOSEPH COLLINS, M.D.,

New York.

The disease now known to the profession as acromegaly is of comparatively recent date, it being scarcely six years since it was first accurately described as it showed itself in two of Charcot's patients, both females, at the Salpêtrière. It is unnecessary to say that homage is due to P. Marie for first describing the disease with precision; indeed, Verstraten has given to it the name of Marie's disease, and many writers following him have referred to it in the same way. However much we may deprecate the application of personal names to particular diseases, it would probably be advisable to continue des-

ignating the symptom complex known as acromegaly, as Marie's disease, until we have learned something about its causation and pathology, and then give to it a name based on such acquired knowledge, particularly the latter. The term acromegaly is especially inappropriate, its literal meaning being simply large extremities. In the disease to which the name is applied, however, the changes in the face are as of much importance and as frequent occurrence as in the extremities. Therefore the clinical picture called up by the term acromegaly does not literally suggest the disease described by Marie. Mosler, in an extensive critical study of the disease, and particularly of the arguments put forth by Souza-Leite in his well-known monograph, argues that we should refer to the disease as pachyacria, a word which would in a way signify that the change was a thickening of all the structures of a part involved, and this not alone of the extremities. Recklinghausen was the first to put forward this term pachyacria, and Mosler concurs in its propriety.

Like a great many other diseases that are seemingly of recent date, acromegaly has probably been noticed and described under other names for many decades past. A disease which was probably acromegaly was reported as seen in two cases by Friedreich, under the title of "hyperostosis of the entire skeleton;" another by Fritsche and Klebs, under the title of "gigantism;" another by Lambroso as "general hypertrophy, macrosomia." Then no doubt some of the cases that have come to light under the heading of myxœdema have been of this nature, particularly a case by Henrot reported under this title. Souza-Leite thinks that there can be no question but that a case by Saucerotte-Noel in 1772 was really a typical case of this disease. A critical search of the literature of the past century would be likely to yield other cases which were quite as characteristic as those reported by Marie. This does not in the least detract from the honor due Marie for calling the attention of the profession to this pathological entity in such a lucid and explicit way, that although a great number of cases have since been reported, no additional important symptoms have been produced.

In Souza-Leite's brochure, which is a most complete exposition of the subject up to the date of its appearance, a complete record of all the published cases are to be found. More recently a translation of Souza-Leite's

article under the auspices of the New Sydenham Society has appended to it the notes of nine cases, the reports of which had appeared in the journals between the time of the appearance of the brochure and its translation. This brought the sum total of cases up to forty-eight, which probably includes every case on record till near January, 1891. The principal object of this present digest is to bring the bibliography of acromegaly up to date, and to note and discuss any advances that have been made in respect to the disease, either in its etiology or pathology. It is a deserved compliment to Marie to say, that so thorough and critical was his description of the disease as originally observed by him, that really nothing of importance has been added to the symptomatology of the disease since that time. Although the clinical picture presented by acromegaly is very well known, it will probably be of no disadvantage if we take a glance at the history and development of the disease before we present the notes of the cases that have been recently published.

As regards the etiology of the disease it may be said that it occurs somewhat more often in males than in females. Although we cannot speak definitely concerning its appearance in the different races, racial differences probably offers no exemption. A case has recently been described in a negro (Berkley). Although it may appear at any age, the larger proportion of cases by far first show themselves between puberty and early adolescence (19 to 26, Souza-Leite). It may show itself at a very early age (Rolletson's case), or it may first present itself long after middle age (Erb's, Waldo's, and Hare's cases). Congenital and hereditary influences are excluded from etiological significance in this disease by Marie. He is particularly decided on the point that those cases that have been described under the heading of congenital acromegaly do not belong to the category of true acromegaly as he has described it. Recently, however, some cases have been reported that would tend to show that hereditary influences may have some place in its causation.

Among the exciting causes mentioned are fright (Pel), mental depression (Pechardre), syphilis (Marie, Brigidi, *et al*), exposure to cold (Marie, Verstraeten), traumatisms, either falls or direct injury (Minkowski), rheumatism or gout (Hadden, Ballance, Marie, Souza-Leite, Goodlee, *et al*), alcoholism (Brigidi, Osborne), the

exanthematous fevers, such as variola, scarlatina, malaria, and bronchitis, epistaxis, multiple abscess, severe and prolonged cardiac palpitation, abortion, have all been mentioned by one or more writers as having preceded the appearance of the symptoms of acromegaly, but their connection would not seem to be a very intimate one. In most of the cases published within the last two years there has been absolutely no apparent connection between an etiological influence and the appearance of the disease. In some few cases (Hare, Rolletson, Litt-hauer), the extremities had been large so long as the patient could remember.

The theories that have been forwarded concerning the actual causation of the disease may be stated as follows:

1. Marie's. That the disease is a systematic dystrophy dependent in some obscure way upon the disease of an organ, the pituitary body, the functions of which are as yet unknown. It is only just to say that although a comparatively large number of autopsies have shown that there was a lesion in the hypophysis cerebri in many of the cases, Marie does not support this theory with any more vigor than formerly; that is, he is still inclined to believe that the pathogeny of the disease is yet to be demonstrated. A considerable number of cases yet under observation, of which may be mentioned Osborne's, Bignami's, Bard's, Hare's, present symptoms that are apparently dependent on and traceable to a lesion in the pituitary body.

2. Klebs'. This pathologist, astonished by the existence of the thymus in his case, and taking into consideration a symptom known as post-sternal dullness which had some importance attributed to it by Erb, formulated the theory that the thymus persisting was the seat of vascular budding and proliferation, angioblasts were produced and these formed the foci of the change that went on in the affected parts of the body.

There are serious objections to this theory, the prominent one being that in many cases it is found on post-mortem that the thymus has not persisted; indeed Klebs had a case sometime after formulating the above hypothesis in which there was no post-sternal dullness. Then again there are other conditions attended with persistence, or even hypertrophy of the thymus, and which are not associated with the development of acromegaly. It must also be remembered that it was not

possible to prove that the thymus was the middle point of the vascular budding.

3. Virchow's. This pathologist thinks that in acromegaly we have only half a disease described, *i. e.*, the latter half, with its degenerative details, while at an earlier period it is often accompanied by an increase of muscular power, and is in some instances apparently hereditary. Freund's case lends considerable support to this view.

4. The nervous theory. In spite of the fact that it is based on no positive knowledge, the theory that acromegaly is a nervous disease has taken a very firm hold on the minds of those who have seen most of the disease.

A favorite supposition of physiologists is that there is no organ in the body in post-uterine life but what has some physiological function to perform. This was the principal reason why Klebs' theory was not greeted with any enthusiasm; the thymus is a body which plays no part in post-uterine life. On the other hand, the pituitary body is always present, and the supposition is that it is a glandular body, particularly the pre-hypophysis. The histological structure of this part of the pituitary gland, being made up of convoluted tubules lined with epithelium and highly vascular as it is and its origin, favors this view. Ragowitsch reasoning from the results obtained by Marmesco by removing the hypophysis in some rabbits, is inclined to favor the view that these glands secrete a substance the retention of which in the system, and particularly its action on the nervous system, facilitates the development of acromegaly. If this theory gain acceptance, as pure theory, it would be easy to follow out a process of reasoning such as is suggested by Marie and Marmesco in the "*Archives de méd. et d'anatom. path.*" for July, 1891, and say that the accumulation of this poison in the extremities and face from some special predilection brings about a condition which is the equivalent of the hyperæmia of inflammation and which is followed by the hypertrophic changes so apparent in the disease. If this theory could be in any way proven, the disease would then take its place as one of the auto-toxæmic diseases.

The symptoms of the disease present themselves very gradually and are preceded or not by perversions of the sexual functions (amenorrhœa in women, sexual weakness in man). The first thing noticed is a gradual enlargement of the fingers and hands, and simultane-

ously, or a little later, the feet are involved. The enlargement affects the soft parts as well as the hard, and goes on to an enormous extent. The fingers become sausage shape; the hands, principally because of the great increase at the wrists, battledore shape. The arms are rarely affected. Except in some cases a widening of the epiphysis at the elbows and atrophy of the muscles. The fingers appear short; the nails, though greatly increased in size, do not generally present any trophic trouble, and there is no deformity of the extremities. In the feet the large toe may become increased out of all proportion to the rest of the foot. The changes in the head are principally confined to the upper and lower jaw, particularly the latter, which gives the prognathous face. The thing that may first attract the patient's notice, however, is that each time that he purchases a new hat it must be a size larger (Osborne). The nose, lips, especially the lower, and the tongue are enormously enlarged. The alveolar processes are frequently widened and the teeth separated. The eyelids become elongated, the lashes coarse and unwieldy, and the color of the lids a mild brown. The shoulder girdle is enlarged principally from the participation of the clavicle in the hypertrophy. Occasionally there is an area of post-sternal dullness at the upper part (Erb). The trunk presents two curvatures, a more or less marked kyphosis in the cervico-dorsal region, and an anterior deformity due to involvement of the lower portion of the thoracic cage. The deformity has been aptly compared by Marie to the figure of Punch. There is frequently a slight degree of scoliosis, and always, if the course of the disease is sufficiently long, a compensating lordosis in the inferior dorsal and upper lumbar region. The pelvis is frequently involved, but only to a moderate extent. Ordinarily there is no disturbance of sensation. The hearing is frequently affected and the ears are greatly enlarged. There is often a considerable degree of exophthalmos, and occasionally unilateral hemianopsia and homonymous hemianopsia. The optic disc is occasionally very pale and the veins of the retina turgid. The larynx is frequently augmented in volume not alone in man, but in woman, and the voice becomes more grave, stronger, of longer duration, and especially disagreeable. The speech is slow, embarrassed, guttural, and seems to stick in the mouth. The face takes on the appearance of an elongated oval, and the sexual organs are fre-

quently atrophied. The disease may be preceded in its appearance by any variety of hybrid symptoms, having more or less intimate relation to the disease. They are headache, malaise, mental apathy, varying pains, feelings of premonition, change in temperament, bodily weakness and loss of ambition, markedly increased appetite, great somnolence (Packard), perversion of *morale*, a dry skin, which has a tendency to crack, and polyuria. Many of these symptoms may attend during the course of the disease.

The symptoms have been elaborately classified by Souza-Leite as follows:

1. Objective.
2. Subjective.
3. General.

Each of these two first groups may be subdivided into:

- (a) Constant, fundamental, or principal symptoms.
- (b) Inconstant, accessory, or secondary.

This classification will be found very useful even though it be somewhat artificial; but it is not possible to here enter into a consideration of it.

The following cases have appeared since the English translation of Souza-Leite's brochure, and will, with the list appended to the end of this paper, I hope, bring the bibliography of the disease up to date. It will be remembered that the number of cases in that translation was brought up to forty-eight. The following cases will be numbered from that number onwards, without respect to priority of publication.

OBSERVATION XLIX.—(Stembo.) Woman, aged forty, who first presented symptoms of the disease concomitantly with the appearance of the climatic at the age of thirty. She first noticed enlargement of the feet and hands so that her shoes and gloves would not fit. The enlargement was after noticed in the nose, the lower jaw, lower lip, and the tongue enormously; likewise the palate and a part of the larynx. The ears remained normal; voice deep and masculine; urine increased in quantity, no sugar or albumen, no disturbance of sensation. Right patellar reflex absent; left, weak; internal organs, normal. No apparent causation for the development of the disease.

OBSERVATION L.—(Litthauer.) Male; good family and personal history; married; three children, two died in infancy. Disease first showed itself thirteen years before.

The hands and feet began to enlarge, and went on increasing for two years and then stopped. In the beginning the hands did not give him any particular trouble, but now they often become "sleepy," and he cannot do his work. Formerly his sexual appetite was enormous and was gratified; now the sexual functions are normal. He denies syphilitic infection. Mucous membranes pale; no œdema; no eruption on the skin; no sensory disturbances. Some kyphoses; ideation slow; speech altered; mental hebetude. Pupils normal. Perverted proportion between the size of face and head. Face has assumed an oval shape mainly from increase in size of the lower jaw. Nose, greatly enlarged; the lower lip likewise and somewhat cyanosed. Neck short and thick, circumference 41 c.m. No post-sternal dullness. Heart and lungs normal. Urine and genital apparatus normal. Red and white blood corpuscles normal in proportion. Hæmaglobin remarkably decreased, 40 per cent. Hands cold to the touch, battledore shape; fingers, sausage shape. Marked anæmia. Myopia of both eyes; field of vision limited; retina normal. No disturbance in sensation or electrical reactions.

OBSERVATION LI.—(Pel.) A girl, previously healthy, of good family history, and with no acquired neuropathic predisposition, received just at the time of menstruation a severe psychological trauma. From this time she dates all the symptoms of the present trouble. First she had general nervous symptoms, such as head and body ache, paræsthesia, psychical depression, apathy, etc. Soon after this the hands, feet, and face began to enlarge. In due time a complete picture of acromegaly was developed. She has had amenorrhœa since the time of the fright, although the genital apparatus seems normal. No post-sternal dullness or other evidence of persistent thymus. Thyroid apparently smaller. Author gives it as his opinion that the trouble is neurotic.

OBSERVATION LII. (Bignani.) Female, forty-nine years old. Symptoms began when twenty-five years old, and came on after an abortion which was accompanied by a great loss of blood. Since that time she has not menstruated. There are no hereditary or etiological factors with the exception of the abortion. At the beginning of the disease there were vague pains, paræsthesia, pains in the bones and joints, headache, and then the remarkable increase in size, which was participated in by the lower jaw, the skull, scapulæ, clavicles, larynx, and extremities.

Muscular power began gradually to decrease; moderate degree of exophthalmos on both sides; opthalmoscope showed simple atrophy of the papillæ, right external strabismus, nystagmus, limitation of the movability of the eyeballs in all directions; appetite enormously increased; general muscular weakness without apparent muscular atrophy; skin thickened in places, and, with the exception of slight sensorial disturbances in the hands, sensibility was intact; weakness of the patellar reflexes; cardia hypertrophy; sensation of taste rather slow in its response. The disease made far greater progress in the first part of its course than the last.

OBSERVATION LIII. (Graham.) Male, aged thirty-nine; presented the following marked features: Enlargement of the face; thickness of the lips, especially the lower one; face elliptical in shape; hypertrophy of superior and inferior maxillæ; atrophy of the optic nerves; enlargement and flatness of the hands, feet likewise enlarged; arms and legs normal in size. The disease had lasted for upward of five years.

OBSERVATION LIV.—(Graham.) Male, died aged fifty-one. The disease commenced when he was about twenty-five years of age. He presented the following conditions: Enlargement of the head and face; prominent eyebrows; nose enlarged; hypertrophied superior and inferior maxillæ; lips thickened, especially the lower one; face elliptical in shape and enlarged in every way, resting upon the upper and anterior portion of the chest; neck short and thick; marked kyphosis of the spine in the upper dorsal and lower cervical vertebræ sternum very prominent; enlargement of the scapulæ, scoliosis, lumbar lordosis. Hands were enormously enlarged, as were also the feet. No marked change in the arms or legs with the exception of the knees which were increased in size. No post-mortem examination was made.

In both these cases the soft parts and bones were enlarged; the vitality diminished; the mental faculties impaired; the voice was altered in both, and there was sexual weakness, but children were begat in both cases after the disease had shown itself.

(To be continued in the January number.)