of subperitoneal fat without any sac. The mechanism of production of this variety has been described by Moschowitz, who shows that the subperitoneal fat is usually derived from that between the peritoneal layers of the falciform ligament of the liver. Such a hernia is always small and appears as a protrusion of fat through a short transverse slit in the linea alba, following the course of a small perforating blood vessel. The same writer states that he rarely diagnoses a true hernia with a sac prior to operation. The second type includes in addition to the subperitoneal fat a process of the peritoneum attached to it, making a true sac, but without contents. The third variety resembles the second with the addition of omentum in the sac. The fourth and least common type contains intestine as well as omentum.

Strangulation of epigastric hernias is exceptionally rare. Moschowitz⁴ reports one case, and after a careful analysis of the literature, he has been able to find only ten others. Small intestine was contained in six of these eleven cases. Transverse colon was present in four. Omentum and strangulated bowel were found in four cases and omentum alone in one. It will be noted that the stomach was not present in a single case, although protrusion of this viscus into the hernial sac is often assigned as one of the causes of the distress complained of by the patient. It is much more likely that such distress is due to traction on the parietal peritoneum, or to increased pressure within the peritoneal sac.

The symptoms of epigastric hernias vary greatly. In many cases there are practically no symptoms, while in others nausea and vomiting, dragging pains in the epigastrium, and distress after meals may be severe enough totally to incapacitate the patient. The symptoms in cases of strangulation do not differ from those of any other hernia similarly strangulated.

The diagnosis is usually not very difficult. Lipomas occurring in this region can usually be demonstrated to have no connection with the abdominal cavity. Strangulation of the contents of an epigastric hernia may simulate an abscess in the anterior abdominal wall.

The following case belongs to the fourth variety of hernias of the linea alba between the ensiform and the umbilicus:

History.—A. S., man, aged 40, a Slav, and a laborer by occupation, entered the Presbyterian Hospital, April 9, 1915, on the service of Dr. A. D. Bevan. Three days prior to admission, the patient was seized with a continuous severe pain in the upper abdomen, and a swelling about as large as a lemon was noticed midway between the umbilicus and the xyphoid cartilage. This mass had remained unchanged since first noticed, except that it had become very tender. Vomiting began shortly after the onset of the pain, and had been almost continuous. The patient complained bitterly of thirst, and stated that he had been unable to retain anything for three days. A small lump in the epigastrium had been present since the patient was a child. He stated that this appeared immediately after an accident in which a wagon passed over him. He also described a number of somewhat similar attacks, none of which had lasted more than a few hours. In each attack the above-mentioned swelling became larger and very painful.

Physical Examination.—The patient was a well-nourished man, retching and vomiting continuously. There was an acetone odor to his breath. His temperature on admission was 99.2 F., but his skin was cold and clammy. The abdomen

was moderately distended. There was no fecal matter in the rectum. About 2½ inches above the umbilicus, there was a firm ovoid mass measuring 2½ inches in diameter. This mass was fixed to the abdominal wall in the median line, and was very tender to the touch. The overlying skin was reddened, but this was attributable to the use of hot applications. The diagnosis of strangulated epigastric hernia was made, and an immediate operation performed by Dr. Bevan.

Operation.—Under novocain (1:200) and epinephrin chlorid (1:100,000) infiltration anesthesia, a transverse incision 6 inches in length was made 3 inches above the umbilicus. The subcutaneous fat was carefully divided until a firm lemon-sized mass was reached. This mass was freed of all fat down to its base. It was found to be a hernial sac protruding through an almost circular aperture in the linea alba. The diameter of this aperture was approximately 1 inch. When the sac was opened, a small amount of dark brown fluid escaped. The peritoneal lining was reddened, granular and lustreless. Within the sac were almost 4 inches of small intestine, probably upper jejunum, in addition to some adherent omentum. The bowel was brownish-black, but improved markedly in color after the constricting ring had been enlarged by a lateral incision. A gray band 1½ inches long and one-fourth inch wide, which showed no tendency to change color, was covered over by Lembert sutures and the bowel replaced. The adherent omentum was ligated and resected. There was a small mass of fat adherent to the upper part of the sac. This may have been subperitoneal fat from the falciform ligament, though no effort was made to determine its origin. Closure was made by the Mayo⁵ method, imbricating from above downward by means of three mattress sutures of heavy catgut. The free margin of the flap was sutured to the aponeurosis below by means of interrupted catgut. A small gutta percha drain was placed in the center of the wound, and closure made with interrupted silk sutures. Primary union was obtained, and the patient made a complete recovery.

Occasionally these hernias are confused with the umbilical type, which are much more common. The extreme rarity of strangulation of the epigastric variety, I believe, justifies this brief report. As far as I have been able to ascertain, this is the twelfth case of strangulation of an epigastric hernia to be reported, and the fifth case in which both omentum and intestine were present.

A NEW METHOD OF INTESTINAL ANASTOMOSIS

WITH A STUDY OF THE INTESTINAL MOVEMENTS AFTER ITS USE*

GOLDER L. McWHORTER, S.B., M.D.

Teaching Fellow in Surgery, University of Minnesota Medical School
MINNEAPOLIS

There is considerable difference of opinion as to the choice of methods of intestinal anastomosis. Some of the advantages claimed for the lateral method are: First, that it permits approximation of broad surfaces of serous membrane; second, that a union can be made without regard to the relative size of the ends of the cut intestine; third, that the opening can be made as large as desired.

Although the opening can be increased in size, as advised by Abbe,¹ there is an extensive cutting of the circular muscular fibers and lack of mechanical reestab-

5. Mayo, W. J.: Further Experience with the Vertical Overlapping Operation for the Radical Cure of Umbilical Hernia, *THE JOURNAL A. M. A.*, July 25, 1903, p. 225.

*From the Department of Experimental Surgery, University of Minnesota Medical School.

1. Abbe, Robert: *Med. Rec.*, New York, 1892, xli, 365.

4. Moschowitz, A. V.: *Surg., Gynec. and Obst.*, October, 1914, p. 520.

ishment, which necessarily impairs the peristaltic activity.

Some of the disadvantages, as observed by animal experimentation, are that indigestible substances, bones, straw and hair (Senn,² Reichel³) may accumulate in the blind proximal end of the bowel. The distal end may become invaginated until this portion fills the opening and produces obstruction (Cannon,⁴ Ashton and Baldy,⁵ Robinson⁶), especially if the blind ends are long and not carefully fixed. There may also be danger of interference with the blood supply to the blind ends if too little of the bowel is left beyond the opening (Corbett⁷). Cannon and Murphy⁸ found stasis in the region of the lateral anastomosis as determined on animals which were kept for four days after operation on a waste-free diet and to which they gave an admixture of bismuth in thin boiled starch, observations being made with the Roentgen ray. They found stasis of this fluid, even after twenty-four hours, in the region of the anastomosis after all of the other unabsorbed material was in the colon. Stasis, as described above, was usually found in animals after from ten days to two weeks. They therefore thought it evident that the two united intestinal loops did not act in unison to propel the food, which in the region of the dilated lumen was pushed forward by peristalsis in the intact intestine, acting on the food proximal to the anastomosis. That there is a tendency of all lateral anastomoses to undergo cicatricial stenosis is generally recognized. Bloodgood⁹ has recently ascribed failures, when anastomoses were done in resection of atonic colons, to the two blind extremities becoming tense and painful diverticuli.

As to the end-to-end anastomosis, it may be more quickly done. Cannon and Murphy⁸ were able to observe at no time a stasis of food after the end-to-end union of the severed bowel, and therefore recommend this procedure in preference to the lateral anastomosis because of the rapid resumption of normal func-

tion. Their observations were made on cats and performed with the Connell suture. Küttner, Senn² and Gibson¹⁰ asserted that it is not possible to get this wide contact of serous surfaces, which is desired, without narrowing the lumen of the bowel so much that obstruction is probable, or that stenosis due to contracture or to a diaphragm may result. There is also the possibility of producing ischemic necrosis from interference with the circulation of the mesenteric vessels. The danger resulting from insufficient approximation of serosa is indicated by a case reported by Elsberg¹¹ in which the end-to-end anastomosis was found, at the secondary operation performed at the end of ten days, to have failed at one point.

Other methods of intestinal anastomosis have not come into general use, and some of them are open to serious objections. Some of the better known are perhaps those of Maunsell,¹² Cheattle,¹³ Ferguson,¹⁴ Grant with Turk's modification,¹⁵ Soresi,¹⁶ Horsley,¹⁷ Maury's¹⁸ twine triangular stitch, and Bloodgood's¹⁹ side-to-side ends together; and there are numerous others.²⁰ The artificial aids, according to Reder,²¹ may be convenient in occasional conditions of intestinal surgery, and they are usually either dissolved or passed. In the end-to-side or lateral implantation, Reder states that the technic is more difficult than the end-to-end or lateral union, and besides has the serious drawback of frequently resulting in stenosis.

On account of the foregoing objections to these methods of intestinal union, I have modified the technic of the end-to-end anastomosis and devised a new method. Before I describe the technic, some of the other modifications may be mentioned.

Fig. 1.—In the upper illustration, the dotted lines between the clamps indicate the direction of the usual division of the intestine. The clamps are put on at a 45-degree angle. The dotted lines on the healthy bowel side of the clamps indicate the sites of division. In the lower illustration, the mattress suture to occlude the uncovered area has been introduced, with a needle on both ends to approximate the richer vascular portions first.

2. Senn: *Ann. Surg.*, vii, 13.
3. Reichel, quoted by Cannon and Murphy: *Ann. Surg.*, 1906, xliii, 519.
4. Cannon, W. B.: *The Mechanical Factors of Digestion*, p. 140.
5. Ashton and Baldy: *Med. News*, 1891, lviii, 235.
6. Robinson, F. B.: *Med. Rec.*, New York, xlii, 188.
7. Corbett, J. F.: Personal communication.
8. Cannon and Murphy: *Ann. Surg.*, 1906, xliii, 512.
9. Bloodgood, J. C.: Condition of Patients After Partial or Complete Resection of the Colon, abstr., *THE JOURNAL A. M. A.*, July 10, 1915, p. 193.

10. Gibson: *Ann. Surg.*, lii, 116.
11. Elsberg: *Ann. Surg.*, lii, 116.
12. Maunsell, quoted by Moynihan: *Abdominal Operations*, 1914, i.
13. Cheattle: *Lancet*, London, 1897, i, 1403.
14. Ferguson: *Am. Gynec. and Obst. Jour.*, 1900, xvi, 488.
15. Turk: *Surg., Gynec. and Obst.*, xvi, 324.
16. Soresi: *Surg., Gynec. and Obst.*, xv, 110; *ibid.*, 1915, xx, 225.
17. Horsley, in Fowler: *Treatise on Surgery*, 1907, ii, 65.
18. Maury: *Ann. Surg.*, June, 1905, p. 914.
19. Bloodgood: *Ann. Surg.*, February, 1909, p. 161.
20. Lundholm, E. M.: A New Instrument for Intestinal Anastomosis, *THE JOURNAL A. M. A.*, Sept. 27, 1902, p. 769. Gilbert: *Med. Rec.*, New York, 1904, lxvi, 957. LaPlace: *Philadelphia Med. Jour.*, 1898, i, 1120. Downes: *Philadelphia Med. Jour.*, 1899, iii, 285. O'Hara, M. G.: *Ann. Surg.*, 1901, xxxiii, 170. French: *Am. Med.*, 1902, iv, 90. McGraw, T. A.: Upon the Use of the Elastic Ligature in the Surgery of the Intestines, *THE JOURNAL A. M. A.*, May 16, 1891, p. 685. Senn, William: Pressure Clamp for Lateral Anastomosis, *THE JOURNAL A. M. A.*, April 20, 1907, p. 1350.
21. Reder: *Lancet-Clinic*, 1914, cxii, 662; abstr., *Surg., Gynec. and Obst.*, May, 1915, p. 490.

In the oblique enterorrhaphy described by R. T. Morris²² and M. E. Connell,²³ with the 180-degree mesenteric twist, the ends are cut more or less obliquely to give a larger internal diameter and lessen the danger of gangrene and stenosis. F. G. Connell²⁴ cut the ends slightly convexo-oblique in some experimental unions. Madelung²⁵ was the first to cut the ends obliquely, in the end-to-end unions, to avoid gangrene and give a larger lumen. Parker and Kerr²⁶ cut the ends of the bowel obliquely, uniting them with their basting stitch method. They state, however, that after making the union, absolutely no angulation was apparent, and even at later removal of these specimens, the bowel at the site of suture was apparently straight. As the method of cutting the ends more or less obliquely is not new, though perhaps not emphasized enough now, the application of it, in securing a right angle result, may aid in impressing its value.

The technic of the right angle is modified from the circular end-to-end, in that the bowel is cut at an angle of 45 degrees leaving the oblique mesenteric sides. The mesenteric portions are approximated first by a mattress suture, which when tied on the inside occludes the uncovered area of the bowel and invaginates the edges. With a needle on each end of this suture, they are continued around so that this mesenteric half is first sutured accurately with a hemostatic buttonhole stitch. These are then continued as Connell sutures, ending at that portion of the bowel opposite the mesentery. A Dupuytren's stitch is now taken outside the first stitch to approximate a wide area of serosa. A few stitches join the mesenteries. If the ends are of unequal size, one may be cut more obliquely or an incision may extend up opposite the mesentery, as Curtis²⁷ suggested in the end-to-end union. Where a lateral implantation is desired, the end may be cut obliquely as above (Figs. 1, 2, 3 and 7). The right angle union has advantages over the usual end-to-end in that it not only

has a richer blood supply, but the angle of the blood vessels toward the union lessens the danger of the second row of sutures causing gangrene of the inverted margin of the bowel. There is, therefore, less danger of leakage at the suture line, and also of contraction and stenosis. This can be done as simply as the circular union. Senn² asked, "Does simple flexion, even if acute, without diminution of the lumen of the bowel, give rise to symptoms of obstruction?" He experimentally observed that it did not. There is a flexion of the bowel, perhaps to more than a right angle, in the "elbow anastomosis" of Jeannel,²⁸ in which in an injury to the bowel opposite the mesenteric portion that near the mesentery is left intact and angular union effected.

C. C., aged 8 years, entered the University Hospital, Jan. 6, 1915, and was operated on for a ruptured gangrenous appendix with drainage of abscess. February 6, he returned with symptoms of obstruction, for which an enterostomy was done. March 13, he was operated on by Dr. Ritchie for closure of the enterostomy. This was dissected out, and conditions requiring a small resection found; the right angle technic described above was used. The patient made an uneventful recovery, and was discharged, April 3. July 12, 1915, the boy was in perfect health and gaining in weight.

In the new method of notch enterorrhaphy, the area to be resected is decided on, soft jawed clamps covered with rubber may be applied to prevent the intestinal contents escaping, and the field is packed off with moistened gauze. The mesenteric vessels supplying the bowel to be resected are ligated. Double clamps are put on the ends of that portion to be resected. The bowel is cut between the clamps with a cautery, leaving them temporarily on the ends to be united. The ends of the bowel that are to be united are overlapped for union as far as desired, restoring the direction of peristalsis. The overlapping of about 2 inches is usually satisfactory. A Dupuytren's stitch taken along the sides near the mesenteries approximates them. The clamps temporarily holding the ends of the bowel closed may now be cut away, leaving healthy bowel ends. The sides of the intestines which approximate each other are nearly cut away, beginning at the end of the lumen, leaving a

Fig. 2.—This specimen with a right angle union (Experiment 3) was removed sixty-seven days after operation. The angle of the union when removed was practically the same as when the operation was performed. There were no adhesions.

Fig. 3.—This specimen of a right angle union from Experiment 3 sixty-seven days after operation is shown after having been divided longitudinally. The lumen at the union is distinctly larger than the remainder.

22. Morris, R. T., in Smith, Stephen: *Operative Surgery*, 1887, p. 502.

23. Connell, M. E.: *Med. Rec.*, New York, xlii, 335; *ibid.*, xliii, 125-413.

24. Connell, F. G.: *Philadelphia Med. Jour.*, 1899, i, 37.

25. Madelung: *Arch. f. klin. Chir.*, 1881-1882, xxvii, 277.

26. Parker and Kerr: *Bull. Johns Hopkins Hosp.*, xix, 136.

27. Curtis, B. F.: *Med. Rec.*, New York, xli, 419.

28. Jeannel, in Moynihan: *Abdominal Operations*, 1905, p. 364.

small edge of the side near the mesentery for the approximation of serosa by an inner stitch through all of the coats of the bowel. A similar width of the bowel is left on the convex side for approximation and invagination by sutures. These notched sections of bowel left should have rounded corners in order that they may be accurately approximated, though nearly a rectangle has been removed. A hemostatic buttonhole stitch is started on the side near the mesentery through all of the coats of the bowel, and continuing around the other side as a Connell suture invaginating their edges. The Dupuytren's stitch previously started is now continued around the first suture line, approximating a safe margin of serosa and completely reestablishing the bowel both as to shape and size. The edges of the two overlapping mesenteries are stitched down with a few interrupted stitches (Figs. 4, 5, 6 and 7).

There are very definite advantages to be gained in the notch intestinal anastomosis in that a large opening can be secured as well as one that functions satisfactorily. Therefore, there is no danger from stenosis and contracture. There is an excellent mechanical and a good physiologic union of their fibers. There are no blind ends, which may be sources of danger, as shown above, in the lateral anastomosis, even though later the ends may atrophy and the lateral opening become more axial (Edmunds and Ballance²⁹). There is a good blood supply, as each notched end has its own mesentery intact. There is less danger of interference with the blood supply of the mesentery from injury than in the end-to-end union, and no danger of injuring the blood supply to blind ends, as in the lateral union, which might result disastrously.

The animals operated on had nothing to eat for twenty-four hours, and with few exceptions were given a moderate physiologic dose of morphin about forty-five minutes before the operation. Ether was used throughout. Following the operation, they were given a general diet, except bones, as desired. The ones used for Roentgen-ray work had not been fed for twenty-four hours. In operating, care was used to prevent drying of the peritoneum, and injury. In the cases used for the Roentgen ray, efforts were made to

avoid any unnecessary stimulation and trauma, so all stripping of the intestine to remove contents was avoided, and no clamps were applied to the bowel. In using the Connell stitch, which was used on the last half of the union of the bowel ends, I found it of value to keep this suture taut by tension on the thread only on the mucosal side. The loop and needle thread can be used as a hammock to swing the serous surfaces together, where they are held by tension on the loop, until the needle comes out of the mucosa on the other bowel. This allows more accuracy in getting the stitches even, and prevents distortion of the edges (Fig. 4, lower illustration). For suture material chromic catgut was found satisfactory for the inner suture, and Pagenstecher linen for the second seromuscular row.

Fig. 4.—In the upper illustration, the sides of the two overlapped ends are being approximated near the mesenteries. The two clamps will be cut off with their enclosed crushed bowel, leaving healthy ends, after this suture has been completed near the mesenteries. In the lower illustration, the clamps have been cut away and the sides of the bowel have been cut out, except a narrow margin along the mesentery and also along the convex side for invagination. The inner suture with the buttonhole stitch has been completed and the Connell stitch for the remainder of closure begun. The serous surfaces are swung together by tension on the sutures only when they are on the mucous sides of the bowel. This loop is held till the needle again comes out on mucosa. It is of advantage to have needles on both ends of the inner stitch. Only one is shown above.

PROTOCOLS OF ANIMAL EXPERIMENTS

EXPERIMENT 1 (A. E. 668).—Dog, weighing 18 kg., was operated on, March 1, 1915, the section and a right angle union of the ileum being performed. Convalescence was uncomplicated. The animal was chloroformed, March 18, 1915, and at necropsy there were no adhesions except one small area on the suture line to the omentum. The union was nearly at a right angle, and there was no evidence of stasis. On sectioning, the union was found healed and practically the same size as the bowel.

EXPERIMENT 2 (A. E. 687).—Dog, weighing 8.7 kg., was operated on, March 15, a high intestinal section being made just below the duodenojejunal angle with a right angle union. Convalescence was uncomplicated. March 24, the dog was chloroformed. At necropsy, the abdomen was found to be clean. There was a small adhesion of the omentum at a part of the suture line. There was no evidence of stasis. On section, a good healed lumen of at least normal size was found. March 16, twenty-four hours after operation, a meal containing 60 gm. of bismuth subnitrate in 300 c.c. of buttermilk was given. At twenty-five and forty minutes after the meal, roentgenoscopy revealed that the shadow of the bismuth had passed the pylorus and had reached about 10 and 15 inches below the duodeno-

jejunal angle, beyond the site of the union, with no evidence of stasis or dilatation. March 19, four days after operation, a similar meal was given. Twenty minutes later, on roentgenoscopy, the bismuth shadow was about 15 inches below the duodenojejunal angle. There was a shadow of bismuth in the colon from the previous meal. Thirty-five minutes after eating, before which the dog had vomited part of the meal, there was some advancement in the bowel. The shadow was lacking somewhat in the fundus and body

29. Edmunds and Ballance: *Med.-Chir. Tr.*, lxxix, 254.

of the stomach, but the antrum was well filled as before. There was very little difference in this good motility on the first and fourth days after this high intestinal section.

EXPERIMENT 3 (A. E. 703).—Dog was operated on, March 26, 1915, a section just below the duodenojejunal angle and a right angle union being made. Convalescence was excellent. The dog was chloroformed, June 2, 1915, and at necropsy there were no adhesions present; the right angle was practically the same as after operation. There was no evidence of stasis; peristalsis could be elicited in this region before death. The union was healed. On the same day of the operation, one hour and thirty minutes after its completion, the usual bismuth meal was given. On roentgenoscopy, ten and twenty minutes later, there were contraction rings and a very little shadow beyond the incisura angularis. There was no shadow beyond the pylorus. Forty minutes after eating, roentgenoscopy revealed the antrum filled, and there was a shadow of bismuth 5 inches below the pylorus in the duodenum. Contraction rings were present in the stomach.

EXPERIMENT 4 (A. E. 750).—Dog was operated on, April 30, 1915. No morphin was given. A section just below the duodenojejunal angle was made, and the oblique ends joined at a right angle. At necropsy, May 13, 1915, there was a small area of two loops of bowel adherent some distance from the union, and a small adhesion of the suture line to the omentum. There was no evidence of stasis or dilatation; the union was still nearly a right angle. On section, the lumen was good and healed over. On the day of the operation, one hour and thirty minutes after its completion, the dog was given the bismuth meal. Twenty minutes after eating, roentgenoscopy revealed the stomach filled with bismuth; a contraction ring was present and the bismuth extended through the pylorus about 5 or 6 inches into the duodenum. On roentgenoscopy thirty-five minutes after eating, a contraction ring was present in the stomach, and there was some advancement of bismuth in the duodenum. On roentgenoscopy one hour and five minutes after eating, there was a definite advance of the bismuth to only about 10 inches.

EXPERIMENT 5 (A. E. 519).—Cat was operated on, March 8, 1915, a piece of the ileum being resected, and the ends united by the notch method, described above. Convalescence was good. The cat was chloroformed, May 7, 1915, and at necropsy the bowel at the union was found of normal size, straight, and there were no adhesions. There was no evidence of stasis. On section, the healed suture lines could be seen by a slight inverted edge along them; the bowel was very well reestablished.

EXPERIMENT 6 (A. E. 683).—Dog was operated on, March 12, 1915, a section of the jejunum being made close below the duodenojejunal angle, and the ends united after the

above-described notch method. The resulting reestablished bowel was slightly larger than the normal. Convalescence was excellent. July 6, 1915, the dog was operated on again. There were no adhesions except one very small area on the suture line to the omentum. The bowel at the union resembled that above and below, although a slightly larger diameter in the region of union. Peristalsis at the union could be elicited. The specimen was resected and examined: at no place was there found a decrease in the lumen at the region of the anastomosis, and it was completely healed. On the day after the first operation, March 13, a bismuth meal was given. On roentgenoscopy fifty minutes after eating, the bismuth had passed the pylorus and reached a point about 15 inches below the duodenojejunal angle. There was no evidence of obstruction. On roentgenoscopy sixty minutes after eating, there was some advancement, which was also present at seventy minutes after eating. March 16, four days after operation, a bismuth meal was given. On roentgenoscopy thirty minutes after eating, the bismuth extended about 30 inches, or about twice as far as on the first day after operation, with no evidence of stasis at the union.

EXPERIMENT 7 (A. E. 746).—Dog was operated on May 7, 1915, a lateral anastomosis being performed just below the duodenojejunal angle. This dog is still living. On the same day of the operation, an hour and a half after its completion, a bismuth meal was given. On roentgenoscopy twenty-five minutes after eating, there was very little bismuth in the antrum and none beyond the pylorus. On roentgenoscopy sixty minutes after eating, the antrum was filled and a contraction ring was present. There was no bismuth beyond the pylorus.

EXPERIMENT 8 (A. E. 773).—Dog was operated on, June 7, 1915, to note the movements and passage of bismuth after an operation on the stomach, a typical posterior short loop gastrojejunostomy being done. On the day of the operation, thirty minutes later, a bismuth meal was given. On roentgenoscopy ten minutes after eating, the bismuth extended very slightly into the antrum, and from this a bismuth shadow extended through the newly made opening about 10 or 12 inches into the jejunum. There was no bismuth beyond the pylorus. On roentgenoscopy forty minutes after eating, there was little change except some advancement in the jejunum. On roentgenoscopy one hour and forty minutes after eating, there was little change in the stomach, still no bismuth below the pylorus, but there was advancement to about 15 or 20 inches in the jejunum below the newly made opening. There was a small contraction ring in the stomach.

EXPERIMENT 9 (A. E. 683).—Dog was operated on, July 6, 1915, a specimen of union being resected from just below the duodenojejunal angle and the proximal end invaginated and closed tightly. The distal end was cut obliquely and

Fig. 5.—This specimen with a notch method union, Experiment 6, was removed 115 days after operation. The only adhesion present is shown. This is a small area on the suture line to the omentum. The size of the bowel at the union is the same as the day when the operation was completed.

Fig. 6.—This specimen of a notch method union from Experiment 6 was split slightly to one side of the middle. The suture lines cannot be seen on the mucosal side. The slight increase in size of the lumen was made at the operation. There is no place where the lumen is narrowed.

anastomosed to the posterior surface of the stomach near the pylorus. The dog was not lively after this operation, and became emaciated. The dog was chloroformed, July 15. At necropsy the stomach was markedly distended with clear fluid, and the duodenum very markedly distended to the tied end of the jejunum. The gastrojejunostomy opening was patent, of good size, and when the stomach was lifted, the fluid contents ran into the jejunum through this opening. The lumen was practically healed and about the size of normal bowel in this region.

MOVEMENTS OF THE STOMACH AND INTESTINES
AFTER HIGH SECTION AND UNION BY
THE FOREGOING METHODS

The right angle unions done close to the duodenojejunal angle caused from no apparent delay of food at the pylorus to slightly over twenty minutes. In the one instance a lateral anastomosis caused a delay of over sixty minutes. In both instances of appreciable delay, morphin had been given. Twenty-four hours and four days after operation by the right angle and notch methods, there was no delay observed at the pylorus, as observed by the Roentgen ray.

In contrast to the foregoing, Cannon and Murphy,⁸ in observations on cats after a high intestinal section and union, found a delay of five or six hours at the pylorus, although gastric peristalsis was good all of the time. They ascribe a protective function to the pylorus, in that its closure prevents the stomach contents coming into contact with any wound of the peritoneum until the peritoneal surfaces have been agglutinated, as after five or six hours they may afford efficient support (Abbe³⁰). They observed only a very small effect on movements of the stomach and small bowel due to etherization, to section of the abdominal wall, exposure to the air, or unusual cooling.

Motor function of the stomach, evidenced by rings of contraction, was present in all of my cases, although when delay at the pylorus occurred there was an early imperfectly filled antrum. In the posterior short loop gastrojejunostomy, in which surgical handling and section of both the stomach and jejunum were done, the bismuth was delayed at the pylorus over two hours after the meal, though the bismuth passed almost immediately through the newly made stoma into the jejunum, where advancement was sluggish. After the most gentle handling of both stomach and intestines, Cannon and Murphy observed no movements of the stomach and no discharge from the pylorus for three hours, while rougher handling caused a longer period of loss of motor function. They observed a local paralyzing effect of section and union of the intestine in that no food came closer than 5 or 6 inches during the seven-hour period of observation.

30. Abbe, quoted by Cannon and Murphy (Footnote 8), p. 516.

The localized loss of propulsing peristaltic contraction waves was present in all of my cases of intestinal section and union from immediately after operation to an hour and over, which was the limit of my observations. However, in the case of the gastro-enterostomy, the bismuth passed the union without noticeable delay, although the advancement in the jejunum was sluggish.

The local loss of motor function after intestinal section and suture may be due largely to the handling, and this stimulation excites possibly Bayliss and Starling's law of the intestines, that of inhibition below and contraction above a point of stimulation. The nerve mechanism for the propulsing peristaltic contraction wave, alone, may be inhibited or fatigued, while possibly the rhythmic pendular movements are not thrown out, as these are probably myogenic in origin. It is probable that a general inhibition of the intestines can be produced reflexly, through the splanchnics, by stimulation of the intestines and also probably of any sensory surface (Bayliss and Starling³¹).

Cannon and Murphy found a twenty-four hour stasis at the site of union on cats fed on a waste-free diet, even four days after operation, when a lateral anastomosis was done. This was more marked on a general diet two weeks later. They found no stasis in the end-to-end union twenty-four hours after operation or later, and recommend its use when a rapid resumption of normal bowel function is desired.

In the foregoing observations on the right angle and the notch methods, there was no evidence of stasis at the union, and these methods are also recommended when a rapid resumption of normal bowel function is desired.³²

Fig. 7.—The specimens of a right angle union sixty-seven days after operation, and a notch method union 115 days after operation, are shown filled with bismuth. The union in neither case has narrowed the normal lumen of the bowel.

CONCLUSIONS

1. There are distinct advantages of the notch and right angle over the circular end-to-end and lateral anastomoses.

2. There is as early a resumption of physiologic function in the notch and right angle anastomoses as in the circular end-to-end.

3. The pylorus, if controlled in dogs by a special protective mechanism, does not delay the food there from five to six hours after a high intestinal union, while the gastric peristalsis is active, as described by Cannon and Murphy in cats. There may be from no demonstrable delay to a short delay in certain cases.

31. Bayliss and Starling: *Jour. Phys.*, xxiv, 108, 114, 121.

32. In addition to the references already given, the following are of interest:

George: *Jour. Michigan State Med. Soc.*, September, 1910, p. 431;

Am. Jour. Surg., November, 1909.

Hedblom and Cannon: *Ann. Surg.*, cxxxviii, 504.

Gatch, W. D.: *Aseptic Intestinal Anastomosis*, *THE JOURNAL A. M. A.*,

July 20, 1912, p. 185.

Walker, F. B.: *Intestinal Anastomosis: Presentation of a New, Simple*

and Aseptic Method, *THE JOURNAL A. M. A.*, Aug. 15, 1908, p. 546.

Keen's *Surgery*, 1914, iv, 713-725; vi, 535-539.

Binnie's *Operative Surgery*, 1912.

Murphy, J. B.: *Med. Rec.*, New York, xlii, 665.

4. Following any intestinal anastomosis, a localized loss of motor function remains, and possibly a general reflex sluggishness of the intestinal tract, varying to some extent with the degree of trauma, emphasizing the importance of delicate surgical technic.

5. There may not, necessarily, be a complete temporary loss of gastric motor function after necessary surgical handling for union to the bowel.

LINITIS PLASTICA HYPERTROPHICA (LEATHER BOTTLE STOMACH)

REPORT OF A CASE *

ALEXIUS McGLANNAN, M.D.

BALTIMORE

Linitis plastica hypertrophica is the rather fanciful name given by Dr. William Brinton in 1854 to a condition of the stomach in which there is great thickening of the walls at the expense of its lumen, due to a fibrosis involving principally the submucous, but to a lesser degree all the coats except the mucosa. The interlacing strands of fibrous tissue, which resemble in appearance coarse linen, suggested the name to Brinton.

Two forms of the disease are recognized, one localized, the other diffuse. In the localized form the pathologic change is limited to a certain area of the stomach, usually the pyloric region, although the indurated plaques may occur anywhere in the wall. The diffuse form involves the entire stomach, which is most often shrunken but may retain the normal size or may be dilated. The wall is rigid, and thickened to six or eight times its normal size. The thickening is sharply limited to the stomach, while the thin walled normal duodenum affords a striking contrast. This rigid thick walled contracted stomach suggested the synonym given to the disease.

In gross appearance, the stomach in this disease shows a smooth, pearly white, dull peritoneal surface. The thick wall cuts with a creaking sound, and the limits of the different layers are distinctly shown, a distinction from the scar tissue of a healed ulcer. The glistening bands of the new connective tissue are visible to the naked eye. The mucous membrane is normal or thin. Atrophy and ulceration are said to be

rare, but Thomson and Graham¹ state that in seven of nine specimens examined "there is situated in the center of the affected area a deeply punched out ulcer; the surrounding mucosa, instead of being as soft as velvet, freely movable on the subjacent coats, and showing prominent rugae, is firm and unyielding, is closely adherent to the submucosa, and presents a hillocky surface, corresponding to variations in thickness of the fibromatosis in the submucous coats."

The submucosa shows the greatest change. This loose layer of areolar tissue becomes solid and tough, and so thick that it makes up a great portion of the gastric wall. Segmentation of the circular muscle by the fibrous tissue is easily seen, the colored muscle bundles separated by the partitions of white fibrous tissue standing out clearly. This segmentation is characteristic of linitis plastica, and distinguishes the hypertrophy of the layer of circular muscle in this disease from that occurring in obstruction of the pylorus.

Late in the disease the mucous membrane always shows the effects of long continued circulatory disturbance.

Microscopically the connective tissue proliferation is the most striking feature. The irregular, interlacing bands of fibers surround the blood vessels and traverse the various layers of the stomach. The endarteritis occasionally observed is probably an evidence of syphilis, which is an unquestioned etiologic factor in some of the cases reported. These cases are all of the localized variety, and probably should be kept in a separate group.

The effect of the fibrosis on the mucous membrane gives a varying microscopic appearance. The difficulty of distinguishing between the microscopic appearance of cancer and that of isolated glands separated from their fellows by connective tissue pro-

liferation is so great that it probably accounts for part of the controversy regarding the benign or malignant nature of this disease. Lyle has made an exhaustive study of linitis plastica, and has collected reports of seventy benign cases and sixty additional cases reported as linitis plastica which are malignant.

The difficulties of microscopic diagnosis have been noted, and further confusion is added by the fact that in several cases in which the microscopic picture was considered definite, the patients are living and well so long after operation that the malignant nature of their

Fig. 1.—Linitis plastica, showing the dense fibrosis of the submucosa and the segmentation of the muscularis by the white fibrous partitions. The specimen is the resected pyloric region from a patient with the localized form of the disease. (Photograph by Schapiro.)

* Read at the semiannual meeting of the Medical and Chirurgical Faculty of Maryland, at Westminster, Oct. 28, 1915.

1. Thomson and Graham: *Ann. Surg.*, 1913, lviii, 10.
2. Lyle: *Ann. Surg.*, 1911, liv, 625.