THE STUDY OF FLAME.*

BY PROF. A. SMITHELLS, F.R.S.

ONE of the questions that first meet us in the study of flames is that of the temperature at which in any given case the phenomenon becomes evident. The old idea that there existed a fixed temperature at which inflammation suddenly took place cannot now be maintained, and the term "ignition temperature" has acquired a different meaning. It is now known that in a very great number of cases a mixture of two flame-forming gases, when gradually raised in temperature, will develop luminosity quite gradually with the chemical combination that is being induced. This phenomenon is, of course, known universally in connection with phosphorus, but it is not so widely known in connection with other combustible substances. Sulphur, arsenic, carbon disulphide, alcohol, ether, paraffin, and a whole host of other compounds, inorganic and organic, will phosphoresce as truly as phosphorus itself; in fact, phosphorescent combustion is the normal phenomenon antecedent to what we ordinarily call flame. This is, after all, only in harmony with the general truth that chemical combination between two gases does not set in suddenly, but comes into evidence quite gradually as the temperature is raised from a point at which the action, if it occurs at all, is so slow as to be negligible. The increase in the rate of combination is, of course, very rapid as compared with the increase of temperature, a difference of about 10 deg. C. (50 deg. F.) serving to double it. The interval between the beginning of phosphorescence and the production of vigorous flames may therefore be very short. In the case of phosphorus this interval, being from 7 deg. to 60 deg. C. (44 deg. to 140 deg. F.) includes ordinary atmospheric temperatures; hence the phosphorescence of phosphorus is a phenomenon that could not well be overlooked. If the prevailing terrestrial temperature were below 7 deg. C. (44 deg. F.), at which, under normal air-pressure, the phosphorescence of phosphorus ceases, it is possible that this element might never have acquired its peculiar reputation; it would not have shone in the dark, and in lighting it with a taper the phosphorescent interval would have been passed over as quickly as is ordinarily the case in the ignition of sulphur, paraffin, and other common combustibles. To make phosphorescence apparent in these last cases it is necessary to take special care to heat up a mixture of the combustible gas and air gently, and to maintain it at a temperature approaching, but not quite reaching, that of ignition. There is no simpler way than that used by Sir William Perkin, who brought the combustible substance near to, or in contact with, a massive metal ball previously heated to the suitable temperature. The change from phosphorescence to ordinary flame is not sudden, but the appearance of ordinary flame is the end point of a continuous, though rapid, development. This end point is the temperature of ignition. What, then, determines the temperature of ignition? The answer to this question has been given with characteristic conciseness by Van't Hoff, as "the temperature at which the initial loss of heat due to conduction, etc., is equal to the heat evolved in the same time by the chemical reaction." We may obtain a clear idea of the meaning of temperature of ignition by supposing a combustible mixture of gases such as that of air and the vapor of carbon disulphide to issue through an orifice into an indifferent atmosphere. If we surround the orifice by a ring of platinum wire, which is gradually heated up by a current of electricity, a flame will gradually make its appearance. If, as soon as this is observed, the heating of the wire by the current be discontinued, the flame will disappear; it is, in fact, not self-supporting, but depends on the accessory supply of heat through the electrically-heated wire. If, now, we raise the ring to a higher temperature, we shall get a brighter flame, owing to an increased rate of chemical action, and at last we shall reach a point where it is possible to cut off the electric current without causing at the same time the extinction of the flame. This is the true temperature of ignition, the temperature at which the reaction proceeds at a rate just sufficient to overbalance the loss of heat by radiation, conduction, and convection from the burning layer of gases, so that the next layer is put in the same state, and steady combustion proceeds. Phosphorescence has been spoken of as degraded combustion, and, though literally the appellation is correct, it is liable to be misunderstood. Again, it is often supposed that phosphorescence is necessarily associated with the formation of incompletely oxidized products. This may be the case in a chemical system which is capable of affording different products at different temperatures, but it is not an essential feature; the phosphorescent combustion of sulphur, for example, affords nothing but sulphur dioxide. Temperature of ignition is, then, neither a temperature at which

combination suddenly begins, nor one dependent solely on the nature of the combining gases. It will vary with the proportion in which the gases are mixed and with their pressure and other circumstances. Notwithstanding the simplicity of this conception, it must be admitted that there are many obscure facts connected with the ignition of gases. The inflammability of gaseous mixtures is not necessarily greatest when they are mixed in the proportions theoretically required for complete combination; the influence of foreign gases does not appear to follow any simple law; the presence of a very small quantity of a foreign gas may exercise a profound influence on the ignition temperature as in the case of the addition of ethylene to hydrogen. When a mixture of methane and air is raised to its ignition temperature, a sensible interval (about ten seconds) elapses before inflammation occurs. These facts are cognate to others which have increased upon us so abundantly in connection with the influence of moisture on chemical change. The study of the oxidation of phosphorus in particular brings us among rocks and shoals. Apart from the influence of moisture on the combination, we have the limitation of the process by a certain tension of oxygen and by minute quantities of a vast number of chemical substances, among which, in spite of much labor, no other common bond can be found. We do not know what oxide is initially formed in the oxidation, and the existence of the oxides P_4O and P_2O is as confidently disputed as it is affirmed. There is some reason for believing that the phosphorescence connected with phosphorus succeeds the formation of one oxide and accompanies the formation of another. The state of the oxygen, whether atomic, ionic, or molecular, which acts on phosphorus, the induced oxidation of other substances, the ionization of air accompanying the oxidation—these are all matters concerning which there exists a bewildering literature that hangs over us like a cloud. The subject, simple as it appears at first sight, is really involved with a vast number of unsolved chemical problems, the elucidation of which would throw much light on chemical action in general. The structure of flames has always been regarded as dependent upon the chemical changes taking place in the differentiated regions, but until recent times little attention has been given to any question beyond the cause of the bright luminosity of hydrocarbon flames. In a flame such as that of hydrogen or carbon monoxide, where we have some reason to suppose that the same kind of chemical transaction is taking place throughout the region of combustion, we should not expect to find a differentiation of structure, and, as a matter of fact, we do not find any. Erroneous ideas have gained currency from the use of impure gases, and hydrogen is still described as burning with a pale blue flame, although Stas long ago stated that if the gas is highly purified, and the air freed from dust, the flame even in a dark-room can only be discovered by feeling for it—a fact consistent with the line spectrum of water lying wholly in the ultra-violet. The presence of a very small quantity of free oxygen in carbon monoxide destroys the perfect simplicity of the single shell of blue flame with which the purified gas burns, and in other flames small quantities of gaseous impurities or of atmospheric dust give rise to features of structure and halos which have been frequently supposed to pertain to the flame of the combining gases. The fringe of a flame in air may be often tinged by the presence of oxides of nitrogen. No flame better illustrates the relation of structure to chemical processes than that of cyanogen, where the two steps in the oxidation of the carbon are clearly marked out in color. Apart from hydrocarbon flames, very few others have been carefully explored from this point of view. The use of single hydrocarbons instead of the mixtures present in coal-gas and other common combustibles has simplified the study considerably. Two problems stand out prominently: one is to trace the steps in the oxidation of the hydrocarbon, the other to account for the bright patch of yellow luminosity. With regard to the question of the luminosity, I do not think there is any longer doubt about its being due essentially to the separation within the flame of minute solid particles of what is practically carbon. The separation seems to be adequately explained by the high temperature of the blue burning walls of the flame, which decomposes the unburned hydrocarbon within. In a similar way, arsenic and sulphur and phosphorus are liberated within flames of their hydrides; but these elements, being volatile, do not appear as solids unless a cold object be placed within the flame. In the case of the hydride of silicon the liberated element at once oxidizes to form the solid non-volatile oxide, which gives a bright glow. The mode in which a hydrocarbon yields carbon by the application of a high temperature has been the subject of experiment and of hypothesis; but neither the view of Berthelot, that the carbon results from a con-

tinuous coalescence of hydrocarbon molecules with elimination of hydrogen, nor that of Lewes, according to which the formation and sudden decomposition of acetylene is the essence of the phenomenon, appears to be in harmony with the experimental facts. It is certainly not easy to ascertain experimentally the changes undergone by a single hydrocarbon as its temperature is raised, and at the last it may be objected that the course of events in contact with the solid walls of a containing vessel is not necessarily the same as that within the gaseous envelope of a flame. I am glad to think that there is promise of further light on this subject from the work of Prof. Bone. Our knowledge concerning the question of flame temperatures has been much improved in recent times, thanks mainly to the admirable work of M. Le Chatelier. The well-known memoir of Mallard and Le Chatelier on the explosion of gases supplied the data which first permitted of a moderately exact calculation of flame temperatures, and the perfection of the thermo-couple by M. Le Chatelier gave us the first instrument that could be used directly for making a satisfactory measurement. The uncertainty connected with this subject may be well illustrated by quoting the temperatures that have at different times been ascribed to the flame of coal-gas when burnt in a Bunsen burner, where we have had values varying from 1,230 deg. to 2,250 deg. C. (2,246 deg. to 4,082 deg. F.) With regard to the use of thermo-couples, the practical difficulties have been successfully met. The chief difficulty is, of course, to secure that the thermo-junction attains as nearly as possible the temperature of the region in which it is immersed. As ordinary flames consist of thin shells of burning gases, on either side of which there is a very rapid fall of temperature, it is necessary to use thin wires, and to dispose them so that there is no appreciable drain of heat from the junction. By using wires of different gage for the couples it is possible by extrapolation to arrive at a temperature for a couple of infinitely small cross-section, and it is also possible to make a correction for the superior radiating power of the couple as compared with the flame-gases. With this last correction a maximum temperature of 1,871 deg. C. (3,400 deg. F.) was obtained as the highest temperature of the flame of a Bunsen burner burning coal-gas. The consideration of flame-temperatures has become of increasing importance in the arts owing to the use of the Welsbach mantle as a means of deriving light from coal-gas. The great improvements which have been made in the efficiency of atmospheric burners depend primarily on the fact that the smaller the external surface we can give to a flame consuming gas at a fixed rate the higher must be the average temperature; and since the emission of light from a mantle is proportional to a high power of the absolute temperature, a small increase of temperature is of great effect on luminosity. The acetylene-oxygen flame in which a temperature of about 3,500 deg. C. (6,332 deg. F.) prevails, not very different from that of the electric arc, is the hottest of the hydrocarbon flames, and finds some important practical uses.

There remains the general question of the luminosity of flames containing nothing but gases. The older explanation of the emission of light from combining gases said no more than that the energy liberated during the reaction and appearing as heat raised the product to incandescence—that is to say, so increased the velocity of its molecules and the violence of their collisions that vibrations were set up whose wave-lengths lay within the limits of visible radiation. This explanation has long been questioned, and there is now, I think, a very general agreement that it will not suffice. The average temperature, in fact, prevailing in a flame, if attained in the product of combustion by the supply of heat from outside, does not suffice to make that substance luminous. We are, therefore, thrown back upon the conclusion that the generation of light in a flame is not a consequence, though it is an accompaniment of the elevation of temperature. The question now is, Can we go any further? To do this we are led to consider individual molecular transactions instead of statistical averages, and the view presents itself that the combining atoms may, in losing their chemical energy, form directly systems of independent vibration where the radiation is such as to fall within the limits of visibility. If we picture such vibrating systems momentarily formed, it is easy to see that by their collision one with another they may acquire in a secondary way increased translational motion, and so lead to a state of things where the greater part of their energy is degraded in the form of heat. The high temperature of a flame would then be a consequence rather than a cause of its light. This subject of the mechanism of luminosity, however, like so many others, has now become involved with the theory of electrons. Some years ago attention was called to the scantiness of our

*Abstract of an address before the Chemical Section of the British Association for the Advancement of Science.

knowledge of the chemical changes that take place when metallic salts are used in flames for the production of spectra. Though there was general agreement that, for example, the yellow flame produced by common salt was due to the liberation and glow of metallic sodium, there was no agreement as to how the sodium was set free. Arrhenius, pursuing the analogy which exists between the laws governing matter in the gaseous state and in the state of dilute solution, had previously been led to the view that the electrical conductivity of flames containing salt vapors was due to ionization of the salt throughout the volume of flame. It appeared possible, therefore, that the luminosity might be ascribed likewise to the metal separated in the ionic state. Experimental investigation undertaken with a view to elicit information on this subject seemed to favor the view that the metal was reduced by chemical processes, and that it glowed in the un-ionized condition. Evidence seemed to point to the conclusion that, for example, when common salt is introduced into the flame of coal-gas the sodium chloride yields sodium by the conjoint action of steam and reducing gases; when liberation of the metal was prevented by adding a large quantity of hydrochloric acid to the flame the glow disappeared, but the conductivity was not always diminished. The fact that sodium salts, including the chloride, impart their characteristic glow to the flame of cyanogen and to other flames in which water is absent leads to some difficulty in finding a chemical explanation, and it must be admitted that a direct thermal dissociation of an alkaline halide or oxide is not out of the question. The interval of detachment of the metallic atom may be exceedingly brief, but it must be remembered that even so short a time as the interval between the molecular encounters in a gas at a high temperature is still sufficient for the emission of thousands of undisturbed characteristic vibrations. Very interesting and important investigations have been carried out by Lenard, who has shown that the stream of luminous vapor produced from a sodium salt in a Bunsen flame is deflected in an electric field in such a way as to indicate that the vapor is positively charged; but he gives reason for believing that the charged condition is intermittent with the neutral condition. The lines in the spectrum of an alkali metal are divisible, as is well known, into distinct groups or series, in each of which the oscillation frequencies corresponding with the lines are in a definite mathematical relationship. The principal series, which include the lines seen individually as such in ordinary flame spectra, are according to Lenard, due to the electrically neutral atoms. In a salted spirit flame, and in other flames of low temperature where only lines of the principal series are represented, the stream of luminous gas does not behave in an electric field as if it were charged. In the flame of coal-gas burned in a Bunsen burner, the salt-vapor gives, in addition to the distinct lines of the principal series, diffuse bands of luminosity on the dark background, which, according to Lenard, represent the undeveloped subordinate series; and it is the atoms emitting these series that are deflected in the electric field. It is inferred, therefore, that the light in a salted Bunsen flame comes from different groups of centers of emission—the principal series from the neutral atom, and the lines of the first, second, and third subordinate series from atoms which have lost respectively one, two, and three electrons. Lenard goes further, and shows that the salt-vapor in a Bunsen flame, as in the flame of the electric arc, emits these different kinds of radiation from different structural regions; thus the vapor at the edge of the flame is electrically neutral and gives only the lines of a principal series. The negative electricity in a salted flame would, according to Lenard, be disembodied, and recent experiments by Gold confirm the view that the negative carrier in flames is a free electron. The subject is obviously one of fundamental importance in relation to spectrum analysis, and my own slight connection with it has only strengthened my opinion that there is still a great deal connected with the genesis of spectra that requires the attention of the chemist even more than that of the physicist.

An invention lately patented by the Deutsche Gasglühlicht Aktiengesellschaft relates to tungsten filament glow lamps, and consists in employing carbon suspenders in place of the suspenders of metal or the refractory metallic oxides which have hitherto been generally used for this purpose. As carbon possesses the property of good radiation for heat rays, this support remains comparatively cool, even near the point of contact, and can, therefore, be made very thin. Carbon suspenders have been found to be unsatisfactory for osmium lamps, owing to the formation of osmium carbon compounds; but this objection is not present when tungsten is used. The suspender is attached to the filament by means of a suitable cement, the carbon being made of such thinness that it will allow sufficient movement in the lamp filament when switching on and off, and rupture of the filaments is thus prevented.—The Engineer.

SCIENCE NOTES.

Uruguay is said to be rich in minerals. So far this source of wealth has been almost entirely neglected. With the exception of granite quarries in various parts of the country and four gold mines at Cuñapirú, in the department of Rivera, the mineral deposits have not been worked. In the past year, however, according to a recent consular report, great interest has been manifested by mining men in the possibilities of various goldfields. Several mining engineers and prospectors from South Africa and elsewhere have reported on the prospects.

Astronomical science loses one of its most eminent workers in the death of the eminent director of the Paris Observatory, M. Maurice Loewy, which occurred suddenly on the 15th of October at the age of 75. For a long time past M. Loewy has been identified with the work of the observatory, having been associated with the celebrated astronomer Leverrier at the outset of his career, and in later times he was especially engaged in observations upon the planets, the moon, about whose formation he advanced some new theories, meteorites, and the determination of longitude. Born at Vienna in 1833, he was naturalized in France in 1864, and was made a member of the Académie des Sciences in 1873. Having the post of sub-director of the observatory in 1878, he assumed the position of director ten years later. To him is due the construction of the great equatorial, for which he designed the plans. Within the last few years he was occupied with lunar photography and the preparation of the photographic atlas of the moon.

Some thirty years ago Lord Rayleigh made an extensive series of experiments in order to ascertain more precisely what are the capabilities of the ears in estimating the directions of sounds. It appeared that even with pure tones, the discrimination between right and left sounds could be made with certainty, and without moving the head. But with pure tones no pronouncement could be made if the sounds were situated in front or behind the observer. With the speaking voice, front and back could often be distinguished. The discrimination between right and left is usually supposed to be explicable by the greater intensity of sensation experienced by the ear which lies nearer to the sound. Lord Rayleigh considers that this explanation is adequate when the pitch is pretty high. But for a frequency of 256, the difference in intensities at the two ears is far from conspicuous, and at 128 it is barely perceptible. Theory indicates that the difference of intensity in the case of a pitch of 128 should be decidedly less than one per cent of the whole; yet the discrimination between right and left is as easy as before. Lord Rayleigh has recently returned to the question, commencing with a frequency of 128. Every precaution was taken to remove bias. The observer, either on his feet or more conveniently seated upon a rotating stool, is turned round until he loses his bearings. It is often necessary to cover the eyes with the hands as well as to close the eyelids, in order sufficiently to exclude the light. With different observers it was a moot point whether there was any balance in intensity for the source on the same side as the open ear in the case of frequency 128, while with 256 the difference was marked. Thus there are cases where it is impossible to attribute the correct judgment of direction to the relative intensity. The only other circumstance in the case of pure tones is the relative phase of the notes at the two ears. The sound reaches one ear later than the other, and the same statement holds good for all the successive condensations in the wave train. If this phase difference is admitted as a source of the power of discrimination, it must be limited to sounds of low pitch. The retardation for frequency 512 would be about half a period, for 1,024 a whole period, and so on; and since any number of whole periods are equivalent to no retardation at all, such a retardation can be of no avail as an explanation. In testing the applicability of the explanation to low pitches, two forks, electrically maintained, were placed in separate rooms, and the sounds led through composition tubing to the observer's ears. The forks were slightly out of tune, so there was a progressive increase of phase difference between the notes. The time for the retardation to change through a whole period was about five seconds. On the very first trial, Lady Rayleigh and Lord Rayleigh at once experienced a distinct right and left effect, the sound appearing to transfer itself alternately from the one side to the other. The beat may be slowed down until it occupies forty or even seventy seconds, with the same results. Further experiments showed that the transitions between right and left effects correspond to agreement and opposition of phase. When the vibration on the right is the quicker, the sensation of right follows agreement of phase and that of left follows opposition; and right and left may be interchanged in the above statement. These results, it may be repeated, apply only to pitches of 128 or under. At a pitch of 256 the effect is still present, but begins to be complicated by the differential intensity effect.

ENGINEERING NOTES.

The position of Russia as the leading producer of manganese ores is now seriously threatened by the extensive development of deposits of manganese ore in India.

A mixture of lead wool and graphite is now being successfully used for packing valves. It is used in the same manner as asbestos or other similar packing is employed. This mixture may be used for high temperatures, and has some advantages over asbestos.

For some time past, says the Naval and Military Record, rumors have been current at Portland and Weymouth that the British Admiralty intended discontinuing the torpedo range in the Roads. The range has been found inadequate to meet the demands of the improved torpedo, the range of which has been greatly increased by the recent hot-air invention, and the authorities have decided to take the range elsewhere—probably to Loch Long, in Scotland. The decision of the Admiralty has caused a good deal of regret at Weymouth, as it is felt that another blow has been struck at the local torpedo industry, which is suffering from acute depression just now.

The Germans are carrying out a plan in the interior of Africa to divert to the German ports on the east coast the trade flowing over the Uganda Railway, a line subsidized by Great Britain. They have already built a railway from Pangani, on the coast, to the base of Mount Kilimanjaro, on the border of British East Africa. Swamps on the one side and mountain ranges on the other prevent this line from being continued to Lake Victoria Nyanza, but to overcome this difficulty a canal through the swamps from a deep inlet of the lake to the railroad is being dug. The canal will be sixty miles long and the overflow from it will be carried to the sea by the River Ruvu. Ordinarily such a work might not be considered a menace, but experts believe that, in the case of Lake Victoria Nyanza, it will do great damage to British interests. The lake has been falling for years, some of the rocks which now stand sixty-five feet above its level having not so many years ago been submerged. If further drained by this canal it is said that it will not be long before the steamers which now ply on the lake will no longer be able to navigate its waters. Some even say that any interference with the waters of the lake will affect the Nile supply, the lake acting as a natural reservoir to be drawn upon in the years when the Abyssinian rains fail.

We recently drew attention to the interesting development that is to be carried out by the International Marine in the construction of a liner equipped with a combination of reciprocating and turbine engines, the latter section of the equipment to operate the steam after it has been exhausted from the cylinders before passing to the condensers, with the object of securing the last ounce of power. Following this action, the Hamburg-American Line have placed an order with the Belfast shipbuilding firm of Harland & Wolff for a transatlantic liner to be engined upon the same principle. This vessel will exceed the "Lusitania" in length, being 840 feet long over all by 86 feet beam. In speed she will be of the intermediate class, similar to the "Oceanic" and sister vessels, the maximum speed desired being about 15 knots. The propelling equipment is to be carried out by Messrs. John Brown & Co., of Clydebank, the builders of the "Lusitania," with whom Harland & Wolff are now allied. The reciprocating engines will be of the quadruple-expansion type exhausting the steam into low-pressure Parsons turbines; and by means of a vacuum intensifier associated with the condenser, it is anticipated that there will be a considerably increased range of total expansion, the result of which will be an appreciable economy in the coal consumption. In fact, it is expected that this economy in fuel will approximate 12 per cent at 15 knots speed.

TABLE OF CONTENTS.

	PAGE
I. AERONAUTICS.—Record Flight of the Farman Aeroplane.—3 illustrations.....	296
II. ARCHAEOLOGY.—Pre-Roman Discoveries in Spain.—3 illustrations.....	300
III. ASTRONOMY.—Cathode Rays and the Aurora.—By C. CHREE.....	295
IV. BOTANY.—Unusual Agents of Pollination.—By G. CLARKE NUTTALL, B.Sc.....	302
V. ELECTRICITY.—Elements of Electrical Engineering.—IV.—By A. E. WATSON, E.E., Ph.D.—4 illustrations.....	293
Electric Metal-testing Pyrometer.—By WILLIAM H. BRISTOL.—3 illustrations.....	297
VI. ENGINEERING.—Eddy Current Brake for Rating Gasoline Motors.—1 illustration.....	295
Steam versus Electric Locomotives.....	298
VII. MEDICINE AND HYGIENE.—Use of Gas from a Hygienic Standpoint.—II.—By Prof. VIVIAN B. LEWIS.....	299
VIII. METALLURGY.—Calcium: Its Properties and Possibilities.....	296
IX. MISCELLANEOUS.—The Great Government Pier at the Jamestown Exposition.—By E. DABNEY TUNIS.—1 illustration.....	290
X. NAVAL ARCHITECTURE.—The Development of Armored War Vessels.—XI.—By J. H. MORRISON.—2 illustrations.....	291
XI. PHYSICS.—The Study of Flame.....	303
XII. TECHNOLOGY.—Qualities and Methods of Using Cement.—By S. B. NEWBERRY.....	290
XIII. ZOOLOGY.—The Manometric Apparatus of Fishes.—By Dr. OTTO THILO.—8 illustrations.....	301