statement of several cases bearing upon it, Garrod gives his conclusions as follows:—

1. Both erythema multiforme and erythema nodosum occasionally appear in direct association with rheumatic fever.

2. They are usually attended by some degree of arthritis.

3. The arthritis which accompanies these erythemata may have any degree of intensity, from the severe joint lesions of acute rheumatism to the slightest joint pains.

4. Collateral evidence, such as the presence of chorea or old heart lesions, is sometimes obtained pointing to the rheumatic nature of the slighter joint lesions.

5. Active cardiac lesions may accompany erythema with the slightest arthritis and probably occur sometimes in the absence of joint symptoms.

6. Erythema without joint lesions is not uncommonly met with in patients who have previously suffered from rheumatism, but in some cases there is no indication of any association with rheumatism. In a word, it appears probable that erythema multiforme and erythema nodosum are often or even usually manifestations of the rheumatic process, and their appearance should always lead us to suspect such causation. We are not, however, justified in concluding that such eruptions have always this origin, nor in concluding from their presence alone that the patient is the subject of rheumatism.

Clinical Department.

ANTISEPTIC IRRIGATION OF THE KNEE-JOINT FOR CHRONIC SYNOVITIS.¹

BY M. H. RICHARDSON, M.D.

The following cases were treated at the Massachusetts General Hospital by Dr. J. Collins Warren and myself in the autumn and winter of 1888-9. The method employed was not a new one.

Many such operations had been done abroad, especially in Germany, with such marked success that I was led to employ the same means. Chronic effusions into the knee-joint have been, in my experience, very disabling to the patient, and liable to produce disorganization of the cartilages or other serious consequences. The ordinary treatment,—compression with or without aspiration, rest, splints, and so on, has rarely been productive of a cure or even of lasting benefit.

One of the first cases I ever treated was that of a woman of about middle age, whom I first saw in the dispensary in 1877. She had an enormous effusion into the knee-joint, which ordinary treatment at her house failed to relieve. I sent her into the Massachusetts General Hospital, where she was kept in bed for a long time and the usual methods of relief tried without any benefit. The joint was aspirated and compression with sponges and fixation on splints tried without any benefit whatever. She finally disappeared. Some few years ago she again appeared at the hospital with the joints in a very much impaired condition. The lateral ligaments were so much stretched that

there was a flail-like motion to the joint. I was quite convinced by this case, among others, that a chronic effusion of any size into the knee was a very serious matter, and that more radical measures were called for if failure followed a reasonable employment of the simpler methods of treatment.

In the following case I made use of a 5 per cent. solution of carbolic acid. The same method employed by me was used by Dr. Warren in his cases. The result was extremely gratifying, and the usefulness of the joint fully restored.

CASE 1. August, 1888. Christina N., æt. twelve, had had for two years a chronic swelling of the knee which followed a blow. There was a distinct scrofulous history. When admitted the knee was greatly distended, with the usual signs of a large effusion. Under ether a large aspirating needle was introduced into the knee-joint above the patella, on the outer side. About five ounces of clear, straw-colored fluid was withdrawn, and the same amount of a 5 per cent. solution of carbolic acid introduced. After massaging the joint well, to be sure that the fluid had come in contact with every portion of the cavity, suction was again used and about eight ounces of milky fluid withdrawn. The action of the acid in coagulating the albumen contained in the effusion caused so much precipitation as to interfere quite seriously with the action of the aspirator, the fluid running slowly and with difficulty. After emptying the joint as completely as possible, the limb was placed upon a ham splint and pressure applied. On the following day there was some pain in the joint and the temperature was slightly higher. Careful examination of the urine showed that there was no absorption of carbolic acid. On the third day there was no pain, and very little fluid could be detected in the joint. At the end of two weeks she was discharged with a plaster bandage on. Two months later she was seen and examined. There was no return of the fluid, and good use of the joint.

CASE 2. Service of Dr. Warren. Irwin L. B., æt. seventeen. December, 1888. Family history good. For one year the left knee had been swollen. The joint was found to be full of fluid and there was some pain and inconvenience. Operation by Dr. Warren. One and a half pints of clear, yellow fluid withdrawn. Five ounces of 5 per cent. solution of carbolic acid injected and after a time withdrawn. Ham splint, antiseptic dressings, and elastic pressure bandage applied. The temperature rose to 102° on the third day. No evidence of absorption of carbolic acid. On the ninth day swelling of the joint much diminished. After a month there was no swelling. Good use of the limb. No signs of a return of the disease.

CASE 3. Service of Dr. Warren. Johanna A. M., æt. forty-two. January, 1889. For thirty years had had suppuration in the left knee, and for the last year and a half weakness and swelling of the right. No pain and no history of injury.

General condition of patient poor. Left knee greatly enlarged. Tibia increased in size. Ankle bent backwards. Right knee much enlarged. Bulging of the supra-patella pouches and floating patella. Operation by Dr. Warren. Needle introduced into the right knee joint on the outer side opposite upper edge of the patella, and six ounces of gelatinous

¹ Read before the Surgical Section of Suffolk District Medical Society, March 6th, 1889.

fluid withdrawn. Three ounces of 5 per cent. solution of carbolic acid injected and withdrawn through needle. Ham splint, elastic pressure, and bandage. Very little pain or disturbance followed. Nine days later bandage removed. No fluid found. Fourteenth day discharged with knee in good condition.

Although the immediate effects of this treatment are good it is too soon to be able to say that there has been a permanent cure. It is quite safe to say, however, that we may expect a permanent cure if we continue this treatment and make use of repeated aspirations should the fluid reappear. It is quite important to use a needle of considerable size because of the precipitation of the albumen in the joint-fluid by the carbolic acid. The best point to introduce the needle is just above the patella on the outer side through the fibres of the vastus externus.

Although this procedure is extremely simple, yet it ought not to be employed indiscriminately. I should never use it till the ordinary means had failed, and then only with the greatest care, especially as to cleanliness and asepsis.

Reports of Societies.

SURGICAL SECTION OF THE SUFFOLK DISTRICT MEDICAL SOCIETY.

GEORGE H. MONKS, M.D., SECRETARY.

MEETING, Wednesday evening, March 6th, Dr. E. H. BRADFORD in the chair.

Dr. C. B. PORTER showed a patient of his, a little girl, upon whom he had very successfully operated for

EXTROVERSION OF THE BLADDER.

Mr. Fisk (of the firm of Fisk & Arnold), introducing the subject of

AMPUTATIONS AND ARTIFICIAL LEGS,

showed, by invitation, specimens of artificial legs as made by them and explained their special features. He said that for the manufacture of artificial limbs he much preferred wood to rubber. He also gave his ideas as to the most satisfactory stumps for the attachment of these limbs.

He said that by far the best place for an amputation of the leg to be made, so far as the satisfactory attachment of a false limb was concerned, was at a point about six or seven inches below the patella.

A satisfactory result, with a good, smooth gait, could be guaranteed in almost every case. If the stump is longer than this there is almost always some trouble with the circulation. (Mr. Fisk here introduced to the section an assistant of his, who wore two artificial legs upon stumps amputated about six inches below the patella. This man could walk, run, go up and down stairs, and with surprisingly little peculiarity in his gait.) Amputation at the knee-joint was satisfactory, if a good thick flap could be got for a pad; but this was difficult, and the usual result was great inconvenience from corns and excoriations. Amputations in the vicinity of the ankle-joint ought likewise to be made so as to ensure thick pads, else great discomfort would result. The tendo Achillis frequently gives trouble, also, by subsequent contraction, drawing the heel

well up, stretching the cicatrix, and changing the point of pressure. While an amputation in the neighborhood of the ankle-joint is in certain cases the best for a patient of the poorer classes, who needs a useful limb only, a patient who can afford to pay for appearance and comfort would be best suited with an amputation performed at the point of election, six or seven inches below the patella. As for amputations of the thigh, an artificial limb can be satisfactorily fitted to almost any stump. Mr. Fisk has even fitted a limb satisfactorily to a thigh stump, where the amputation was made two inches below the perineum. As a rule three to five months must pass after an operation before the stump is in a condition for a false leg to be worn. Until the apparatus is adjusted it is well to keep down œdema and preserve circulation in the stump by bandaging.

Dr. J. C. WARREN said that Mr. Fisk had several times kindly exhibited his collection of false legs before the class at the Medical School at the lecture on amputations. Dr. Warren said that it was his custom to show, on this occasion, many cases of stumps resulting from the various amputations of the lower extremity, from the foot up. He said that the examination of these cases had considerably influenced his views as to the desirability of the different amputations, and that he had now practically the same opinion as to ankle-joint amputations as that just expressed by Mr. Fisk, viz., that certain ankle-joint amputations were the best for the poor, whereas amputation at the point of election (*i.e.*, about seven inches below the patella) was much to be preferred for those who could afford to procure a good artificial limb.

Dr. M. H. RICHARDSON reported several cases of CHRONIC EFFUSION INTO THE KNEE-JOINT TREATED BY THE INJECTION OF A 5 PER CENT. SOLUTION OF CARBOLIC ACID.¹

Dr. W. INGALLS related eight cases, in each of which aspiration followed by compression and absolute rest resulted in recovery. The amount of fluid withdrawn ranged from three to seven or eight ounces. Three of the cases were acute.

Dr. E. W. CUSHING said: It may be of interest in this connection to state that about three years ago I had a case of distention of the hip-joint in a young boy, with well-marked symptoms. At that time I was associated with Dr. Marey, who had had considerable experience in the injection of carbolic acid solution into the knee-joint.

I aspirated the hip-joint and drew off about one ounce of turbid serum, and then injected a 3 per cent. solution of carbolic acid and let it run out, and then again, and let it run out again. The immediate results of this were excellent, and the disease, to the best of my knowledge, was stayed. I think, however, that a 2½ per cent. solution will accomplish as much as a stronger one, without the disadvantage of the coagulation. A coagulum protects bacteria rather than harms them.

THE CHAIRMAN: This irrigation of joints, as Dr. Richardson has said, is not a new one, but I think he is one of the first to apply it here in Boston, though Dr. Marey antedated him. It has been done quite extensively in Germany and to a

¹ See page 488 of this Journal.