competition announced by the "Society for the Prevention of Blindness in London," and to which the prize of 80*l.* offered by the Society was awarded. The book may be described as containing a succinct exposition of the chief causes of blindness, and an endeavour to render them intelligible to non-medical readers; the object being to obtain the cooperation of the public in the removal of these causes, in so far as that desirable end may be attained by improved hygiene, and by a better knowledge of the most favourable conditions of ocular work.

The causes of blindness which may fairly be said to be thus remediable, even including under blindness high degrees of defective vision, are two in number—namely, the purulent ophthalmia of new-born infants, and the progressive short-sight which is not uncommon in schools. The former is a disease which might frequently be prevented, which is always curable if treated in good time, but which, if neglected, is almost certain to destroy the sight; and to neglect of its early stages among the poor, and in remote country districts, probably four-fifths of the blindness which occurs among children in this country may be ascribed. Several months ago the Ophthalmological Society of the United Kingdom, moved thereto by Dr. McKeown of Belfast, sent a deputation to the Home Secretary to call the attention of the Government to the dangerous character and the easy curability of this affection, and to urge that steps should be taken, through the instrumentality of the Registrars of Births, to diffuse a more general knowledge of the importance of early treatment. Partly through the opposition of the Registrar-General, the deputation met with no encouragement; and the information given by Dr. Fuchs is therefore as opportune as it is valuable, and might with great advantage be communicated to the poor by clergymen, schoolmasters, and others. It may be said, however, that many of his recommendations apply chiefly to countries in which the employment of midwives is more general than in England.

The progressive short-sight of the educational period is a matter which has lately attracted much notice in all civilised countries, and Dr. Fuchs has nothing to say concerning it which is original. He presents, nevertheless, a brief and convenient summary of the facts, and a good description of the methods of school lighting and fitting which are most to be commended. This part of his volume may be studied with great advantage by any teachers and managers to whom the more systematic treatises upon the subject are either unknown or inaccessible. The book contains one serious error, which, in the English version, has been slightly modified by a mistranslation. Dr. Dudgeon writes, with reference to the provision for instruction about eye diseases in the medical schools of Great Britain and Ireland—"There are eye departments in all the large hospitals, but as a rule no regular lectures on ophthalmology are delivered." The word rendered "ophthalmology" is in the original not "*ophthalmologie*," but "*augenheilkunde*," and the correct translation would be "the treatment of diseases of the eyes." On this subject, that is to say, upon so much of ophthalmology as has any direct bearing upon the duties of the medical practitioner, systematic lectures are delivered in every medical school in the United Kingdom; and it is difficult to believe that

the translator could have been unacquainted with the fact. "Ophthalmology," of course, takes a much wider range, and embraces branches of optics and of physiology with which the practitioner, unless a specialist in eye disease, has neither time nor reason to concern himself.

OUR BOOK SHELF

Among the Rocks round Glasgow: A Series of Excursion-Sketches and other Papers. By Dugald Bell. Second Edition. (Glasgow: Maclehose, 1885.)

THIS volume furnishes a good example of what a busy man can do in his few intervals of leisure. The volume is mainly based on notes of excursions kept by the author while acting as secretary to the Glasgow Geological Society. It affords a fairly accurate idea of the geological structure of the country round about Glasgow, and of the principal features of interest which the rocks of the district present. The excursions extend as far as Stirling, take in the course of the Clyde and not a few districts on its banks. Many of the papers are pleasant reading; and even geological specialists may find something in the pages to interest and inform.

Three Martyrs of Science of the Nineteenth Century. Studies from the Lives of Livingstone, Gordon, and Patterson. By the Author of "Chronicles of the Schönberg-Cotta Family." (London: S.P.C.K., 1885.)

THE author of this volume tells the story of these three remarkable lives very pleasantly and instructively, more, however, from the religious than the scientific standpoint. A very fair account is given of the work accomplished by Livingstone in Africa, though the author does not seem to be quite aware of the value of the geographical work accomplished by Gordon on the Upper Nile.

LETTERS TO THE EDITOR

[*The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts. No notice is taken of anonymous communications.*]

[*The Editor urgently requests correspondents to keep their letters as short as possible. The pressure on his space is so great that it is impossible otherwise to insure the appearance even of communications containing interesting and novel facts.*]

Upper Wind Currents over the Equator

THE importance of an accurate knowledge of the general circulation of the atmosphere over the equator is so obvious and so little known that the following observations, taken on a voyage from Aden to Australia in February, 1885, will be of interest:—

Over the north-east monsoon, north of the line, the surface wind was east-north-east, while the low clouds came from due east. No high cirrus was ever seen.

In 2° N. lat. the surface wind lowest clouds came from N.N.E., the next layer of cirro-stratus from E. rather fast, while the highest cirri drove very slowly from E.S.E.

In about 1° S. lat. the surface came from N.W. (the N.W. monsoon), small flecks of low cloud from N.E., while some high cirri moved from E. at a moderate rate.

In 5° S. lat. the surface wind still blew from N.W., the lowest cumulus moved from N.N.W., the next layer of cirro-stratus from N., while a still higher layer of cirrus came slowly from E. or E.S.E.

In 10° S. lat. the surface wind came still from N.W., and the clouds at moderate altitude from S.E.

In the "Doldrums," which we only reached in 13° S. lat., the surface wind was from S. and the clouds from S.E.

After we entered the S.E. trade, while the wind came from S.E., the clouds drove from S., and when about 25° S. lat. the trade drew into E.; the clouds came from S.E.

The relation of upper to surface winds in the N.E. monsoon is just what might have been expected; but the discovery of an

easterly current over the N.W. monsoon and of an upper current over the S.E. trade, more southerly than the surface wind, is not only altogether new, but also quite anomalous.

In Australia, and the Southern Hemisphere generally, the upper current over a N.W. wind is from about W. and over a S.E. wind from about E.

On my way home I ran a section across the Atlantic from Rio to Teneriffe, but the absence of cirrus prevented any satisfactory determination of the upper winds in that region.

The matter is, however, so important that I start again in a few days for the hurricane region of Mauritius, where I hope to observe one of these exceptional cyclones. Then I hope to repeat a section of the Indian Ocean between Mauritius and Bombay, and afterwards, if all goes well, to get some sections in the Pacific to see what the meaning of this curious discovery may be.

RALPH ABERCROMBY

21, Chapel Street, S.W., October 26

The Hellgate Explosion and Rackarock

THE statement in NATURE of the 15th inst. (p. 575) that rackarock is "blasting gelatine" or "nitroglycerine with compressed gun-cotton," is incorrect. Rackarock is simply powdered potassium chlorate, impregnated with an inexpensive oily combustible, such as coal-tar oil, and is one of my safety-explosives, which I discovered in 1870, patented in England, April 6 and October 5, 1871, and described more fully in the *Journal of the Chemical Society* for August, 1873, under the title: "On a New Class of Explosives, which are non-Explosive during their Manufacture, Storage, and Transport."

I am not responsible for the quaint name which the Americans have been pleased to give to my child.

As the so-called "rackarock" is not very sensitive or easy to explode, it requires a strong primer or detonator to set it off. This property, which I have fully discussed and particularly accentuated in my paper of 1873, explains why Gen. Newton, the Chief Engineer of the Hellgate mine, took the precaution of placing as a primer such a powerful charge (33 tons) of expensive dynamite on the cheaper charge of the potassium chlorate mixture (107 tons), a precaution carried here perhaps a little too far.

Still it is satisfactory to see that my safety-explosive performed the main part of the labour and rendered good service in the advancement of the works of peace.

H. SPRENGEL

Savile Club, 107, Piccadilly

[We are very pleased to insert Dr. Sprengel's correction as to the composition of "rackarock." Up to the time of our notice about the explosion going to press the only information we could obtain was that it was the same substance as blasting gelatine, but with a less portentous name.—ED.]

An Earthquake Invention

IN your number for October 15 (p. 573) your numerous scientific readers will be interested to find a pretty long letter under the above heading from so able a seismologist as Prof. John Milne, of Tokio, Japan. Yet, his invitation notwithstanding, I must decline any discussion with him, either about my old letters which he refers to, or his own much changed opinion on their subject, since the occasion for my writing them occurred.

Those points, Mr. D. A. Stevenson, who is also invited, may, or may not, take up. My letters were impersonal, and dealt only with a British Association Report. I desire also to continue to keep them strictly to that, even to the very words of the particular Report as given forth to the world with all the usually unquestioned authority of that mighty Association, in their B.A. volume for 1884, p. 248, Section entitled "Experiments on a Building to resist Earthquake Motion."

C. PIAZZI SMYTH

15, Royal Terrace, Edinburgh, October 16

On the Behaviour of Stretched Indiarubber when Heated

SOME time ago (NATURE, vol. xxv. p. 507) you permitted me to express a doubt as to the invariable success of an often-quoted experiment with cylinders of bismuth and iron, intended to illustrate some relations between specific heat and thermal conductivity.¹ I regret that I have made further progress along

¹ Perhaps I may mention in passing that if lead is substituted for the bismuth the experiment succeeds perfectly, as theoretically it should do.

the evil road of scepticism. I should like, in fact, to ask whether it is absolutely true to say without qualification, as is done in many text-books, that india-rubber (when stretched) forms an exception to the general law that the volume of a body is increased when the temperature is increased. The usual form of the experiment supposed to prove this is well known: a piece of india-rubber tube or cord is stretched by a weight connected with a long light index-lever, and it is shown that when heat is applied the india-rubber gets decidedly shorter.

I have always had some hesitation in showing and explaining the result of the experiment in the above way, especially as I could not find any proof given that the contraction in length was not compensated, or more than compensated, by an expansion in other directions (like that of a worm in its creeping progress, or of a dry rope when wetted). I had, in fact, lately arranged an apparatus for determining the coefficient of expansion of india-rubber, whether positive or negative, when I found that the subject has been very fully investigated by Dr. J. Russner, of Chemnitz (see Carl's *Repertorium* for 1882, pp. 161 and 196).

His results are briefly these:—

(1) That india-rubber (of which several kinds were examined) has without exception a definite coefficient of expansion which is always positive; experiments made at temperatures varying from 0° to 53°·4 gave, for its value at 10°, 0·000657; at 30°, 0·000670.

(2) That india-rubber in a stretched state expands to the same extent as when it is not stretched. No point of minimum density was observed, such as Puschl supposed to exist.

(3) That the apparently anomalous behaviour of stretched india-rubber when heated is simply a case analogous to those of anisotropic crystals, which expand to different extents in different directions. Iceland spar, for instance, as Mitscherlich showed, actually contracts in a direction at right angles to its principal axis when heated, although its volume is, on the whole, increased.

Although ordinary india-rubber is, of course, isotropic, yet when stretched it becomes anisotropic, as may easily be shown by stretching a piece until it is semi-transparent, and placing it between crossed Nicols; the direction of the strain lying at an angle of 45° with the plane of polarisation. Distinct colours, as with a selenite film, will be seen, varying from red to blue with the amount of strain.

The fact that india-rubber becomes hot when stretched, and especially if stretched and allowed to contract several times in succession, may perhaps be accounted for by molecular friction. It would almost seem, then, that in the account given in many books the truth, as well as the india-rubber, has been slightly "stretched."

H. G. MADAN

Eton College, October 23

The Resting Position of Oysters

IN carrying out a series of experiments on the artificial breeding of oysters in my private aquaria, I noticed that the young oysters born in the tanks rested on the flatter shell when they obtained a flat surface, such as a tile, to adhere to, but when I so arranged that they had irregular surfaces to deal with, such as little bundles of twigs, some adhered one way, and some the other. But where young oysters, nearly two years old, were moved from their original supports, and were compelled to find new ones, they selected the flat shell to rest upon in every instance, except where they were placed on sand, in which case they rested on the convex shell, in order apparently to avoid clogging the mouth of the shell with sand. Is it not possible from these observations that adult oysters vary their position according to the nature of the ground they are on. I have seen adult oysters on muddy ground lying on the convex shell, while where adhesion to a flat surface could be obtained, they were all on the flat shell, and pectens are dredged with Balari and other growths on the flat shell in some instances, and on the convex shell in others, principally, however, on the latter.

H. STUART-WORTLEY

South Kensington Museum, October 23

The Value of the Testimony to the Aurora-Sound

I HAVE read with much interest the descriptions of this sound as given by Dr. Sophus Tromholt's correspondents in NATURE of September 24. I was, however, struck by the similarity of these descriptions to the well-known phenomena of *tinnitus*