details his own experiments which were made on the popliteal nerves of dogs. The results of the reunion after division, and suturing in various combinations, such as the central end of the internal popliteal to the peripheral portions of both internal and external popliteal nerves, are investigated both from an anatomical and physiological standpoint. The author summarizes his conclusions as follows: "It is possible to function the two opposing groups of muscles by a single nerve, which previously supplied one group only. Second, when the central end of one nerve is joined to the peripheral ends of two nerves there are many more fibers in the peripheral nerves than in the central nerve, so that the nerve fibers in the proximal trunk divide on going to the distal trunks. Third, in some cases, at least, some of the branches from one nerve fiber go to supply one set and others the opposing set of muscles. This may prevent very delicate movements being restored. Fourth, after this form of suturing the arrangement of the nerve fasciculi in the peripheral nerves is considerably altered.

C. D. CAMP (Philadelphia).

CEREBRAL SYMPTOMS IN MEASLES. . Guy J. Branson (British Medical Journal, April 29, 1905).

The case appeared to be a simple, uncomplicated attack of measles, but on the next day after the appearance of the rash there were noted frequent twitchings of the face and limbs, and the day following a severe general convulsion accompanied by unconsciousness. There were no paralyzes or localizing symptoms, but the attack was followed by intense restlessness and cerebral irritability. All the symptoms completely disappeared as the measles rash and fever cleared up.

C. D. CAMP (Philadelphia).

DISSEMINATED SYPHILITIC ENCEPHALITIS. Albert M. Barrett (The American Journal of the Medical Sciences, March, 1905).

A case of brain syphilis occurring in the secondary stage of the disease. The symptoms were those of gradually progressive dementia accompanied by ocular symptoms and general spasticity of the limbs, which was more marked on the left side. Death occurred in coma, and at autopsy the greater affection of the left side was explained by the discovery of a small hemorrhage in the right side of the pons. A detailed histological examination of the brain revealed areas of acute localized encephalitis not apparently connected with the meningitis, which was also present. There was also found a typical diffuse encephalitis, syphilitic arteritis, etc.

C. D. CAMP (Philadelphia).

REGENERATION OF NERVES. Mott, Halliburton and Edwards (Proceedings of the Physiological Society, March 19, 1904).

These authors have followed their studies in nerve degeneration with others in nerve regeneration. Early in the degeneration of the nerve fibers the neurilemma cells multiply, and later seem to assist the phagocytes in the removal of the broken-up myelin drops. The neurilemma cells are found in elongated strands concealing the newly formed axis cylinder. The authors conclude that this cylinder proceeds alone from the central end of the divided nerve, and that the peripheral cells simply act as a scaffolding for it. They call attention to the fact that histological evidence is not sufficient to establish the identity of a nerve fiber; that a strand which looks like one is not really such unless capable of responding to excitation and conducting nerve impulses. The peripheral theory of nerve regeneration has been held to be supported by clinical evidence, on the ground that after freshening up and suturing a divided nerve, sensation returns within forty-eight hours. A case is noted, however, in which the patient stated shortly after the operation that he was able to feel, but these sensations disappeared again in a short time and feeling did not really