Clinical Notes :

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A NOTE ON DECOMPOSITION OF CHLOROFORM, AND SICKNESS.

BY DAVID NEWMAN, M.D.,
SURGEON TO THE GLASGOW ROYAL INFIRMARY;

AND
WILLIAM RAMSAY, PH. D., F.R.S.

EARLY in the history of chloroform it was noticed that the agent had a tendency to undergo spontaneous decomposition on exposure to air and light, and although within recent years great care has been taken by manufacturers of the drug to produce as pure an article as possible, nevertheless, the fact remains that chloroform is not stable and rapidly deteriorates as an anæsthetic. When freshly distilled chloroform is administered, the vapour is easily inhaled, having a sweetish and not unpleasant taste and odour. It produces little or no irritation in the air passages, little excitement of the patient is observed, and if the agent is carefully given sickness, fainting, and irregularity of the pulse and respiration seldom occur during its administration. But if chloroform is kept in a bottle containing air, exposed even to feeble light, changes are produced in the drug which render it less suitable as an anæsthetic. The change is due to the formation of carbonyl chloride and hydrogen chloride, according to the equation $\text{CHCl}_3 + \text{O} = \text{HCl} + \text{COCl}_2$. These substances render the vapour slightly pungent in odour, and on inhalation it produces irritation in the air passages. The period of excitement of the patient is more marked and the liability to sickness during and after anæsthesia is greater.

During the last six months the effects of freshly distilled chloroform and of chloroform supplied in the ordinary way have been contrasted, greatly to the advantage of the former. It has been found, however, that distillation is not necessary, as by shaking the chloroform with slaked lime and filtering the irritating products of decomposition are eliminated, as, indeed, theory would predict; for the hydrochloric acid is, of course, at once acted on by the lime, giving calcium chloride and water, and the carbonyl chloride, to the presence of which the sickness is probably to be ascribed, is resolved into carbonate and chloride of calcium, thus— $\text{COCl}_2 + \text{Ca}(\text{OH})_2 = \text{CaCO}_3 + \text{CaCl}_2$. In illustration the three following cases may be mentioned:—

CASE 1.—A man was operated on three occasions. For the two first the chloroform given had been in the wards for not more than ten days, and was kept in a well-stoppered bottle away from direct light; but the bottle contained some air. Each operation lasted fully an hour, and on both occasions the pulse became very irregular and feeble so as to necessitate artificial respiration. While recovering from the effect of the chloroform he was very sick, and the sickness continued for two days after the first operation, and for three days after the second. The third operation lasted for one and a half hours. Chloroform which had been kept with slaked lime in it and filtered before use was employed. The patient, although much feebler through disease, remained well during the operation, and there was no sickness for twelve hours, when he suddenly became sick, but this passed off in a short time.

CASE 2.—A woman had been under chloroform four times and she was invariably very sick while coming out and afterwards, the sickness lasting for twenty-four hours or more. When chloroform treated with lime was used, on coming out of the anæsthetic sickness occurred for a few minutes only.

CASE 3.—A man was under chloroform four times. On the first and third chloroform as supplied by the druggist was used, and it was not more than twelve days old. After both operations there was severe sickness, lasting for thirty-six hours. On the second and fourth administrations freshly distilled chloroform was employed with no sickness resulting.

As a matter of experience it was found that freshly prepared chloroform is more reliable as an anæsthetic, and is less likely to produce sickness during or after administration, and the risks of anæsthesia are probably increased by the decomposition of chloroform when kept for any considerable time exposed to air and light. Even with fresh, chloroform sickness may occur if the patient is allowed to come out during the operation, but the quantity of acrid mucus is very small in amount as compared with what is seen when older chloroform is given. Perfectly pure chloroform is said to be a much less stable body than chloroform containing alcohol. As an instance of the instability of pure chloroform it may be mentioned that a sample left in a half-full stoppered bottle from July till October, 1896, smelt as pungent as hydrochloric acid, gave with baryta water a copious precipitate of carbonate, and was absolutely irrespirable.

Glasgow.

UNUSUAL CASE OF CARCINOMA OF THE BREAST.

BY W. S. CRAWFORD, B.A. CANTAB., F.R.C.S. EDIN.,
SENIOR ASSISTANT SURGEON, LIVERPOOL CANCER AND SKIN HOSPITAL.

IN the beginning of September, 1892, a fresh-complexioned, thin woman, aged thirty-one years, came under my care at the Liverpool Cancer and Skin Hospital. She was unmarried. Some months before admission the patient had received a blow on the right breast which caused her a good deal of pain for some time. She did not notice any swelling. Three months after receiving the blow she found a lump in her right breast about the size of a marble, situated above and to the outer side of the nipple; it was freely moveable and quite free from pain. On the patient's admission to hospital, some four months later, the lump had attained the size of a tennis ball. It was very hard and adherent to the skin. The nipple was retracted. The axillary glands were hard and enlarged. It was a well-marked scirrhus of the breast. The breast was removed and the axilla cleared. The sections were found to be hard, spheroidal-celled carcinoma. On Nov. 25th, 1896 (four years later), the patient again presented herself at the hospital. Her general appearance was but little changed. She gave a history of having wrenched her right arm and hurt her right breast eight months before. There was no lump of any kind then. The present local condition is as follows. There is a large oval-shaped tumour distending the skin, which is ulcerated in a circular patch two inches in diameter over the most prominent part of the growth. The ulcerated area is concave; it bleeds readily on the slightest interference. Towards the axilla a flattened portion of growth is felt extending from the main mass and firmly adherent to the pectoral muscles, but not certainly to the bony wall of the chest. About two inches lower down than the main mass there is a small area somewhat discoloured about the size of a penny, intensely hard, and two or three small nodules in the immediate neighbourhood, evidently recurrences in the superficial lymphatics. The right armpit is clear, but the left contains a mass of enlarged and undoubtedly cancerous glands. The main recurrence commenced, according to the patient's description, immediately above the old cicatrix. It now involves the scar and the tissues on each side.

Remarks.—Her father's sister died from carcinoma of the breast which followed an injury. The patient's primary growth, as well as recurrence, followed definite injuries. In so young a patient it was pleasing to find so long an immunity period—three and a half years.

Liverpool.

NOTE ON A CASE OF RUPTURE OF THE HEART.

BY CECIL ROBERTSON, M.B., C.M. ABERD.

CASES of rupture of the heart from external violence without penetration of the chest are by no means of frequent occurrence. Erichsen, in the last edition of his "Science and Art of Surgery," says that Gamgee was only able to collect twenty-two published cases. In the majority of these cases the blow was directly over the cardiac region. For two reasons I think the following case is not unworthy of being recorded—(1) the small number of recorded cases; and (2) that the heart in this case was ruptured by indirect violence.