initely suspended, I found a substitute in horticulture; and am confident that I owe it in good measure to the kindly influence of that gracious pursuit that the demon in the brain was gradually soothed into comparative quiet. In 1861 I was able, with frequent interruptions, to take up my work again. At the same time there was such amendment as regards sight that I could bear the sunlight without blinking and read for several minutes at once without stopping to rest the eyes, though my chief dependence was still in those of others. In 1865 "The Pioneers" was finished, and the capacity of work both of brain and eye had much increased. "The Jesuits" was finished in 1867; "The Discovery of the Great West," in 1869; "The Old Régime," in 1874; and "Frontenac," in 1877. "Montcalm and Wolfe," which involved more labor, was not ready till 1884.

While engaged on these books I made many journeys in the United States and Canada in search of material, and went four times to Europe with a similar object. The task of exploring archives and collecting documents, to me repulsive at the best, was, under the circumstances, difficult, and would have been impossible but for the aid of competent assistants working under my direction.

Taking the last forty years as a whole, the capacity of literary work which during that time has fallen to my share has, I am confident, been considerably less than a fourth part of what it would have been under normal conditions. Whether the historical series in hand will ever be finished I do not know, but shall finish it if I can.

Yours faithfully,

F. PARKMAN.

JAMAICA PLAIN, 28 Oct., 1886.

Clinical Department.

REPORT OF CASES FROM THE SURGICAL SERVICE OF THE CHILDREN'S HOSPITAL.

BY DRs. H. L. BURRELL, H. W. CUSHING AND J. S. STONE.

Reported by Dr. W. C. Howe.

1. Simple papilloma of abdominal wall; 2, scalp wound: concussion; 3, branchial cyst; 4, bullet wound of left chest; 5, abscess of abdominal parietes; 6, congenital deformity of hands and feet.

SIMPLE PAPILOMA OF ABDOMINAL WALL.

O. M., a boy of eleven years, admitted Feb. 5, 1903. Family history negative; previous history negative. At birth showed a minute verrucous growth on abdominal wall, which gradually grew to size of a pea. Four weeks ago it was irritated by the child's finger-nails, some bleeding occurred and it has since then increased in size (Fig. 1). Physical examination: Well developed and nourished. Marked glandular enlargement in neck, glands size of beans. No glands in axillæ; few in groin. Heart sounds regular, normal, no murmurs. Lungs show numerous wheezy râles, scattered throughout front and back; percussion normal. Abdomen negative. Upper and lower extremities normal. On abdominal wall, 2 inches below lower end of sternum, is a reddened area $1\frac{3}{4}$ inches long, $1\frac{1}{2}$ inches wide. In the center of this area is a verrucous growth $\frac{3}{4}$ inch long, $\frac{1}{2}$ inch wide, elevated $\frac{1}{4}$ inch. This growth is very vascular and bleeds easily; not painful. Operation: Dr. Cushing: The papilloma, with the surrounding skin at its base, was removed. The growth evidently did not extend below the skin. Slight

bleeding. Wound closed with a subcutaneous catgut suture. Crêpe lisse dressing. Bandage. Sixth day: The wound was dressed and found closed by first intention. Linear cicatrix firmly healed. No bulging on cough. Tenth day discharged well. Pathological report from Dr. H. A. Christian stated that the growth was a simple papilloma with lymphoid and plasma cell infiltration.

FIG. 1.—Simple papilloma of abdominal wall.

tion. We are also indebted to Dr. Christian for the photomicrographs which follow (Fig. 2 and Fig. 3). This case is of interest in that it had the gross appearance of an epithelial carcinoma. It was a simple papilloma which had increased in size as the result of irritation.

SCALP WOUND: CONCUSSION.

F. K., a boy of ten years, admitted March 30, 1903. Family history negative. Previous history: Diphtheria six years ago; following this there has been internal strabismus of right eye. There has also been paralysis of right arm since birth. Present illness: At 5.30 p.m. to-day, boy was struck on head with stone about size of a fist. At 6 p.m. was seen by a doctor, who cleaned up the wound, put in one suture and dusted wound with iodoform. While being dressed, the boy became drowsy and could be aroused only with difficulty. He was brought to the hospital at 9 p.m. Physical examination: Well developed and nourished. Pupils semi-dilated, left slightly larger than right, both react to light. Internal strabismus of right eye. Pulse regular, 76, good volume and tension, alike in the two wrists. Temperature 98.8°. On posterior aspect of head, at base of left parietal bone, is a contused wound 1 inch long. No depression or fracture can be made out. Boy not drowsy; is clear mentally. Heart, lungs and abdomen negative. Knee jerks normal. Right arm 21 inches, left arm 23 inches in length.

No muscular atrophy. The wound was cleaned with corrosive solution: corrosive dressing. On third day the wound was dressed; it was found to be clean. No symptoms. On sixth day the boy was up and about the ward. He showed no symptoms. On tenth day the patient was seen by Dr. W. N. Bullard, who made diagnosis of obstetrical paralysis, right arm. The patient was discharged well.

Injuries of this description in children are not uncommon, and unless one has seen the clinical picture of concussion, with pupils semi-dilated, one slightly larger than the other, internal strabismus of one eye, one might be led to localize a cerebral injury and operate. This patient, immediately following the injury, was mentally clear; but while the wound was being dressed stupor, a serious symptom, appeared. We have seen instances in which convulsions occurred which were so alarming as to cause the attending surgeon to operate.

BRANCHIAL CYST.

T. S., a girl of seven years admitted April 6, 1903. Family history and previous history negative. Present illness: When two years of age, the mother noticed a small lump, size of a pea, on left side of neck. This lump ruptured spontaneously and discharged. The patient was operated four years ago at another hospital by an incision and curetting. Two weeks ago the tumor reappeared and began to discharge a thin, purulent fluid.

FIG. 4. — Branchial cyst, showing opening of sinus just to left of median line.

Physical examination: Situated on anterior aspect of neck, at the level of the thyroid cartilage and at the apex of the left submaxillary triangle, is a raised reddened area about the size of a pea, covered with a slight crust in the center. It is superficial and freely movable and not tender. It has a deep attachment running in toward the median line (Fig. 4).

April 13, operation, Dr. Burrell. An elliptical incision was made circumscribing this raised, reddened area and a dense, tubular sinus found which extended downward to the wing of the left thyroid cartilage. The sac was dissected free from the cartilage. In removing the tubular sac, the dissection opened the crico-thyroid membrane, which was sutured with fine silk. The dissection was difficult, and to remove the whole of the branchial cyst it was necessary to recognize clearly the difference between the cyst wall and the adjacent tissues. The separated muscular tissue and fascia were sutured with catgut and the skin was closed with silkworm-gut sutures. In a week's time the wound was united and the patient was discharged well. In removing a branchial cyst it is necessary to dissect out the continuity of the sac. Otherwise there will be a relapse from the secreting membrane that is left from the incomplete removal.

BULLET WOUND OF LEFT CHEST.

C. G., a boy ten years of age, admitted March 23, 1903. Family history and previous history negative. On March 20, 1903, the patient was one of a band of "white settlers" who were defending a town on Cape Cod from the attack of "Indians;" the Indians ranged in age from eight to twelve years; both the white and red men were armed for the most part with bows and arrows, but the Indians were fortunate in the possession of one 22 cal. revolver. The attacking party was rapidly being repulsed, until their solitary firearm, which throughout the conflict had been discharged over the heads of the whites, was turned against the patient, at a distance of about six feet. The bullet entered his left chest. The wound was dressed with gauze and corrosive sublimate; and was not probed by the attending physician. In the afternoon he had considerable pain in the left chest. No evidence of hemorrhage. Temperature 99.5. Pain on respiration. Morning of 21st, temperature normal, respiration 30, pulse 110; evening of 21st, temperature 100, pulse 120, respiration 32; 22d, temperature 101, pulse 120, respiration 35, wound looked well; evening of 22d, restless, temperature 102, pulse 124, respiration 40; 23d, temperature 101, pulse 124, respiration 36. Physical examination: Well developed and nourished. Expression anxious; tongue slightly coated; pupils equal and react. Slight glandular enlargement in left axilla and groin. Heart rapid, regular, no murmur. No friction rub. Heart's dulness extends to right, $\frac{1}{4}$ inch beyond the right border of sternum. Right

lung normal. Left lung shows good resonance in front, becoming dull in posterior axillary line, dullness over lower third of back. Just inside nipple on the left is a slight discoloration in fourth interspace; 1 inch inside and $\frac{1}{2}$ inch below nipple is a circular wound in the skin, with surrounding yellowish discoloration (Fig. 5.). Wound $\frac{1}{4}$ inch

FIG. 5.

in diameter. Swelling and tenderness over pectoral muscle. No râles in lung. Slight discoloration in back over lower end of scapula. March 24, some pain in left chest. Physical signs in left lung have increased and lower third of back is flat, with bronchial breathing and increased voice sounds. March 25, seen by Dr. T. M. Rotch, who made diagnosis of pericarditis with effusion. Cardiac dullness now extends to line midway between right nipple and sternum. Slight friction rub over third space on left. Lung on left is now flat for lower two-thirds of back. March 27, respirations have dropped to 30. No râles in lung. Physical signs in heart and lungs the same. Small amount of bloody sputum to-day. March 28, heart's dullness $\frac{1}{4}$ inch to right of sternum. Lung is not so flat. No râles. March 29, just below angle of left scapula, slough seen in back. Bullet not felt. April 1, pericardium evidently not so markedly stretched to right as before. April 5, temperature 99.5, respiration 30, child comfortable. Dullness in lower left back, several moist râles. Condition satisfactory. April 10, steady improvement since last note. April 13, child is improving in general condition and temperature is falling. Dullness in left back persists. April 17, this morning child developed swelling in region of right parotid. Some tenderness to touch and pain on opening mouth. Isolated on suspicion of epidemic parotitis. Dr. Rotch confirms diagnosis. April 22, temperature down. Throat

not sore. Isolation stopped. April 27, last night temperature was up. No cause found. Up on go-cart in sun parlor. May 2 boy has been up and dressed about the ward. May 6, x-ray shows bullet about in middle of lung. May 7, physical examination: Heart sounds regular, not rapid, no murmurs. Relative dullness from anterior axillary line, third rib down to seventh rib. Area of dullness below and to right of angle of left scapula. Breathing somewhat diminished in left back in region of scapular angle. In left axilla, from fourth to seventh rib, extending to post-axillary line, is a region where breathing is almost entirely absent. Boy discharged relieved.

It is a fair presumption, from the situation of the wound externally, the situation of the bullet in the lung and the existence of pericarditis, that the apex of the heart muscle was perforated. A few years ago such injuries were assumed to be fatal, but to-day we know that punctured wounds of the heart may recover under suitable treatment. The immediate dangers are profuse, overwhelming hemorrhage, or a hemorrhage which trickles into the pericardial sac and thus gradually compresses the heart and stops its action. Life has been saved by operating and removing the clots which interfered with the action of the heart and suturing the heart muscle.¹

ABSCESS OF ABDOMINAL PARIETES.

A. A., age six years, was admitted March 2, 1903. Family history and previous history negative. Present illness: Three weeks ago was struck with a club on right side of abdomen. Has not been in bed; has walked, but with considerable pain, leaning to one side. Bowels have moved with aid of castor oil. Sleep is prevented by pain; appetite very poor. Physical examination: Fairly developed and nourished. Looks sick. Lies by preference with thighs flexed. Pupils equal and react. No efflorescence. Throat normal. Linear scars behind each ear and on left side, in posterior triangle. Heart and lungs normal. Glands in each groin. Hyperextension of right hip limited. Limbs otherwise normal. Spine held somewhat stiff, especially to hyperextension. Child cannot be made to stoop or bend over while sitting. Walks rather guardedly, but not with the characteristic gait of Pott's disease. In right side of abdomen, just above McBurney's point, is a visible and somewhat indefinitely palpable tumor about two inches in diameter. No redness, increased surface temperature or fluctuation. Tumor moves with respiratory movement of abdominal wall and appears confined to abdominal parietes. Extremely sensitive. No general abdominal tenderness or distention. No evidence

¹ L. L. Hill: N. York Med. Rec., Dec. 15, 1900, p. 921. Case of C. H. Mastin: Transactions of American Surgical Association, vol. xiii, p. 273.

FIG. 2. Photomicrograph of section of papilloma.

FIG. 3. Photomicrograph of section of papilloma, showing that the growth is superficial, and does not involve the deeper layers.

FIG. 7. Radiograph of hands.

FIG. 8.

of ascites. Right rectus more tense than left. Whole abdomen, except upon pressure over tender area, is flaccid. Operation, Dr. Cushing: Ether. Incision one inch long over abscess and small amount of pus evacuated. Abscess evidently between external and internal oblique. Irrigated with peroxide. Iodoform wick. Dressing. Good recovery from ether. One week later the wound was nearly healed, the granulations were healthy, slight discharge. Motions of hip apparently normal. Flexibility of lumbar spine normal. Lumbar spine slightly flattened. Discharged relieved. Blood examination, March 3: 21,400 leucocytes. Bacteriological record: *Streptococcus pyogenes aureus*. Apparently this was a hematoma, following the injury, which later suppurated. The case was sent to the hospital with the diagnosis of appendicitis with abscess.

CONGENITAL DEFORMITY OF HANDS AND FEET.

P. F., a boy of eleven years, admitted Dec. 5, 1902. Family history negative. Only child. Previous history: Normal labor, full term, measles. Present illness: Right hand operated on at another hospital when one year old. No other treatment. Since that time, so far as known, the child has had no trouble of any sort except the deformity. Physical examination: Peculiarly developed, strong, well nourished, good color. Tongue large, flat and fissured. Intelligence good. Ears outstanding. Forehead low and slightly retreating. Head, normal shape, rather small. Chest flattened at the sides. Costal cartilages prominent. Heart and lungs normal. Abdomen soft and full; normal. Penis small. Testicles in scrotum. Thighs large. Spine normal. Right arm normal. Right thumb, terminal phalanx, turned out at angle of

30°. Right little finger abnormally wide at base, as though there were an extra metacarpal. At base, scar of old operation for supernumerary finger. Left arm normal. Left thumb similar to right. On left hand the supernumerary finger beyond little finger, which seems to consist

FIG. 8.

entirely of an extra metacarpal; between it and main body of hand is a web $\frac{1}{2}$ inch wide, which runs up on outer side of little finger (Figs. 6 and 7). Movements of fingers not interfered with. Each foot presents six webbed, rudimentary toes, with imperfectly formed nails. Right foot apparently has five distinct metatarsals with corresponding phalanges and on inner side of these, an imperfectly formed metatarsal, with which articulate distally and internally irregular phalanges, three or four in number, forming a broad, flattened great toe. The left foot is similar in conformation. (Figs. 8 and 9). The great toe of

FIG. 6.

FIG. 10.

each foot is turned in at an angle of 30° with the axis of the foot. Operation, Dr. Burrell: Ether. *Left hand:* Incision starting at end of ulna and coming up around extra finger, forming an oval. The dissection was carried down to the joint of supernumerary finger and the finger was removed. Skin sutured with interrupted catgut sutures. Sterile dressing. *Feet:* Great toes removed. Incision at joint of toe and metatarsal up and around toe. Skin dissected back. Muscles cut and toes excised at joint. Skin sutured with catgut. Sterile dressing. On the seventh day the wounds were dressed and found to be healed by first intention. On the fourteenth day the patient was discharged (Figs. 10 and 11).

FIG. 11.

Medical Progress.

RECENT PROGRESS IN OPHTHALMOLOGY.

BY MYLES STANDISH, M.D., AND W. D. HALL, M.D., BOSTON.

(Concluded from No. 13, page 347.)

FUNCTIONAL OR HYSTERICAL AMBLYOPIA.

Gunn⁶ divides this class of patients into idiopathic, usually occurring in women, sometimes in children, and traumatic, occurring probably to an equal degree in both sexes, but always in adults, according to his experience. He thinks that we do not understand the nature of the affections of the nervous system resulting in hysterical manifestations, although many forms admittedly

have an element of simulation. In many cases a single central lesion will not account for all the symptoms. When such lesions actually exist it is not a pure functional disorder, but an obscure early manifestation of real organic disease. Many hysterical persons subsequently become affected with an ailment such as insular sclerosis, consequently these symptoms should be considered carefully. It may be assumed that in most of these cases there is a temporary, partial or complete loss of the power of conduction between the lower or middle visual centers and the highest conscious centers. Gunn has noticed in cases of supposed blindness in very young children that the child pays no special attention to moving objects, although ophthalmoscopically nothing appears to be wrong. He thinks the prognosis to be favorable in such cases, and considers it an instance of delayed visual consciousness, in which there may be retarded development of higher association fibers. He thinks that there may be some similarity between this class of cases and those which later in life are called "hysterical amblyopia," inasmuch as in both the retinae are perfect, the optic nerve seems to be perfectly able to transmit impressions, the tracts seem normal, the sight centers presumably are fit for work and actually are receiving impressions, and yet there is no consciousness of these impressions, or a consciousness lessened to the degree occasioning no response. In hysteria the association fibers have been normally developed and efficient, but from some unknown cause their functions seem to have been suspended. This loss of conducting power may be re-established by shock, suggestion, surprise or even a voluntary exertion, on the part of the patient, of his own will power. In hysteria a peculiarly susceptible nervous organization is to be considered. As regards the treatment, change of scene if it can be arranged; but if not, friends must be careful to avoid drawing attention to, or showing regard for, the symptoms. Nourishing diet, assafetida and faradism may be of value.

THIOSINAMINE.

Dr. George F. Suker⁷ has carried on a series of experiments to determine the efficacy of this remedy as an absorbent of corneal opacities, cicatricial contraction of the lids, exudative chorioiditis, symblepharon, capsular opacities and cicatricial ectropion. It is an active alterative belonging to the same group as iodide of potash and mercury, and has been favorably mentioned as a solvent of cicatricial tissue in other locations. Experiments made to determine whether it would prevent the maturing of cataracts were without results. The best mode of administration is in 3-gr. capsules once or twice a day, and after it has been used for five or six weeks, it is well to intermit for a week or ten days. It occasionally produces vertigo when first taken, which is an indication for diminishing the dose. Therapeutically it is a marked tonic, favors the absorption of exudates, transudates

⁶ Ophthal. Rev., October, 1902.

⁷ Journ. Amer. Med. Assoc. Aug. 9, 1902.