The breathing now became deep and regular, and colour returned to the cheeks and lips. He seemed to sleep until 7 A.M., when he roused and asked for some toys. At 7.30 there was an extremely severe seizure, caused apparently by a sudden profuse secretion of mucus in the fauces. This collection, after a tremendous effort, he swallowed. He spoke of some water which he saw in a glass standing on the mantelpiece six or eight feet from him, but was not affected by looking at it. Another attack threatening at 7.35, I again gave one-tenth of a grain of morphia. He remained quiet, talking occasionally, and vomited some brown fluid. At 7.50 he inquired whether I had gone away, and then asked to be held so that he could see the gas. At 8 A.M., without loss of consciousness, there was a clonic convulsion of the left arm and of both hands. Both thumbs were flexed into the palms, but not firmly. Secretion in fauces more and more profuse, but no spitting. Breathing consisted of violent shallow inspirations, rapid expiration, and long pauses. Vomiting of brown fluid continued, which was now darker and contained altered blood. At 8.30 A.M. a grand attack was imminent. Morphia was again given, after which he dozed. No further violent spasms, but constant starting and continual vomiting. Pupils slightly contracted. At 10 A.M. corneæ muddy, and commencing ulceration of lower external quadrant on left side. Profuse muco-purulent discharge from conjunctivæ. Large quantities of grumous fluid vomited. No further secretion of mucus in the fauces occurred. He lay quietly in bed, disturbed from time to time by convulsion of the left side of the body and face until now, when in a violent and general convulsion he died.

I have detailed this case with what may seem tedious minuteness; but while it falls to the lot of but few practitioners to observe any considerable number of cases of hydrophobia, there are scarcely any opportunities for observing the disease in infancy. This fact may be my excuse. It is only in childhood that the course of the malady is uncomplicated by the mental condition arising out of knowledge of the inevitable termination close at hand; and although this condition is often, if not generally, betrayed by but few outward signs, it cannot fail to influence the psychical symptoms manifested.

Shanghai.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF HÆMOPHILIA.

BY HERBERT W. PAGE, M.A. CANTAB., F.R.C.S. ENG.,
LECTURER ON SURGERY, ST. MARY'S HOSPITAL.

THE publication by Dr. Skelton of his "Three Cases of Hæmophilia in the same Family" induces me to publish a clinical note of a case recently under my care in St. Mary's Hospital, where the patient was a member of a family in which ten members, males and females, of three successive generations were hæmophilic, and in which there was some departure from the ordinary rule of transmission.

A young man aged twenty-two, tall, anæmic, and thin, was admitted with an extensive swelling of the left thigh reaching from the knee to the groin, and from its physical characteristics evidently due to blood extravasated beneath the muscles. Seven weeks previously he had been laid up with so-called rheumatism in his left knee, which left it stiff; and five days before admission he had stumbled and fallen, and his leg was bent forcibly under him. It seemed to him as if "the flesh was being torn from the bone"; he suffered great pain, and the thigh rapidly swelled. We very early suspected he was a bleeder, and learned from him that he had had profuse hæmorrhage at one time from the extraction of a tooth, that he was subject to epistaxis, had bled furiously some two years ago from the opening of an abscess, and three years before had been laid up for many weeks with a swelling of the right knee, which, he said, was thought at first to be a sarcoma, but which in all probability was an intra-articular hæmorrhage. His high temperature, and the considerable tenderness with some redness at one point of the thigh, led us to fear that suppuration might

supervene; and as the necessity for making an incision was above all things to be avoided, the limb was placed at perfect rest on a Macintyre splint, after having been carefully bandaged from toes to groin. He was soon easier, the swelling at once began to diminish, and in seven weeks had entirely gone, some slight stiffness of the knee alone remaining when he was discharged at the end of two months.

His family history was very carefully worked out by my dresser and friend, Mr. Davis; and although it was impossible for him to see the various members of the family, he nevertheless elicited the following facts. Paternal grandparents healthy; maternal grandmother was in early life a bleeder, though in what precise way it was not known. She and her husband were not related to each other, nor were her own parents. She had eleven children, of whom the first, second, fourth, sixth, seventh, and eighth (all males) are dead—a startling mortality; but whether any were bleeders or not it was impossible to discover. As, however, the ninth, a living male, is a bleeder, it seems probable that some of his elder brothers may have been bleeders also. One of these brothers had died in infancy, and none are known to have died of hæmorrhage. The third, fifth, tenth, and eleventh (females) are alive, and all are bleeders. The third child, and eldest daughter, now aged sixty, is the patient's mother, and some of her children alone, of the third generation, are, so far, bleeders. Her family consists of four daughters and three sons. The three eldest are daughters, all bleeders, if a history of severe epistaxis and tooth-bleeding is sufficient to establish this fact. The fourth is the patient; then come a daughter and two sons, of whom none are bleeders. It is true that these three, the younger brother and sisters of the patient, may perchance not yet have had the opportunity of showing their diathesis, but it may be hoped that their immunity is an indication that the hæmophilia is tending to die out. What may have been the original cause of the hæmophilia in the first generation it is of course impossible to say; but this pedigree affords, at any rate, a good example of a morbid diathesis becoming rapidly developed to an extreme degree in the second generation—so much so that the children were affected certainly from the third down to the very youngest member (the eleventh) of the family; and possibly declining again, and it may be disappearing altogether, in the third generation, the three youngest grandchildren not being affected. Of the fourth generation, there are two, a male and a female, children of the patient's eldest sister, who are not bleeders. A noticeable feature, also, in this history is the fact that males and females seem to have been affected in an equal degree, contrary to the supposed usual rule that the males alone or most chiefly suffer, and that when the males inherit the disease it is through their mothers, who themselves remain unaffected.

SOME REMARKS ON A CASE OF SMALL-POX ARISING IN MID-OCEAN.

BY CHARLES J. POWER, M.A. CANTAB., M.D. DUB.,
LATE SENIOR HOUSE-SURGEON, METROPOLITAN FREE HOSPITAL.

THE following communication may be interesting in connexion with the leading article on "Small-pox in Australasia," which appeared in THE LANCET of Jan. 22nd of this year.

In April of last year, two days after leaving Aden, a case of variola arose amongst the passengers of the Orient Company's R.M.S. *Chimborazo*. The patient was an Italian (a steerage passenger, who embarked at Naples). Before the diagnosis of variola was certain I had the patient removed to one of the boats, which was partly covered in, and otherwise fitted up for his accommodation. One of the patient's compatriots volunteered to act as nurse, and he was also strictly confined to the boat. No one was allowed to visit the patient but myself, and the part of the deck beneath the boat was fenced off. The food of both patient and nurse was drawn up by the latter, and all scraps of food, dejecta, &c., were lowered down at night and at once thrown overboard. In addition to this, the ship was repeatedly fumigated with sulphur, steam, &c. The supply of lymph on board enabled me to vaccinate all the most urgent cases. As I was the only surgeon on board I had to move about amongst the passengers in the exercise of my duties; but I always kept a special suit of clothes, well sprinkled with carbolic solu-

tion, for visiting the patient, avoided touching him, and took a bath immediately after my visit. On our arrival at Adelaide those of the passengers who had reached their destination were placed in quarantine, and the boat in which the patient was (who was now becoming convalescent, the scabs commencing to fall off, showing well-marked "pits") lowered into the sea, and taken off to a separate island to undergo his quarantine. We subsequently arrived at Sydney, where we were placed in quarantine for three weeks, the time to date from our arrival at Adelaide. All the rest of our passengers were landed on the quarantine ground at Melbourne and Sydney. The patient, who had never been vaccinated, had confluent small-pox, highest temperature 104° , never had any delirium, and made a good recovery. At the commencement of disease there was an uncertain history of vomiting and pain in the back. The "secondary fever" was well marked. None of the passengers, officers, or crew subsequently contracted variola. I believe the main factor in preventing the spread of the disease was the early isolation of the patient. Variola, never easy to diagnose, is rendered much more difficult in hot climates. I have seen cases of "prickly heat" closely simulating variola, and a mosquito bite which has not been scratched has exactly the hard, shot-like feel of a papule in small-pox. Of course these cases will not have the constitutional disturbance which variola usually has, though I have seen constitutional symptoms in bad cases of prickly heat, and these symptoms in modified variola are sometimes, at any rate at the commencement of the disease, very slight. I can strongly recommend the treatment of patients suffering from variola in a boat which is partly covered in, thus forming a tent, provided the climate is suitable.

The Australians have a great dread of variola, yet vaccination is not compulsory amongst the inhabitants. The sanitary authorities made a great point about the vaccination of our passengers and crew. All who had not been successfully vaccinated within two years had to undergo the operation or remain in quarantine for an indefinite period. Some of the quarantine regulations are very strict, but in other matters the sanitary authorities are lax.

Ald-erney-street, S.W.

CASE OF THYROTOMY FOR EPITHELIOMA OF THE LARYNX.

By W. R. H. STEWART,

AURAL SURGEON TO THE GREAT NORTHERN CENTRAL HOSPITAL,
SURGEON TO THE LONDON THROAT HOSPITAL, ETC.

B. K.—, aged forty-five, a chimney sweep, came under my care in 1883. A slightly-made, delicate-looking man, married, no children; does not remember his parents; one brother and two sisters living, all healthy; no history of hereditary taint; had been a sweep from boyhood; had suffered from difficulty of breathing for the last six months, which was much worse lately; paroxysmal cough; had lost flesh. When first seen there were marked laryngeal stridor, harshness of voice, and severe attacks of dyspnoea. There were no enlarged glands in the neck. Laryngeal examination revealed a rather large, irregular growth, springing from below the right vocal cord, and projecting across the larynx. A portion removed by laryngeal forceps and placed under the microscope proved to be epithelioma. A preliminary tracheotomy was performed, and when the patient was sufficiently recovered, the growth being large, thyrotomy was determined on. Ether having been administered and a tampon tube inserted in place of the ordinary one, the superficial structures were divided in the middle line, the thyroid cartilage was then carefully cut through in its whole length with a knife (there was no ossification), and the *alæ* drawn aside with retractors. Through this small opening the tumour was seen to be growing from the under surface of the right vocal cord, and involving the soft tissues around. It was seized with forceps and as much as possible of the tumour, with the vocal cord and adjacent structures, cut away with small curved scissors; the remains of the growth were then scraped away with a sharp spoon, and solid nitrate of silver applied. The hæmorrhage was rather profuse at first, but soon stopped. The *alæ* were then carefully adapted and brought together with silver sutures, strapping applied to the external wound, a drainage tube inserted, and antiseptic dressings used. The tampon tube was left in for twenty-

four hours, after which an ordinary tube was used for a couple of days. The patient was entirely fed for the first week by nutrient enemata. There was no sign of perichondritis, and the patient made an uninterrupted recovery. At the end of the fourth week, when he left at his own desire for domestic reasons, the external wound had completely healed and the internal one was cicatrised over, but up to that time he had not regained his voice. The cartilages, however, having firmly united, would lead one soon to expect a complete return. There was no sign of a recurrence of the growth.

I unfortunately lost sight of the case, as I was obliged soon after to leave England on account of my own health, and subsequent inquiries have failed to find the man. Where possible, either the above operation, or removal of half the thyroid, as illustrated in a case recently published in a medical journal, is, I think, preferable to the extremely fatal one of excision of the larynx. I strongly advise preliminary tracheotomy with retention of the tube for some days after the operation, and also feeding per rectum.

Devonshire-street, W.

HERNIOTOMY IN A VERY YOUNG CHILD.

By J. LIONEL STRETTON.

H. G.—, born July 28th, 1886; male child. On Aug. 16th a swelling was noticed in the right scrotum; this was found to be a rupture, and was reduced by my assistant. On Aug. 25th the rupture again appeared, and all attempts at reduction failed. Fæcal vomiting set in, and the child was evidently losing ground. There had been no evacuation, and, so far as the mother could tell, no passage of flatus since the rupture appeared, thirty hours previously to my seeing him. Under such circumstances I determined upon immediate operation. Chloroform was administered and an attempt made to reduce the rupture; this failing, I made an incision over the swelling about three-quarters of an inch in length, and exposed the bowel, which was an exceedingly difficult proceeding in such a small space. The stricture, which appeared to be situated at the inner ring, was exceedingly tight, and required two or three nicks before the bowel, which was in a healthy-looking state, could be reduced. Silk sutures were employed and the wound dressed with oiled lint. Ordered half a minim of tincture of opium every four hours; the mother to continue suckling.—Aug. 27th: Passed flatus last night; no sickness since operation. Temperature normal. Takes breast well. Wound looks healthy.—28th: There were two natural evacuations during the night. Child seems quite well. A little local inflammation round the wound.—Sept. 1st: The bowels have acted regularly. Sutures removed. Slight gaping.—7th: The wound has quite healed; the child has gained much in flesh, and appears to be quite well. There is no bulging, even on violent crying.

Remarks.—This case is remarkable on account of the age of the child, the immediate subsidence of all symptoms after operation, and the rapid recovery. I have been unable to find any account of operation in so young a child; but when I take into consideration all the disadvantages I laboured under in this case I should not despair of success in even younger subjects. He was the child of poor parents, and was brought to my house for me to examine him, and knowing that he was in better air and surroundings I operated at once, and he was carried home in a blanket, and certainly did not receive any skilled nursing afterwards, nor was it possible to prevent the dressings being soaked in urine.

Kidderminster.

A RARE COMPLICATION IN A CASE OF PLACENTA PRÆVIA.

By GEORGE D. LOGAN, M.B. & C.M. ED.

IN July last I was called to see a woman who lived four miles away, and who was said to be in labour and bleeding to death. On arrival I found her blanched and in a cold sweat. Her pulse was quick, irregular, and feeble. I was informed by an attendant that the patient, who was a primipara, had fainted twice, and that the pains were neither so strong as they had been, nor so frequent. She