irritation and the commencement of the contraction determined in both cases.

For this purpose, a projecting metal point on the axis of the cylinder is made to open an electrical current in the primary coil of an induction apparatus, and the momentary current induced in the secondary coil is used to irritate the nerve.

The cylinder has therefore at every revolution the same position at the moment when the nerve is irritated, and the curves of muscular contraction will coincide when the time elapsing between the irritation and the commencement of the contraction is the same.

Now it is found that the two curves obtained by irritating the nerve at the above-mentioned points do not coincide, but are separated on the cylinder by a distance which is the measure of the time occupied by the nerve force in passing between the two points. To obtain the value in time of this distance it is only necessary to know the rate at which the cylinder revolves. This is determined by allowing a tuning-fork, vibrating at a known rate, to record its vibrations on the revolving cylinder. We have, then, as the data of the problem,

I. The length measured on the cylinder of one vibration of known duration, for example, .01".

II. The distance measured on the cylinder between the commencement of the two curves of muscular contraction caused by irritating the nerve at two different points.

III. The distance measured on the nerve between the two points thus irritated.

From these three data the rate of transmission of nerve force can be calculated.

Let A = length in mm. of a vibration of .01";

Let B = distance in mm. between the commencements of the two curves;

Let C = distance in cm. between the two points irritated.

Then $\frac{AC}{B}$ = distance in metres travelled by the nerve force in 1".

Dr. White asked if the rapidity of nerve force could be properly measured by means of a circle of persons holding hands.

Dr. Bowditch responded that it could. The time required to pass a hand pressure round the circle diminishes by practice, for a while, but a limit is soon reached, and then about the same time is occupied each time that the experiment is tried.

Selected Papers.

ON THE DIFFERENT FORMS OF PNEUMONIA.

By H. C. Wood, Jr., M.D., Clinical Lecturer at the Philadelphia Hospital.

LIKE many other affections, pneumonia occurs both in an acute and a chronic form, though, as we shall see directly, what is called chronic pneumonia is often rather the result of acute pneumonia than a distinct, true inflammatory disease. The amount of cellular tissue between the air-sacs is so small, the union of bloodvessel, nerve and cells is so close, and the cavities of the air-vessels are so ready to receive secretion, that it is *a priori* inconceivable that an acute inflammation should occur which should confine itself to the inter-vesicular tissue, and whose exudation should be poured out into the walls—not into the cavities—of the air-sacs; and, as clinical observation bears out this anatomical inference, we are justified in our first assertion, that in all acute pneumonias the exudation takes place into the air-sacs.

A thought as to the anatomical structure of the lungs will at once show what the nature of such exudation must be. The blood-vessels abounding everywhere pour forth the exudation material of inflammation—fibrinous serum—which soon coagulates and forms the major part of the matters filling up the air-sacs. At the same time with this fibrinous serum no doubt white blood-corpuscles escape, and the epithelial cells lining the air-sacs, responding to the general irritation, multiply, so that under the microscope the exuded material is seen to be made up both of fibrinous filaments and cellular elements.

In most inflammations, as is well known, there are two diverse possible exudations, whose nature is largely determined by, or at least associated with, a certain condition of the general system.

Thus, pleurisy in a debauchee produces a profuse serous exudation, while in the strong man it plasters the membrane with lymph. The same thing occurs in pneumonia. In the vigorous adult the exudation, even as first poured out, is almost a jelly, while in the old or prematurely broken-down it is a thin liquid, giving origin to the so-called prune-juice sputum.

There can be no doubt that in these different forms of ordinary fibrinous pneumonia the proportion of the cell-element varies

in the exudation; yet it is always a subservient, secondary part of the exuded material.

There is, however, a third form of exudation, in which the cellular element predominates, and whose character is due not so much to any recognizable condition of the system as to the anatomical construction of the lung.

Remembering that the lining membrane of the bronchial tube is continuous with that of the air-cells, and bearing in mind also the tendency which all inflammations have to travel along a tissue or membrane affected rather than to pass to the diverse tissues upon which such membrane rests, you will see at once that *a priori* we would expect to find a form of pneumonia whose origin should be a catarrhal or epithelial inflammation of the bronchial tubes, and whose anatomical characters should be catarrhal rather than fibrinous, or, in other words, whose exudation should be composed chiefly of imperfectly-developed epithelial cells or mucous corpuscles. It also seems to be an anatomical necessity that such disease should at first affect small isolated portions of the lungs, as the minute lobules, although by the coalescence of these the whole lobe of a lung may at last be consolidated. Every bronchial tube has subject to it as it were a certain territory of lobules, forming, it is evident, in regard to a bronchial tube of any size, a triangular mass, whose apex is the tube in question, its base the outer portion of the lungs.

Clinical observation has continually demonstrated that in children this form of acute pneumonia is exceedingly common. * * * * It is rare in adults; partly, no doubt, because the development of activity of the child is greater than that of the adult, and the epithelial tissues partaking of this excessive developmental power are more ready for that rapid production of cells which we call catarrhal inflammation, and partly because the formation of such pneumonic consolidation in the child is often due primarily not so much to a passage of the inflammation from the bronchial tubes to the air-vesicles as to a collapse of the latter from the child's want of power to expel the secretions of the bronchial tubes and to the anatomical changes consequent on such collapse. In the adult, collapse of the lobules of the lung from such cause is so rare, except in certain low fevers, that it may be practically said not to occur. * * *

Ordinary pneumonia—that which is known by the German as croupous, by the French as fibrinous—is normally a cyclical disease,

in which, at the end of a period varying from three to seven days, the exudation being completed, there is a sudden fall in the temperature and abatement of the general symptoms, the patient entering as it were upon convalescence, or at least a period of repair, with the road to health open to him only by the freeing of the vesicles from their contained exudation. As the little plugs that fill the air-sacs are mostly much larger than the openings into the bronchial tubes, it is necessary that the exudation matter should be softened if it is to be expelled or absorbed. Rindfleisch believes that but a comparatively trifling portion of the exudation is absorbed, and that nearly all of it is cast out by expectoration. But, from clinical observation, I think that absorption bears a very much more important part in the removal of the exudation than this. However this may be, there can be no doubt that both absorption and expectoration aid in the removal, and that to facilitate these the fibrinous plug undergoes two distinct changes—one a fatty degeneration, the second the conversion of the fibrinous filaments into a soft gelatinous mass, chemically allied to mucin; and that while this is occurring there is a free formation of cells upon, and, to some extent, even within, the alveolar walls. If matters progress favorably after the exudation is softened, it is gradually removed by expectoration and absorption, so that at last the lung is free and the patient well. Unfortunately, however, not always does the disease end so favorably; for, instead of resolution following exudation, the latter may remain unchanged, and the result be, first, chronic pneumonia; or, secondly, suppurative changes in the lungs may occur, and the so-called gray hepatization be the result; or, thirdly, caseous degeneration may ensue, and the so-called infiltrated tubercle result; or, fourthly, gangrene of the lung may take place.

The anatomical cause of the chronic pneumonia remaining appears to be simply the failure of the ordinary changes to occur. The fibrin does not undergo change into mucin, and the cells seem simply to shrivel away, but not to be in sufficient quantity to cause disintegration of the entire mass. Why these changes fail to occur I do not know. Possibly there is something in the original constitution of the exudation which interferes; possibly the ancients were not altogether wrong in their belief in the absorbents playing an active part in the physiological and pathological economy. * * *

Niemeyer makes no distinct allusion to it

in the article in his *Practice upon Pneumonia*. Jaccoud, however (*Traité de Pathologie interne*, tome ii. p. 49), describes it very distinctly. He states that there are two forms of it: one in which the arrest of the acute process seems to occur at the end of the stage of exudation; the other in which it takes place during the period of liquefaction and absorption. He says that in the first case the lung-tissue is compact, homogeneous, impermeable, vaguely granular. The appearance is very much that of the acute disease, the only difference being in the color and contained liquids; the hyperæmia has disappeared, and the color becomes slaty-gray, with black spots, due to the deposition of pigment granules; the absorption of the liquid is total, so that the tissue is dry and condensed. In the other case, the aspect of the lesion is different: it does not form a compact mass all in one block; the permeability of the lung is established in places, and the alveolar cavities contain some fibrin, dried and fatty purulent globules, and atrophied alveolar cells: the breaking down of the exudation has taken place, but the elimination has been arrested.

Of the second of these forms I know nothing from experience; the first I have seen repeatedly. How long this form of chronic pneumonia will endure I do not know, but I have seen many cases in which the history pointed to its having existed for a long time. It is probable that in many instances it finally causes chronic interstitial pneumonia, and, at last, contraction and a cirrhotic condition of the lung, or else ends in caseous degeneration. That it often does not do so for a very long time—that patients far advanced in life die, and portions of the lungs are found affected with chronic pneumonia, whose origin is obscured in the far past and whose existence was not expected during life—I most certainly know. * * *

The second termination of pneumonia in our schedule is in diffuse suppuration, if such term is allowable—that known as gray hepatization. * * * It is always to be feared in any case when the sudden fall of temperature—the crisis—fails to develop, or, developing itself, fails to persist, and, at the same time, very low typhoid symptoms are manifested; and especially is the danger imminent when along with these symptoms coarse moist râles are to be heard in the affected lung. Some authorities ascribe importance to distinct chills or repeated slight shiverings as diagnostic of the formation of pus in the lungs. * * *

Now, what are the anatomical differences

between the processes of resolution and gray hepatization? It must be borne in mind that in cases of resolution during the time when the exuded material is being removed there is a free formation of cells in the epithelium lining the air-sacs, and that there is also a tendency to over-rapid cell-increase within the alveolar walls. Now, in the degeneration of which we are speaking, this tendency being exaggerated, cell-multiplication takes place in both of these positions with great rapidity. Within and without the walls of the vesicles cells multiply. More than this, I am convinced, though not in a position absolutely to affirm it as proven, that in all these cases from the very onset the inflammatory products are largely cellular; that the exuded material is very poor in fibrin and rich in cells; that, in other words, the affection approaches closely the catarrhal type. The sputum in such cases certainly indicates it; and it is inconceivable that a mass composed chiefly of fibrin filament should be changed into a purulent, cellular liquid.

However this may be, excessive cell-multiplication—a multiplication so rapid that reproduction takes place at the expense of vitality—occurs on and within the walls of the alveoli of a lung about to undergo the purulent degradation. Under its influence the parenchyma of the lungs swells up and becomes more dense and heavy. At the same time the granular appearance which the filled alveoli produce in the red hepatized lung becomes less marked, from the swelling of the alveolar walls, and the color changes from reddish to whitish or yellowish. This alteration of the color is evidently due in some measure directly to the rapidly increasing cells within and without the alveoli, but chiefly to the pressure which they exert upon the bloodvessels, preventing the passage of blood into the affected tissues, which therefore become exceedingly anæmic.

The degeneration of these cells into pus or pus-like globules is of double origin. In the first place, their own native want of vitality facilitates their death; but I believe an even more important factor is the pressure which they themselves exert, cutting off their supply of blood—i. e. of nutriment—and causing death by starvation. I want to lay stress upon this point, because this purulent degeneration of the lung is the same in kind as the caseous degeneration—the so-called infiltrated tubercle—caused by the same anatomical changes, only that in the one case the course is rapid, in the other slow; in the one the cells mul-

tiply quickly, drive out the blood, and produce rapid local death, with exudation of more or less serum; in the other the cell-multiplication is slower, the blood-current is more gradually lessened, the death takes place more slowly, and is accompanied with desiccation. The two cases bear almost the same relation to each other that the moist gangrene which follows sudden arrest of circulation does to the dry gangrene caused by atheromatous arteries. * * *

It is in cases of cellular exudation especially that the last termination of acute pneumonia of which I shall to-day speak is especially apt to occur—that which was formerly known as acute infiltrated tubercle.

There is in such cases of pneumonia, from the very outset, an intense and rapid multiplication upon the alveolar walls of the epithelial cells, which, accumulating more and more, at last entirely fill up the alveoli affected. It is self-evident that such an exudation is not nearly so easily gotten rid of as a fibrinous one. The latter may soften and break down into a mucoid mass, but cells do not do this readily. More than this, the tendency to reproduction of cells does not confine itself to the epithelial tissue, but affects also the inter-alveolar tissue, until at last a solid mass is formed, the cells within crowding those without the alveoli. As a result, pressure upon the bloodvessels is exerted, until their calibre is obliterated or the current in them stagnated and the supply of blood to the part is cut off. What then happens? Death of the part affected. The deprivation of blood is, however, gradual, not sudden, and the death is slow. The cells in the interior of the alveoli are the first to suffer, because they are the farthest removed from the nutritive supplies; having from their very birth a tendency to fatty degeneration, they slowly undergo it, and at the same time shrivel up under the pressure, which squeezes out of the tissue the fluids contained in it. Even with the naked eye the change can be marked, commencing as a whitish spot in the middle of a dark mass of consolidated tissue, and spreading from within outwards. * * *

The degeneration of the cells of the affected part, continuing, spreads from the centre of an alveolar cavity to the alveolar walls, which are soon involved in the common death. Then from the portions of lung not yet affected serum begins to be thrown out, and to permeate the mass, until at last the whole diseased tissue breaks

down, melting away alveolar walls and all, and a cavity results.—*Philadelphia Medical Times*.

Medical and Surgical Journal.

BOSTON: THURSDAY, JANUARY 25, 1872.

THE ETIOLOGY OF TYPHOID FEVER.

DISEASE makes no discrimination of social position in its selection of victims, and, other things being equal, prince and peasant are in this respect identically human. The very serious and at one time alarming seizure of typhoid from which H. R. H. the Prince of Wales is at present convalescing exemplifies this equality; but while the mode of attack, the natural course of the disease and its complications have in no essential particular differed from what might have fallen to the lot of the meanest of the heir apparent's prospective subjects, the high position of the royal patient has stimulated a rigid investigation into the conditions which may be regarded as causative of the attack and has suggested renewed discussion of the subject in its general bearings. Medical commissioners were despatched to Londesborough Lodge, where the Prince was thought to have contracted the disease, and it is curious to observe the unanimity which characterizes the results of the inspection, as they are reported in recent English medical journals. A single quotation from the *Lancet* will serve as a sample of the opinions expressed.

"The most damning fact is that two healthy persons (the Prince and Lord Chesterfield) should have been attacked with typhoid fever, the only history common to them both being the circumstance of their having occupied the same bed-room in successive weeks, that bed-room being placed on the summit of a closed sewer with nothing but a few inches of water to protect the sleepers from the probable source of their disease."

Without at present criticizing the justice of such conclusions (which, by the way, reflect pretty clearly the prevailing English