staphylococcus pyogenes aureus, *spirillum cholerae*, *bacillus phosphorescens*, *bacillus pyocyaneus*, a *sarcina*, and a yeast.

The above organisms in broth culture were sealed in thin glass tubes and introduced directly into liquid hydrogen contained in a vacuum jacketed vessel immersed in liquid air. Under these conditions they were exposed to a temperature of about -252°C . (21° absolute) for ten hours. At the end of the experiment the tubes were opened and the contents were examined microscopically and by culture. The results were entirely negative as regards any alteration in appearance or in vigour of growth of the micro-organisms. It would appear, therefore, that an exposure of ten hours to a temperature of about -252°C . has no appreciable effect on the vitality of micro-organisms.

We hope to extend these observations upon the influence of the temperature of liquid hydrogen on vital phenomena and to make them the subject of a future communication, and to discuss their bearing upon problems of vitality.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF ACTINOMYCOSIS.

BY EDWARD T. JONES, M.R.C.S. ENG.,

AND

JOHN C. MACKAY, M.D. ST. AND.

THIS case appears to be worth recording because of (1) the uncommonness of the disease and (2) the enormous doses of iodide of potassium taken during treatment. The patient was a male, aged 42 years, who is the driver of a traction-engine used with a threshing-machine. Early in November, 1898, he had an old stump of a tooth removed from the right side of the lower jaw. This was followed by enormous swelling of the tongue accompanied by great pain. This apparently subsided under treatment and he came under our care a month later with a large swelling under the jaw in the anterior triangle, which was rapidly increasing. The skin was very red, thickened, and oedematous; the tongue was bound down and when protruded came out markedly on the right side. At one spot there was a patch of softening, this was incised and was found to contain thick curdy pus loaded with actinomycotic granules (confirmed by the Clinical Research Association). The treatment adopted was to open all places of softening as they appeared, to swab out the cavities with 1 in 3 carbolic solution, and to administer iodide of potassium internally. The dose of the iodide was increased from 30 grains per day to a maximum of four drachms per day in the course of six weeks. The patient never had any unpleasant symptoms from the iodide; in fact he put on flesh. The changes in the discharge and granules due to the iodide as recorded by Fairweather¹ were observed. The disease proved extremely chronic, spreading in the subcutaneous tissue downwards and forwards over the larynx, crossing the middle line. Finally it cleared up and early in March, 1899, the patient was discharged with no further sign of the disease and with nearly all the induration and thickness of the skin gone. Whether the iodide had much to do with the apparent cure it is difficult to say. Indications for treatment were not easy to find. Veterinary surgeons appear to treat their cases in cattle with enormous doses of iodide with success. Perhaps more radical surgical treatment at the beginning, as recommended by Ransome,² would have produced a good result more quickly, but on the other hand it is worth noting that with the exception of one week at the commencement of treatment our patient was able to keep at his work during the whole time.

St. Columb, Cornwall.

NOTE ON THROMBOSIS OF CEREBRAL VEINS OCCURRING IN A CASE OF CHLOROSIS.

BY EDWARD F. M. NEAVE, M.B., CH.B. EDIN.

THROMBOSIS of cerebral veins or sinuses is a very rare complication of chlorosis. Dr. Byrom Bramwell in his book on anæmia states that "thrombosis of the veins of the leg occasionally occurs [in chlorosis], but thrombosis of the cerebral sinuses very rarely." He does not record a single case of the latter. I have not been able to ascertain the frequency of this complication, but judging from the scantiness of the literature on the subject it must certainly be very rare. On consulting Osler's "Practice of Medicine" I find that Brayton Ball has recently reported a case and has collected 10 or 11 others from the literature of the subject. All were in girls with anæmia or chlorosis. Ball's patient was dull and stupid and had vomiting, dilatation of pupils, and double-choked discs. Slight paresis of the left side occurred. In some of the other cases the patients had headache, vomiting, and delirium. In Douglas Powell's case with similar symptoms there was loss of power on the left side.

The diagnosis of cerebral thrombosis is very difficult and in many of the cases it was made only at the necropsy. The symptoms pointing to thrombosis in the cerebral sinuses and veins are described by Sir Dyce Duckworth¹ as follows: "(1) headache, at times very severe, and sometimes plainly localised; (2) vomiting, generally severe; (3) motor symptoms, indicative generally of irritation, either cortical or central—hence tremors, jerking, spasms, and contractures in other parts, usually of one side, ending in paresis or paralysis; (4) drowsiness, delirium, coma; (5) variations in the size of the pupils."

The following case, in which most of the above symptoms were present, seems worthy of record:—

The patient was a married woman, aged 31 years, and she had no children. She complained of breathlessness, palpitation, and weakness. The duration of the illness was indefinite, but she had felt ill for some months and ultimately became so weak that she had to take to her bed. The family history was good. The patient had anæmia when a girl several times, but she always quickly recovered under a course of iron. When I visited her on May 19th, 1900, I found her lying in bed on her back, looking extremely exhausted and presenting most of the typical features of chlorosis. Her development and nutrition were good, but her face was extremely blanched and the lips and conjunctivæ were very pale. Pulsation was visible in the neck and a *bruit de diable* was well marked. On the 20th the patient had a few fainting fits, from which she always quickly recovered. On the next day (the 21st) she had a severe attack of vomiting, with severe headache. A blue pill, followed by a saline draught, was given with some benefit. On the evening of the 22nd she seemed very weak and could not turn on to her side. She slept fairly well but on awakening next morning she complained of a feeling of numbness in the left arm and leg, saying that those members appeared to be "dead." At the time she complained of great pain over the vertex and frontal region. This pain could not be localised to any one side. Her memory was confused—for instance, she did not know the day of the week or what she had had for breakfast. She was slightly delirious at times. The speech was unaffected. There was paralysis of the left arm and leg. As regards the leg the paralysis was complete, while the paralysis of the arm was partial. The opponens and adductor of the thumb were in action, the flexors of the wrist were slightly in action, while all the other muscles of the limb were paralysed. There was no facial paralysis. When the tongue was protruded it pointed distinctly to the right side. The left knee-jerk was slightly exaggerated. There was no ankle clonus. At this time there was incontinence of urine. The pupils were very large and they reacted to both light and accommodation. The discs, beyond being pale, seemed normal. There was no optic neuritis. The urine contained no albumin or sugar. The pulse was 120 and the temperature was 100°F . As regards the sensory symptoms there was slight anæsthesia of the left arm and leg. There was no hyperæsthesia.

¹ Brit. Med. Jour., June 27th, 1896.

² Ibid.

¹ THE LANCET, Feb. 22nd, 1890, p. 404.