vomited. Her mother died from phthisis. Nothing abnormal could be made out on auscultation of the lungs, but the heart's action was rapid and heaving, and there was a slight systolic murmur at the apex conducted outwards. In order to relieve such of the symptoms as were referable to the stomach I ordered her a mixture of papain, and the vomiting forthwith ceased. She is still under treatment.

CASE 13.—A man aged fifty-one years came on Sept. 15th, 1894, complaining principally of pain and vomiting after meals, from which he had suffered off and on for seven years. There was some consolidation at the apex of the left lung. The vomiting ceased at once under papain, though the pain did not entirely disappear under the treatment. I then combined small doses of morphine with the papain, under which the pain and other symptoms disappeared, and he was soon enabled to discontinue his attendance.

I have chosen these cases pretty well at random in order to show in what instances papain did good and also those in which it either did not afford relief or had for some other reason to be discontinued. In uncomplicated cases of atonic dyspepsia papain appeared to exert very beneficial effects, and in the instances in which this was the case I was struck by the almost immediate relief thus obtained. In only three cases in which I have prescribed the drug did it apparently fail, but in two of these (Cases 7 and 9) the patients were emotional women in whom the dyspepsia was associated with subjective symptoms of a hysterical character. The third case was that of a man aged forty-six years, an alcoholic, who suffered from dyspepsia with waterbrash, which yielded to bismuth.

Mecklenburgh-square, W.C.

Clinical Notes :

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF DIABETES INSIPIDUS.

BY ROBERT J. LEE, M.D. CAMB., F.R.C.P. LOND.,
LATE PHYSICIAN TO THE HOSPITAL FOR CHILDREN, GREAT ORMOND-STREET.

A YOUNG woman nineteen years of age was for many years the subject of diabetes insipidus. The malady was attributed to a fall from a hayrick when she was about four years old, and up to the time when she had the attack about to be described, which occurred last June, there was nothing unusual to record. The chief trouble during those years was thirst, particularly during the night, which led to the drinking of between three and four quarts of water after going to bed and getting up in the morning. The illness, which commenced in June, followed immediately after exposure to the strain of an examination, and retention during some hours, without permission to leave the room, caused great suffering each time it occurred. Great weakness followed immediately; the symptoms of diabetes ceased, and there was no thirst during the night. When the case came under the care of my friend, Mr. James Kingston Barton, the great state of prostration and feverishness led to the suspicion that this was some form of remittent fever, and soon it was considered proper to report the case to the medical officer of the parish. The skin was highly congested, the pulse quick, and the temperature, when Mr. Barton first took it, in the morning was 102.5° F. and in the evening 104.5°. It continued like this for a week and then fell for a fortnight two degrees morning and evening. Then it rose three degrees—that is, to 103° and 105°—for a week, and fell again lower than it was before. In this way from June 22nd to Aug. 22nd the state of fever continued without any symptoms of intestinal disorder or of other serious nature except attacks of nasal hæmorrhage so violent as to cause great alarm. During the next month a gradual fall of temperature to the normal took place, and gradually the night thirst returned, so that by the middle of September there was nothing to note except great prostration and not enough strength to allow of the patient being moved to the seaside till some weeks further had passed. By degrees during the winter the patient's strength has been restored to some extent, but not entirely to its former state, and the case seems to deserve notice in the interest of those who may be exposed in examinations at

schools, where such easily prevented accidents are possible. So far as treatment deserves notice, it may be remarked that no effect seemed to be satisfactory, and that not until the symptoms of returning diabetes displayed themselves did there appear to be any sign of improvement. It is strange to have a person in whom the healthy condition of the kidneys is one favouring excessive secretion of urine, and that any interference with this action is liable to serious constitutional disturbance. If this case recalls similar cases to others it would be interesting if they were recorded, and would help us to comprehend the curious and inexplicable phenomena that appear in diabetes in its various forms.

West Kensington, W.

SWALLOWING OF A PIN BY A CHILD.

BY JAMES TILY, M.D. ST. AND., L.R.C.P. EDIN.,
M.R.C.S. ENG.

INSTANCES of a pointed metallic body passing harmlessly through the alimentary tract are sufficiently unusual to justify the publication of the following case. A child aged five years having swallowed a pin on the evening of Nov. 21st, I directed that solid food, such as rice pudding &c., should be taken, and also gave a mixture containing tincture of opium, tincture of lavender, and chloroform water. On the evening of Nov. 23rd the pin was voided per anum, firmly packed in a solid cylinder of fæces, having consequently traversed the alimentary canal in forty-eight hours. It may be described as a small scarf-pin, having a total length of almost an inch and a half. The head is circular, flat, very thin, and nearly three-eighths of an inch in diameter; the stem is of brass, slender, and very sharp, and its upper part is bent so that the stem is parallel to the plane of the head. The diagram gives a good idea of the size and shape of the pin.

Chiswick.

ON ABSORBENT DRESSINGS, ASEPTIC AND CHEAP.

BY ARTHUR NEVE, F.R.C.S. EDIN.,
SURGEON, KASHMIR MISSION HOSPITAL, INDIA.

FIFTEEN years ago there was no consensus of opinion with regard to the best materials for surgical dressings. Listerism was introducing everywhere the use of gauze with mackintosh outside. Gamgee and others were working with absorbent material, whilst tow and ordinary cotton-wool were the staple dressings elsewhere, and carbolic oil the favourite application. As a rule the dry dressings were not aseptic, and the antiseptic dressings were not dry. The introduction of medicated absorbent cotton-wool and wood-wool reconciled these two methods; gauze dressings have also been used dry for many years, and many of the old crude applications have been banished, so that in the hospitals of Europe the same cardinal principles are everywhere recognised and to a great extent the same materials adopted. It may be assumed that surgeons are now satisfied with the ample choice of sterilised or antiseptic and absorbent gauzes or wools, but that managers of hospitals and war officials are less content with these changes, which have involved "an altogether astounding increase in expense." The question of cheapness cannot be altogether set on one side. It is not a question of antiseptic *versus* septic surgery, but of *cheap* *versus* *expensive* antiseptic appliances. The French War Department has adopted peat as the best cheap absorbent dressing. Peat-moss appears to have been used successfully in Russia for the last ten years. Suggested by Leisrink, it was soon used by Schede, Hagedorn, and others. I tried it for a time, but soon fell back upon sawdust pads, which have now been our staple dressing material for over ten years. Thomas of Liverpool and others have used them in England. These have now been introduced by Surgeon-Lieutenant-Colonel Lawrie into the hospitals and dispensaries of Hyderabad State. They are cheap. In Kashmir sawdust costs about sixpence a hundredweight. The muslin bags can be made at the rate of about 500 for £1. They can be made aseptic or antiseptic. We impregnate the pads the day before use with a 1 in 2000 solution of mercuric zinc cyanide. They can be easily