

This article was downloaded by: [University of Florida]

On: 03 October 2014, At: 00:58

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Annals and Magazine of Natural History: Series 6

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

LXXIII.—Report upon the
stomatopod crustaceans
obtained by P. W. Basset-
Smith, Esq., Surgeon R.
N., during the Crusie, in
the Australian and China
Seas, of H.M.S. 'Penguin,'
Commander W. U. Moore

R.I. Pocock ^a

^a British (Nat. Hist.) Museum

Published online: 06 Oct 2009.

To cite this article: R.I. Pocock (1893) LXXIII.—Report upon the stomatopod crustaceans obtained by P. W. Basset-Smith, Esq., Surgeon R. N., during the Crusie, in the Australian and China Seas, of H.M.S. 'Penguin,' Commander W. U. Moore , Annals and Magazine of Natural History: Series 6, 11:66, 473-479, DOI: [10.1080/00222939308677559](https://doi.org/10.1080/00222939308677559)

To link to this article: <http://dx.doi.org/10.1080/00222939308677559>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

A retrospect of the speculations which have been set up and the views which have been expressed will allow much to appear as requiring to be confirmed by further observation. Nevertheless we should not under-estimate the difficulty of proof in the treatment of a question in which sometimes even but slight indications of old peculiarities must be of importance. This difficulty, however, will not be permitted to suppress the attempt at an elucidation. The circumstance that arguments may really be pointed out everywhere will allow this consideration of the case to appear admissible, while the fact that every attempt at an elucidation ought to be made will show that it is justifiable.

LXXIII.—*Report upon the Stomatopod Crustaceans obtained by P. W. Basset-Smith, Esq., Surgeon R.N., during the Cruise, in the Australian and China Seas, of H.M.S. 'Penguin,' Commander W. U. Moore. By R. I. POCOCK, of the British (Nat. Hist.) Museum.*

[Plate XX. B.]

DURING the past two years the Trustees of the British Museum have received from the Lords of the Admiralty an immense and very valuable series of Crustacea obtained by Mr. P. W. Basset-Smith, of H.M.S. 'Penguin,' in the Australian and China Seas.

The Stomatopoda alone of this series form the subject of the present communication; but it is probable that when the rest of the material is examined, the remainder of the orders will be found to be equally well represented by new and interesting forms.

I may add that, during a recent visit to the British Museum, Dr. H. J. Hansen, of Copenhagen, made a thorough revision of our extensive collection of Squillidæ. Amongst other important innovations, Dr. Hansen has suggested more than one new generic name for certain species that have been hitherto referred to previously existing genera; but until he has himself published the descriptions of these genera, I have not considered it advisable to adopt them.

(1) *Squilla fasciata*, De Haan.

Two specimens. Chusan (10–14 fath.); Holothuria Bank (34–36 fath.).

(2) *Squilla affinis*, Berthold.

Chusan (10-14 fath.); Kowloon Bay (Hong Kong), Holothuria Bank (China Sea) (38-52 fath.).

The specimens from Kowloon Bay and Holothuria Bank closely resemble the specimen named *gracilis* by Miers in the distribution of the patches of black pigment upon the segments of the abdomen and on the telson, as also in having the telson somewhat narrower than appears to be normally the case in *Sq. affinis*, and its spines longer. They resemble the typical form, however, in having only four or five spines on the raptorial limbs, and not six or seven, as in the example of *gracilis*. The specimens in question thus appear to be intermediate in characters between *gracilis* and *affinis*, from which we may conclude that the former can scarcely be permitted to rank as a distinct variety.

(3) *Pseudosquilla oculata* (Brullé).

A single example from Macclesfield Bank (China Sea).

(4) *Pseudosquilla ciliata* (Fabr.).

Three young examples from the Arafura Sea.

(5) *Gonodactylus chiragra* (Fabr.).

Baudin Island, Troughton Island, Damma Island, Baleine Bank (15-20 fath.), Amboina Bay, Arafura Sea, Macclesfield Bank (32 fath.).

A large number of specimens.

I doubtfully refer to this species a semi-larval form, measuring 9 millim. long, from Baleine Bank, which differs from the adult in having only a single median crest upon the telson.

(6) *Gonodactylus graphurus*, Miers.

N.W. Australia (20 fath.); Baudin Island (8-15 fath.); Baleine Bank, N.W. Australia (15-20 fath.); Arafura Sea; Holothuria Bank, China Sea (15-24 fath.).

A great number of examples.

The Squillidæ in the British Museum included by Mr. Miers under this head contained examples of genuine *graphurus* and others which have subsequently been made into a distinct species, *G. glaber* (erroneously spelt *glabrous*), by Brooks. Still later Dr. de Man has united the two again; but since the dorsal grooves characteristic of *graphurus* are well marked, without signs of failing, in all the adult speci-

mens obtained by Mr. Bassett-Smith, it has not seemed to me advisable at present to adopt Dr. de Man's opinion.

In one semi-larval form (10 millim. in length) from Baleine Bank the groove is represented only upon the fifth abdominal tergite.

(7) *Gonodactylus Smithii*, sp. n. (Pl. XX. B. fig. 1.)
(? Var. of *G. chiragra*.)

Colour a deep green, obscurely mottled, the manus of the raptorial limb reddish purple, the distal extremity of the penultimate segment indigo-blue; a reddish-purple spot on the inner side of the distal extremity of the large segment of the appendage.

This species is closely allied to *G. chiragra*, from which it appears to differ in being more robust and in the ornamentation of the telson and of the sixth abdominal tergite. The crests upon these two plates are the same in number as in *chiragra*, but are much more compressed and carinate. Thus on the sixth abdominal tergite the crests are ridges, flat above and produced without constriction into long spines, which considerably overlap the hinder border of the tergite; in *chiragra* these crests are wider, convex above and rounded behind, the spines being very short and sharply defined from the rest of the crest. In the telson, again, the median prominence is a strongly compressed ridge, the upper edge of which is almost straight and the posterior angle is regularly produced into a long strong spine; all the other crests are similarly compressed and carinate; in *chiragra* the median crest is convex above, sometimes tipped with a small spine, rounded behind and scarcely compressed.

The following measurements (in millimetres) will show the stouter build of this new form:—Total length (from base of rostrum to notch of telson) 23·5; width of fifth and first abdominal somites and of carapace 4·5, of second free thoracic somite 4·3; length of carapace 6·3, of eye 2·8.

An example of *chiragra* measuring 23 millim. in length has the fifth abdominal somite and the carapace only 3·8 millim. wide, the latter plate being 5·6 millim. long and the eye-stalk 2·5. Again, an example of *chiragra* with the abdomen 4·5 millim. wide measures 27 millim. long, the carapace being 7 millim. long and the eye-stalk 2·8. A second example of *Smithii* measuring only 17 millim. long shows corresponding differences when compared with an example of *chiragra* of the same length. These comparisons show that, in addition to being stouter in body, the eyes are longer in *Smithii*.

Two examples (♂ ♀) from the Arafura Sea, both possibly young.

A number of young examples of *chiragra*, both surpassing and falling short of these in size, were taken at the same time and in the same locality, and there is no difficulty whatever in distinguishing those that are here named *Smithii*. Dr. Hansen, however, when naming the Museum collection of Stomatopoda referred an exactly similar form to *G. chiragra*, so that it is possible that the view here expressed may be erroneous.

(8) *Gonodactylus excavatus*, Miers.

A single male example dredged at a depth of 26 fath. on Macclesfield Bank.

The rediscovery of this species is interesting, the locality of the type being unknown. The figure of the telson of this species given on pl. iii. of Mr. Miers's paper is very inaccurate. The excavation is represented as much wider than it really is, and the posterior angles of the telson appear to be single, whereas in reality they are double, consisting of two subequal processes.

(9) *Gonodactylus trispinosus*, Dana.


Two small examples from Baleine Bank (N.W. Australia).

Under this species, in his revision of the Squillidæ, Mr. Miers mentions two males that were obtained at Sharks Bay (W. Australia). According to Dr. Hansen, these examples belong to a different species, namely *G. stoliura* of F. Müller.


(10) *Gonodactylus tuberosus*, sp. n. (Pl. XX. B. fig. 2.)
(? Var. of *trispinosus*.)

Very closely allied to the preceding, *G. trispinosus*, the only differences between the two that I have noticed being found in the form of the rostrum and of the telson. In *trispinosus* the lateral spines of the rostrum are longer, thinner, and are directed forwards and outwards, so that if prolonged in the same direction they would pass along the outer border of the eyes. In *tuberosus*, on the other hand, the spines are much shorter and stouter, being directed forwards, so that the prolongation of their axes would pass through the external half of the eyes. Again the telson of *tuberosus* is more squared and its posterior angles less rounded; the prominences, too, are considerably larger, the median being almost spherical and so high that its summit is on a level with the summits of the median tubercles of the sixth abdominal tergite, although

A.



B.



Downloaded by [University of Florida] at 00:58 03 October 2014

these tubercles are themselves considerably more elevated than in *trispinosus*; moreover, the lateral tubercles of the telson are also much less elongate and more spherical; the notches separating the lobes of the telson are wider than in *trispinosus*, the external notch being particularly well-marked and widening towards the margin of the plate.

Measurements in millimetres of largest specimen:—Length from apex of eyes to end of telson 40; length of eye 3·3, width of cornea 2; length of telson 7·5, width 8·2.

Two examples from Macclesfield Bank, at a depth of 37 fath.

(11) *Gonodactylus Hansenii*, sp. n. (Pl. XX. B. fig. 3.)

Belonging to the same group as *G. Edwardsii*, *scyllarus*, *cultrifer*, &c.

Colour (in alcohol) a uniform yellowish pink, with a black patch of colour on the second segment of the external ramus of the swimmeret.

Eyes very large and globular, at least as large as in *G. Edwardsii*. The dorsal plate of the ophthalmic somite deeply bifid in front. *Antennal scales* about the same relative size as in *G. scyllarus*, being considerably shorter than the carapace.

Rostrum with evenly convex anterior border and evenly rounded angles.

Manus of the raptorial limbs moderately dilated at the base and armed with nine distinct, sharp, subequal teeth. The margins of the three posterior thoracic tergites narrowed and not dilated even as in *G. scyllarus*.

The margins of the abdominal somites thickened as in *G. scyllarus*, the posterior angles spined as in that species. The sixth abdominal tergite and telson closely resembling these plates in *G. scyllarus*, the median being elevated, carinate, and posteriorly spined as in that species; the crests on each side of the median one not so close to it as in *scyllarus*, and not divided by a groove; the median crest not continuous with the crest upon the internal spiniferous process, while the external crest, which in *scyllarus* lies close to the external edge of the anterior half of the telson, is situated more internally, being in the same line as the external edge of the median process on each side.

The movable spines on the exopodite of the swimmeret are longer than in *scyllarus*, the apex of the distal one being on a level with the margin of the distal segment of this appendage; moreover the crests upon the endopodite of the swimmerets are unequal in size, the exterior being considerably

longer and stronger than the interior; in *scyllarus* they are subequal.

Measurements in millimetres:—Length (from anterior margin of rostrum to the median notch of telson) 60; length of carapace at the side 14, of antennal scale 11; width of rostrum 4, length 2; length of eye and width of cornea 4; width of telson 10, length from anterior end of median keel to median notch 8.

A single female example from Macclesfield Bank, 35 fath.

On the whole this species seems to come nearest to *G. scyllarus*. It differs, however, in the arrangement of the crests of the telson, the size of the eyes, the form of the rostrum, the bifid ophthalmic tergite, and the number of teeth on the raptorial limbs. In having large eyes and many teeth on the raptorial limbs it resembles *G. Edwardsii*, Berth., but differs in the structure of the telson, the small antennal scales, form of rostrum, &c.

(12) *Gonodactylus carinifer*, sp. n. (Pl. XX. B. fig. 4.)

Belonging to the same section as the preceding, *G. Hansenii*, but closely related to *G. cultrifer*.

Colour (in alcohol) a uniform ochraceous tint, the dorsal surface marked everywhere with fine, close-set, irregular star-shaped patches of pigment.

Eyes very large and globular. The dorsal plate of the ophthalmic somite with its antero-lateral margins directed upwards.

Rostrum ovately convex, its extremity bent downwards. Antennal scale rather large, a little shorter than the carapace measured along the dorsal middle line. The manus of the raptorial limb moderately dilated and armed with three distinct sharp subequal teeth. The margins of the free cephalothoracic tergites not dilated; the posterior angles of the posterior abdominal tergites spinous. The sixth abdominal tergite furnished with six posteriorly spined crests.

The telson almost exactly like that of *G. cultrifer* (White), except that the median crest is much lower, being only about half as high as it is long, with its upper edge only lightly convex, and that there is no trace of a crest near the external margin of the anterior half of the telson.

Measurements in millimetres:—Length from anterior edge of rostrum to median notch of telson 27, of carapace at the side 7, of antennal scale 5·6; length of eye 3, width of cornea 2·8; width of telson 5, length from median notch to anterior edge of median keel 3·8.

A single male example from Holothuria Bank (China Sea), 24 fath.

Judging by its size the example here described is young, and it is necessary to bear in mind that it may prove to be the young of *G. cultrifer*. Since, however, it appears to have all the characters of an adult, I have decided, at all events provisionally, to regard it as a distinct form. It may be at once recognized from *G. cultrifer* by the presence of *three*, and not two, teeth on the hand of the raptorial limb, and by the lowness of the median crest of the telson.

EXPLANATION OF PLATE XX. B.

- Fig. 1. Gonodactylus Smithii*, sp. n.; last abdominal tergite and telson from above, $\times 2$. 1 a. Ditto, ditto, from the side, $\times 2$.
Fig. 2. Gonodactylus tuberosus, sp. n.; last abdominal tergite and telson from above, $\times \frac{3}{2}$. 2 a. Ditto, ditto, from the side, $\times \frac{3}{2}$. 2 b. Ditto, rostrum.
Fig. 3. Gonodactylus Hansenii, sp. n.; anterior end of body. 3 a. Ditto, last abdominal tergite and telson, from above, $\times 2$. 3 b. Ditto, ditto, from the side.
Fig. 4. Gonodactylus carinifer, sp. n.; anterior end of body. 4 a. Ditto, last abdominal tergite and telson, from above. 4 b. Ditto, ditto, from the side; nat. size.

BIBLIOGRAPHICAL NOTICE.

Éléments de Paléontologie. Par FÉLIX BERNARD &c. Première Partie (pages 1 à 528), avec 266 figures dans le Texte. Svo. Baillière, Paris, 1893.

PALÆONTOLOGY is here treated in a very full and masterly manner. The object and history of the Science are first noticed. Its relationship to Biology and Geology is clearly defined, and the necessity of a knowledge of these, as collateral studies, is insisted on. Its history extends from the ancient recognition of fossil shells having once been marine, to the systems of modern philosophy, comprehending far-advanced theories of natural history in its many branches, and the now-established doctrine of Evolution. Chapter II. deals with the definition of "Species," gradations of varieties, and transitional forms; also how far natural selection and the influence of locality and requirements have had effect. Correlation of structure and rudimentary organs are considered; and parallelism and convergence among forms, also aberrant and synthetic types, are carefully exemplified. How far there may be a vital force in species and in larger groups, and whether the perfection of races and individuals may be looked for, are among the problems still waiting for elucidation by the study of facts. In Chapters III. and IV. the Author defines natural classification and phylogeny, referring to the help of comparative anatomy in the study of the evolution of organs,