XL. On the existence of carbon in the sun

Professor John Trowbridge & C.C. Hutchins

To cite this article: Professor John Trowbridge & C.C. Hutchins (1887) XL. On the existence of carbon in the sun , Philosophical Magazine Series 5, 24:149, 310-313, DOI: [10.1080/14786448708628105](https://doi.org/10.1080/14786448708628105)

To link to this article: <http://dx.doi.org/10.1080/14786448708628105>

Published online: 29 Apr 2009.

Submit your article to this journal [↗](#)

Article views: 2

View related articles [↗](#)

In regard to the other three spectra of oxygen of Schuster we have nothing to say ; but so far as concerns the spark spectrum in air and the solar spectrum from wave-lengths 3749.8 to 5033.85 we can safely affirm that there is no physical connexion between them.

XL. *On the Existence of Carbon in the Sun.*

By Professor JOHN TROWBRIDGE and C. C. HUTCHINS*.

FROM the presence of absorption-bands in the solar spectrum at high altitudes, Captain Abney has been led to believe in the existence of certain hydrocarbons between the earth and the sun ; and Siemens's theory of the conservation of solar energy depends upon the supposed existence of carbon vapour in interplanetary space. It is not our purpose to discuss Abney's observations, or the truth of Siemens's hypothesis. We wish to call attention to the remarkable character of the carbon spectrum, formed by the Voltaic arc in air between carbon terminals ; and to draw attention to the evidence presented by the juxtaposed solar spectrum of the existence of carbon in the sun.

In our early experiments the carbon terminals between which the Voltaic arc was formed were heated several hours, while a stream of chlorine gas was passed over them. This operation was not entirely successful in removing metallic impurities. Subsequently we discovered that the spectra of these impurities could be readily distinguished from the marked fluted carbon spectrum, and we therefore employed the ordinary compressed carbon sticks employed in electric lighting.

For our work the nicest adjustment of slit was necessary, in order that no displacement of spectrum lines could possibly occur when the carbon spectrum was photographed in juxtaposition with the solar spectrum. This was accomplished by the use of a slit, the jaws of which opened equally.

One of Rowland's concave gratings, of 21 feet 6 inches in curvature and 14,000 lines to the inch, was employed. In order to avoid any possible displacement of the photographic camera during the operation of photographing the carbon spectrum immediately below the solar spectrum, a drop-shutter was arranged directly in front of the sensitive plate, the movement of which was independent of any movement

* From an advance proof from the Proceedings of the American Academy of Arts and Sciences, vol. xxiii. Communicated by Professor J. Trowbridge, of Harvard University, Cambridge, Mass., U.S.

of the camera. Preliminary experiments showed us the importance in this work of employing a spectroscope of great dispersion and of fine definition, giving also a normal spectrum. The use of a prism-spectroscope would undoubtedly have masked the phenomenon we have observed. For our purposes, therefore, Rowland's apparatus was peculiarly advantageous.

Our experiments led us to conclude that there is positive evidence in the solar spectrum of the existence of carbon in the sun. Before giving an account of our experiments in detail, a few observations may not be considered out of place.

One who studies the solar spectrum by itself, and who has had no experience in the formation and observation of metallic spectra, is apt to regard the dark lines in the solar spectrum as fixed in character and condition. A line which is seen by one observer, and not by another, is generally regarded as a terrestrial line formed by absorption in the earth's atmosphere. Certain lines are well known to be due to the terrestrial absorption, as can be easily proved by their appearance when the sun is observed at sunset, when the rays of light have to penetrate a greater thickness of the earth's atmosphere than at midday. The shifting layers of vapour in the sun's atmosphere also may, in certain cases, obliterate or strengthen certain lines of a metal. To understand this it is only necessary to extend the reasoning of the conservation of energy to the subject. It is a common lecture experiment to reverse the metallic lines by passing the rays of light produced by the vapour of the element through a layer of vapour colder than that of the source of the rays. The energy of the rays is thus absorbed in heating the colder layer. When the temperature of the vapour is increased, and becomes equal to that of the source, no reversal takes place. Thus, on the sun's surface the conditions for a reversal may be wanting at certain times, and faint lines may become bright. Their brightness may not be sufficient to affect the general illumination of the solar spectrum of which they form a part. Conditions may arise, moreover, in which the temperature of the reversing vapour may be called critical—at such a temperature that the faint reversal is sufficient to extinguish the bright line of a metal without producing a well-defined dark line. At certain epochs, also, the temperature of the vapour of any element in the sun may be higher than at other times; and certain lines may thus appear which are wanting when the temperature falls. One is forced to these conclusions in observing the conditions under which the varying character of metallic spectra are produced. For instance,

we have caused the rays from iron vapour to traverse a long and dense layer of iron vapour, and have observed that the strength of the lines and the number of reversals have been largely increased. In another experiment, the lower carbon of the electric lamp we employed occupied the centre of an electro-magnet. This was accomplished by passing the carbon through a hollow iron cone, and surrounding the latter by layers of wire, through which the electric current employed in generating the light passed. In this case the electric arc was spread out at right angles to the pole of the magnet, into a fan-like, intensely hot flame, which roared loudly, and which rarefied, so to speak, the iron vapour between the carbon terminals. The strength of the lines and the number of reversals were diminished under this new condition.

Another phenomenon may happen. When an excess of the vapour of one metal floats over or is mixed with that of another, the lines of one metal are superimposed upon those of another in the solar spectrum, and the stronger spectrum of one element may easily obliterate the weaker spectrum of another. Thus we have succeeded in completely obliterating the fluted spectrum of carbon in the green and blue, by photographing upon it the spectrum of iron, of nickel, and of cerium. A species of composite photograph was thus obtained. It is possible that in the future Galton's ingenious method of composite photography may be applied to the solar spectrum; and by a judicious selection of photographs of the elements, a composite photograph may be obtained which will closely resemble portions of the solar spectrum, and will enable us to judge of the composition of the reversing layers of the sun.

To the varying conditions which we have thus outlined are due, we believe, the disappearance in the sun's spectrum of the marked fluted spectrum of carbon in the green and blue portions.

A careful examination of the fluted spectrum of carbon, however, with the juxtaposed solar spectrum, discloses a remarkable fact: while traces of obliteration of the evidence of carbon vapour are seen, yet the general character of the lines in the solar spectrum immediately juxtaposed with the fluted spectrum of carbon near H lead us to believe that there is unmistakable evidence of the existence of carbon vapour in the sun. When the arrangement of the fine lines of the spectrum of carbon is plotted as a curve, and that of the dark lines in the solar spectrum immediately above the carbon spectrum is also plotted, the two curves have a

remarkable similarity in character, running with a slight convexity toward one axis.

In the first fluting at wave-length 3883·7 within the limit of ten wave-lengths, over 28 of the spaces between the fine bright lines of the flutings coincide with dark lines immediately in juxtaposition in the solar spectrum. When we consider that the progressive arrangement of these lines is exactly the same both in the spectrum of carbon and that of the sun, we cannot believe that this coincidence is the result of chance. On examining the spectrum of carbon in the region near H still further, a remarkable number of coincidences of the spaces between the bright lines of the carbon spectrum with dark lines in the solar spectrum will be observed. We are led, therefore, to conclude that the fluted spectrum of carbon is an example of the reversal of the lines of a vapour in its own vapour. Fluted spectra occur at comparatively low temperatures. When carbon is ignited, we have at first a continuous spectrum. When the temperature increases and the carbon is volatilized, fluted spectra occur, which consist of interruptions of the continuous spectrum by fine line reversals occurring in harmonic order. The same phenomenon can be observed in the spectrum of iron lines: through the centre of an iron line, when a sufficient amount of iron vapour surrounds the Voltaic arc in which iron is volatilized, reversal lines are always seen. Now if the iron lines were arranged in regular order, the reversals would also be in like regular order, and would coincide with similar reversals in the solar spectrum. Assuming the conditions at the sun's surface to be the same as those we have in the Voltaic arc when carbon is volatilized, the character of the carbon spectrum should exactly agree with the character of the solar spectrum juxtaposed. This is found to be true to a remarkable degree in comparing portions of the solar spectrum with portions of the fluted spectrum of carbon beginning at wave-length 3883·7.

Our hypothesis leads us to conclude, that, at the point of the sun's atmosphere where carbon is volatilized, so as to produce the peculiar arrangement of reversals observed, the temperature of the sun approximates to that of the Voltaic arc.