tubercles or bacilli they might very well represent the result of some tuberculous process. In view of the small number and size of erythroblastic foci in the spleen pulp one cannot conclude that the spleen was exercising any important function as a red blood cell forming organ. Nor was there much evidence of active red cell destruction except as already noted in the vicinity of the caseous areas.

Marrow.—The fat cells of the marrow have not entirely disappeared. The most notable feature is the dilatation of the thin-walled blood-vessels with red blood corpuscles. These congested blood-vessels traverse the whole section but there are no hæmorrhages into the substance of the marrow proper. The endothelium of the sinuses remains quite intact. The following cell types were noted. 1. Large mononuclear cells with somewhat vesicular nuclei and basophile protoplasm were most frequent. In morphology these cells correspond to the basophile myelocytes or undifferentiated cells of Dominici. Neutrophile myelocytes and polynuclear cells were present in fairly large numbers, though the granulation was difficult to demonstrate. Transition forms can readily be traced between these basophile cells which may sometimes attain a large size and the giant cells presently to be described. 2. Eosinophile cells, both mononuclear and polynuclear, occur in large numbers and in Giemsa preparations make a remarkably beautiful picture. 3. The giant cells present are of two types. Sometimes as many as ten may be seen in one field of the oil immersion Leitz $\frac{1}{2}$ th, oc. 4. Cells with large-lobed nuclei assuming the most bizarre shapes are most numerous. The protoplasm of these cells is slightly basophile. Many of them appear to be developed directly from the basophile undifferentiated cells already noted and in which mitotic figures are frequent. The second type of giant cell possesses a straggling irregular nucleus which is intensely pyknotic and a cytoplasm which is inclined to be acidophile. These cells frequently contain entire leucocytes in their interior. They are evidently degenerated megakaryocytes. 5. Lymphoid cells are not very numerous. 6. Mast cells occur in small numbers. 7. With the Unna-Pappenheim stain plasma cells are found to be as scarce in the marrow as in the spleen pulp. 8. Here and there one sees erythroblastic foci, generally of small size but sometimes fairly large clumps of developing nucleated red cells are met with. The nuclei stain intensely while the cytoplasm takes on a yellowish stain. Frequently the nucleus is broken up or presents a budding appearance. It is possible that this irregular division is the explanation of the great variations in size and shape of the red cells in blood films *intra vitam*. One could not conclude from a single section that red blood cell formation was actively proceeding, but when one reflects that large tracts of marrow were in a similar condition one must infer that the polycythæmia is to be explained by a general increase of the erythroblastic functions of the marrow. The large number of granular cells in the marrow certainly indicates an increase in the leucoblastic function of that tissue and would explain the polynuclear leucocytosis in the circulating blood that is so marked a feature of most of the recorded cases.

Liver.—The interacinar capillaries are greatly congested with red blood cells, though polynuclear cells are also frequent. The liver lobules and acini are quite intact and the cells apart from very slight fatty infiltration appear healthy. Clumps of bile pigment occur in some of the liver cells and here and there one sees the lumina of the smaller bile channels filled with inspissated bile thrombi. There are no focal necrotic areas and no micro-organisms."

Remarks.—There can be no doubt that clinically this case belongs to the group characterised by chronic cyanosis with polycythæmia and splenomegaly, of which examples have been recorded by Vaquez, Saundby and Russell, Rosengart, Osler, Parkes Weber, and others, and for which Parkes Weber² has suggested the name of "myelogenic polycythæmia." The increased number of red cells (from $7\frac{1}{2}$ to 8 millions per cubic millimetre) and the splenic enlargement together pointed to such a diagnosis. In addition, there was found by Parkes Weber an increase in the viscosity of the blood such as he has described in a previous case of his own, whilst a polynuclear leucocytosis which was present in our case has also been met with in some previously recorded examples of the disease. The cerebral and mental symptoms, too, which were such a marked feature in our patient have also been remarked in such cases before.

We are fortunate in being able to add another to the few

recorded necropsies on cases of this group, and as far as an explanation of some of the most striking clinical symptoms is concerned the result was satisfactory enough. The loss of vision, for instance, and along with it probably the attacks of giddiness and the final stupor, may be held to be explained by the extensive areas of thrombotic softening in the occipital lobes of the brain. It cannot be said, however, that much light is thrown upon the pathogeny of the disease by a study of its morbid anatomy. That the amount of blood-forming tissue in the bone marrow was increased there can be no doubt, which, with the absence of any signs of increased blood destruction in the liver, would point to the polycythæmia being the result of an increased formation of red cells. The enhanced viscosity of the blood which is the consequence of the polycythæmia and the abnormal strain which it throws upon the arterioles may perhaps explain the high degree of vascular fibrosis met with. The fibrotic degeneration of the heart was certainly a most striking feature of the morbid anatomy but was perhaps merely secondary to the thickening of the walls of the coronaries.

As regards the softening in the occipital lobes of the brain and the large pulpy areas in the spleen, there can be no doubt, we think, that they were due to thrombotic necrosis, the consequence of stagnation of blood which the high viscosity of the latter tended to induce. We found no evidence of any primary tuberculosis of the spleen such as has been alleged to occur in some of these cases and are inclined to think that in such descriptions areas of infarction may have been mistaken for tuberculous foci.

In conclusion, we would like to take the opportunity of expressing our indebtedness to those who so kindly helped us in the investigation of the case; to Dr. Parkes Weber especially for much advice and suggestion and for estimating the viscosity of the blood; to Dr. Boycott for his careful reports on the histological features of the blood; and to Dr. Ledingham for his minute investigation of the histology of some of the more important organs.

London Hospital.

SPIROCHÆTA PALLIDA (SPIRONEMA PALLIDUM) IN SYPHILIS.*

By THEODORE SHENNAN, M.D. EDIN., F.R.C.S. EDIN.,
SENIOR PATHOLOGIST, ROYAL INFIRMARY, EDINBURGH; LECTURER
ON PATHOLOGY AND BACTERIOLOGY, SCHOOL OF MEDICINE
OF THE ROYAL COLLEGES, EDINBURGH, ETC.

(Concluded from p. 667.)

IN Siegel's papers,^{2 3 4} in addition to the methods of staining and the description of the characters of his cytorhynchus luis, some details of previous discoveries in syphilis are given⁴—e.g., Klebs, Döhle, Stassano, &c.

Fr. Schultze,¹²⁹ Freund,¹³⁰ and Merk¹³¹ contribute papers on the cytorhynchus.

Spirochæta in yaws.—Castellani in Ceylon^{113 114 115} and Wellmann in Central Africa¹¹⁶ about the same time found spirochætæ in films from the cutaneous papules of yaws some of which in form correspond closely to those of syphilis. I have seen one of Castellani's preparations through the kindness of Dr. J. O. Wakelin Barratt. There is no question as to the great similarity to spirochæta pallida. Castellani will not draw any conclusions as to any possible relationship between yaws and syphilis, though in this connexion it is of interest that Mr. Jonathan Hutchinson¹³¹ holds in the face of great opposition that they are closely related, probably identical diseases.

Staining methods.—As I have indicated, Giemsa's method, or various modifications of it, is generally considered the best, but Leishman's modification of Romanowsky's stain also gives good results (Castellani,¹¹⁵ Dudgeon¹²²), while in the hands of Leishman himself it gives *better* results than Giemsa's stain. It must be acknowledged that the new Giemsa's stain, in containing methyl alcohol as a solvent, approximates to Leishman's stain which existed before it. Of Giemsa's new solution 15 cubic centimetres are added to 10 cubic centimetres of distilled water. The films which have been previously fixed in alcohol are stained for an hour at room temperature. The films may be

² THE LANCET, May 13th, 1905, p. 1259.

* The "superior," or raised, figures occurring throughout refer to the bibliography at the end of the article.

left in the stain over-night. After washing and drying with filter paper they are examined in cedar oil or balsam. Details are given in his papers.^{117 118} Davidsohn¹²³ strongly recommends cresyl violet, Herxheimer⁴⁵ Nile blue, and later⁴⁷ an aqueous solution of gentian violet after preliminary fixation in alcohol. Hoffmann,⁶ Oppenheim and Sachs,⁸⁵ and Ploeger⁶² stain with anilin water or carbolic gentian violet, Poleger finding that preliminary fixation in alcohol is not necessary. Carbol-fuchsin is also used by several investigators. Marino's azur solution is used by others—e.g., Metchnikoff. Weitlauer¹²¹ claims success with Löffler's methylene blue. Bertarelli and Volpino,⁸⁸ Levaditi,²⁹ Paschen,¹¹¹ and Buschke and Fischer¹⁷ have succeeded in demonstrating spirochætæ in sections by silver impregnation methods and such as are used for the demonstration of the flagella of bacteria, while Reitmann,¹¹⁹ with a modification of Sclavo's method, and Proca and Vasilescu⁶⁰ with a modification of Pitfield's method for staining flagella have also been successful. In my experience the last gives good results. In addition to the authors named above, Schaudinn,⁷ Nicolas, Favre and André,⁸⁹ Burnet,⁹⁰ Bandi and Simonelli⁹² discuss and criticise the various staining methods.

Spirochætæ in other conditions.—Spirochætæ, sometimes approximating in form to spirochæta pallida but mostly differing from it, have been found, amongst others, by Polland¹²⁷ in hospital gangrene, by Buday¹²⁸ in noma and gangrenous tonsillitis, by Bonhoff¹²⁵ and Süpfle¹²⁶ in vaccine pustules, by Borrel⁸⁶ in several cases of mouse carcinoma, by Hoffmann and Mulzer¹¹ in ulcerated carcinoma, by Sobernheim and Tomaszewski⁷⁵ in cancer of the lips and balanitis, and by Kiolomenoglou and von Cube⁴⁹ in various venereal and non-venereal conditions. O. Moritz⁸⁷ found somewhat coarse spirochætæ in the bone marrow and in the muscular coat of the small intestine in severe anæmias and also in carcinomatous lymphangitis. He could not prove any relationship. As a rule all these spirochætæ except those described in the last two papers are readily stained and by other characters easily distinguishable from spirochæta pallida.

Gathering together the details from this not quite complete survey of the literature on the subject, and from my own experience, I may now piece together an account of the characters of these spirochætæ as at present understood. The names "spirochæta pallida," or "spirochæta pallidum," and spirochæta refringens are descriptive, the former staining faintly, the latter being markedly refractile in the living state. The *spirochæta pallida* is an extremely delicate organism, weakly refractile when living, vigorously motile, stained with considerable difficulty and difficult to demonstrate, requiring at least a magnification of 800, with good illumination. It is long, extremely thin and filamentous, of a distinct spiral or corkscrew shape, both when living and when fixed on the slide. It tapers at the ends to a sharp point. In length it varies from 4 to 20 μ ; in breadth it measures about 0.25 μ . The spirals number from six to 26, averaging from eight to ten. Not only are these spirals numerous but very narrow—not over about 1 μ —regular and deep. Schaudinn, Hoffmann, Herxheimer, and others describe terminal flagella, but all observers are not agreed upon this point. When these are double they may really represent the commencement of a longitudinal fission. The spirochætæ may lie in clumps and occasionally form a felted mass. Löwenthal thinks that they are made up of several shorter (from 3 to 4 μ) individuals, or possibly these may be degenerative forms. With Herxheimer, Mulzer, and others, he describes a granule or localised swelling between the middle and one terminal third, which may represent blepharoblast rather than nucleus. Some describe a terminal globular body, the real nature of which has not yet been ascertained. It may simply be a loop at the end of the spirochæta. Herxheimer has described other bodies in, on, or near the spirochæta. At first it was thought to possess a vibratile membrane attached throughout its whole length, but this idea is given up by most authors, with the exception of Dalous, the appearance being put down as artefact. The absence of such membrane along with the possession of flagella are held to be characters distinguishing it from other spirochætæ which commonly occur on ulcerated surfaces. Its motility is of three kinds: corkscrew, to and fro, and bending or lashing. In a film dried on the surface of glass the spirals may become stretched out owing to its becoming attached first at its extremities. By silver impregnation methods it has been demonstrated in sections of the

liver in congenital syphilis and in the primary and secondary stages of the acquired disease. It lies in the capillaries of the papillæ of the corium, chiefly lying up against the endothelium or in the lumen. Thence it makes its way into the surrounding tissues and works up between the cells of the Malpighian layer to reach but not to penetrate the stratum corneum. With Giemsa's stain, it takes a rose-pink colour, or slightly lilac if the staining be too prolonged. It may also be demonstrated by "flagella" stains and impregnation methods, by gentian violet, carbol fuchsin, cresyl violet, Nile blue, &c. Schaudinn places spirochæta pallida with the protozoa rather than with spirillar bacteria, allying it to the trypanosome group.

The *spirochæta refringens* is generally acknowledged to be of saprophytic nature and corresponds to the forms already described in normal smegma, in balanoposthitis, vulvo-vaginitis, on the surface of ulcers about the genitals, on venereal warts, in the mouth secretions, on the teeth, tonsils, in ulcerated carcinomas, and probably also the forms found in gangrene are closely related. It is thicker than spirochæta pallida, distinctly refractile. It may have a marked corkscrew shape and motion when alive but flattens out when dried in films, so as to have usually a wavy outline. The curves are flatter and longer than those of spirochæta pallida, measuring from 1.5 to 2 μ or more. It varies in length like the spirochæta pallida, measuring most commonly from 8 to 10 μ . It is usually described as having truncated ends but often it has tapering ends or one end is sharp and the other blunt. It possesses a vibratile membrane but no flagella. It stains easily, quickly, and deeply by all methods and with Giemsa's stain which is the only differential one as yet known it takes a purplish or violet colour. It occurs usually only on the surface of moist and ulcerated syphilitic lesions. Some describe it as invading the glands.

Demonstration of these parasites.—From the literature I have picked out the details as to 758 lesions examined by different workers and find that spirochæta pallida has been found most often in moist papules, mucous plaques, and condylomata, genital and extra-genital—159 positive and 26 negative. Next come hard chancres, excised or scraped—150 positive and 55 negative; non-ulcerated papules—92 positive and 31 negative. Success has not been as frequent in films made from excised or punctured buboes—52 positive and 30 negative—and rarely has spirochæta pallida been found in roseolar blood—four positive and 17 negative—or in the circulating blood—five positive and 37 negative.

In lesions with moist or ulcerating surfaces the spirochæta pallida is commonly accompanied by the spirochæta refringens and it is a very significant fact that most successes have been attained in such circumstances. From one's own experience the number of forms intermediate between the two, partaking to some extent now of the one, now of the other, is remarkable. It is sometimes extremely difficult, often impossible, to refer these forms to the one group or the other and it is easy to imagine how readily one's percentage of successful demonstrations of the spirochæta pallida in such lesions might be raised by having regard solely to tinctorial or morphological characters. Frequently, in perfectly stained Giemsa preparations I have seen forms, staining pink like spirochæta pallida, sometimes delicate but wavy or almost straight in outline like the spirochæta refringens and, moreover, not in isolated examples which might be explained as stretched spirochætæ pallidæ, but in very considerable numbers. Other forms stain deeply like spirochæta refringens but they have a spiral outline like typical spirochætæ pallidæ. As a rule, these forms are thicker than spirochætæ pallida and do not present so much difficulty as the other class of intermediate forms. Still, the fact remains that some authors figure such forms as spirochætæ pallidæ. As group characters one must take both tinctorial and morphological characters into consideration. The refinements of technique necessary to demonstrate flagella are too great to make their presence a point of great importance for the ordinary observer. When their presence is more widely confirmed, the methods of demonstration simplified, and staining results more constant, they might be of use in the diagnosis of the intermediate forms just referred to. It is regrettable that in condylomata and mucous plaques, the most infective lesions in the adult, these difficulties of diagnosis should be experienced.

Before making films from such lesions it is well thoroughly to cleanse the surface, as the better this is done the fewer saprophytic spirochætæ are met with. This refers also to hard sores, ulcerated papules, and rupioid or crusty

syphilides. After cleansing the surface it should be scraped fairly deeply with a sharp spoon, the bleeding allowed in great measure to stop, and films (from six to eight) made from the material scraped away and from the bloody fluid which exudes. Some authors dilute these products previously to making the films with a drop or two of distilled water.

Hard chancres and papules, whether ulcerated or closed, may be excised, cut open from their under surface, and the expressed juice smeared over slides or treated by scraping, as in making films from moist or ulcerated lesions.

Blood.—Though several observers have found scanty spirochætæ in the blood during life, either by puncture of the skin or of the spleen, it is necessary in general to take a larger quantity of blood than is available by these means and to treat it in the manner indicated first by Noeggerath and Staehelin. They adopted the method employed for the detection of trypanosomes in circulating blood. They take at least one cubic centimetre of blood from an arm vein, mix it with ten cubic centimetres of one-third per cent. acetic acid, centrifugalise and examine the deposit. One may remark that it is also possible, judging by analogy from the fact that many trypanosomes being of less specific gravity than the corpuscles remain in the supernatant fluid, that similarly all the spirochætæ present may not be sent down by centrifugalising. Roseolar eruptions may be scraped or punctured and the blood examined, or following Levaditi and Petresco³⁰ small blisters may be raised over them by means of cantharides and the fluid examined after centrifugalising. Several authors—e.g., Bodin¹¹⁰—report the failure of Levaditi's method in their hands. If, as some think,¹⁰⁸ this lesion is caused by impaction of plugs of spirochætæ in the capillaries of the papillæ they ought to be more easily demonstrable than they are at present.

Indurated buboes may be excised or punctured with a fairly stout hypodermic needle. After the usual antiseptic precautions the gland is punctured and the point of the needle kept near its surface so as if possible to receive some lymph from the peripheral sinus. The gland meanwhile should be massaged between the finger and thumb of the other hand. In view of the fact that some observers mix scrapings with a drop or two of distilled water to facilitate securing a thin film it is probably not of importance to insure the syringe being absolutely dry before using.

Congenital syphilis.—It is now a well-established fact that spirochætæ pallida occurs very constantly in the blood and tissues of infants suffering from congenital syphilis. It has been found in the blood during life and in the organs after death, especially in the liver, spleen, adrenals, kidneys, bone marrow, lungs, and even once in the cerebro-spinal fluid. In the bullæ of pemphigus occurring in such infants Levaditi, Salmon, Hoffmann, and others have found spirochætæ abundantly, often in clumps apparently the result of agglutinative action. This is very important in consideration of the great infectivity and virulence of congenital syphilitic lesions and is a fact greatly favouring an etiological relationship between spirochætæ pallida and syphilis, as is also the fact that in tertiary syphilis, which is little or non-infective, most authors fail to find spirochætæ pallida at all.

Fixation.—Most authors agree that the films, after drying in the air, should be fixed in absolute alcohol or alcohol-ether mixture for from ten minutes to 20 minutes or methyl alcohol for about five minutes. Schaudinn recommends that the films as soon as made should be exposed for two or three seconds to the vapour of osmic acid before drying and fixation in alcohol. For staining Giemsa's medium is most used. It is related to Leishman's modified Romanowsky's stain, which gives as good results when one is expert in its use. Modifications of Pitfield's flagella stain—e.g.⁶⁰—give good results, but the spirochætæ appear thicker and coarser as well as darker than by other methods. This is still more evident in films and sections treated by silver impregnation methods.^{29 33 17} Schaudinn demonstrates the terminal flagella by Löffler's special method. Films are mounted in cedar oil or balsam.

The following points may be taken as axioms to be borne in mind. (a) Saprophytic spirochætæ are always present on ulcerated lesions, on moist papules and mucous plaques. These are often difficult to distinguish from spirochætæ pallida. (b) The more thorough the preliminary cleansing the fewer of these seen. (c) Spirochætæ pallida may be somewhat deeply in the lesions and apparently in clumps. (d) On account of this one must make six to eight films. (e) A little blood is usually an advantage in the film as it assists in focussing and,

moreover, spirochætæ are often seen in apparent relationship to red blood corpuscles. (f) In the case of excised chancres and papules it is well to bear in mind that spirochætæ pallida is not demonstrable after from six to ten hours, even though the tissue has been kept on ice. They soon degenerate. (g) The films should be thin and should be fixed as soon as possible. (h) When using Giemsa's method the nuclei of the leucocytes should be stained of a deep brownish purple, almost black, colour, otherwise the spirochætæ will not be seen. (i) Decolourisers such as alcohol or weak acetic acid should, if possible, not be used as they tend rapidly to discharge the stain from spirochætæ pallida. This does not apply when the mordanting (flagella) methods are employed.

Cultivation.—Several attempts have been made to cultivate these spirochætæ but so far without success; judging, however, from the successful cultivation of trypanosomes this should not be impossible. De Souza and Pereira⁸⁰ used human blood with 5 per cent. each of citrate of sodium and chloride of sodium. Bertarelli and Volpino³³ added gland juice to human blood with 50 per cent. of citrate of sodium, incubating at 27°, 30°, and 37° C.

Control observations.—A very large number of these have been made by numerous workers. Most agree that spirochætæ pallida is not found in gonorrhœal pus and buboes, epididymitis, soft chancres and associated buboes, phagedænic ulcer, venereal warts, non-syphilitic ulcers of the glans, inflamed phimosis (except in one case in which a masked hard chancre was suspected), balanitis, and balanoposthitis; nor in smegma in man or woman, scabies, iodide rash, verruca juvenilis, scrofuloderma and lupus, phthisical sputum, furunculosis, acne, sycosis, dental suppuration, acute dermatitis, dermatitis herpetiformis, herpes zoster, eczema acute and chronic, eczema seborrhœicum, eczema papulo-crustosum, erythema exudativum multiforme, erythema nodosum, lichen ruber planus, psoriasis, pemphigus neonatorum, molluscum contagiosum, impetigo simplex and contagiosum, sarcoma and carcinoma of the penis and labia, epithelioma cutis, carcinoma of the lip, the rectum, the neck, and the upper jaw, fibroma of the mamma, hypertrophy of the tonsils, Vincent's angina, throat mucus and mouth secretions, normal skin, organs of newly born non-syphilitic children, and many other pathological and non-pathological tissues.

My own investigations embrace a comparatively small number of cases, the greater number of which have been obtained for me by Dr. Alexander M. McIntosh, clinical assistant in the lock wards of the infirmary, who has also assisted me in the microscopical investigation of several of them. I am also indebted to him for furnishing me with histories of most of the cases.

CASE 1.—A male, aged 18 years. He had had six weeks' illness. There were ulceration under the prepuce and gonorrhœa. On admission in October, 1905, there was a roseolar and papular rash over the whole of the body. There were œdema of the prepuce and a concealed hard chancre near the corona. The indurated groin glands were tender to the touch. Treatment consisted of mercurial inunctions; the prepuce was slit up, &c. The patient was readmitted at the end of November with gonorrhœal arthritis. Spirochætæ pallida was found in the scrapings from a non-ulcerated papule on the thorax. The fluid from the groin gland proved negative. (Stain: Giemsa.)

CASE 2.—A male, aged 25 years. He had had frequent connexions, the last about three weeks before admission. There were sores on the penis and a rash appeared a few days later on the trunk and extremities (papular syphilide). There were mucous plaques over the pharynx. There were indurated buboes. On examination fluid from the glands gave a negative result. In scrapings from the hard chancre spirochætæ pallida was present along with spirochætæ refringens. Scraping from the dry papulo-squamous syphilide proved negative. (Stain: Giemsa.)

CASE 3.—A male, aged 22 years. Infection occurred about 12 months before admission. There were typical peri-anal condylomata. Scrapings from these showed numerous spirochætæ refringentes and many intermediate forms, many delicate, staining pink with Giemsa, but straight or wavy in contour. Some forms were thin, with narrow spirals, but staining of a bluish or purplish colour.

CASE 4.—A male, aged 19 years. There was a two months' history. There was a diffuse macular and papular rash over the face and the body. The prepuce was œdematous and there was a concealed hard chancre. The glands were

FIG. 1.

Spirochæta pallida in apparent relation to a red blood corpuscle. Stained by Giemsa's solution. The organism appears in the centre of the field, sloping downwards from left to right.

FIG. 2.

Spirochæta pallida in relation to red blood corpuscles. Reproduction from an untouched negative. Film stained with Rossi's stain. One-twelfth 4 oc. Zeiss. Bellows 35 centimetres. The organism appears at the extreme top edge of the illustration, just under the figure 2.

FIG. 3.

Film from condyloma latum showing numerous forms, mostly intermediate between *spirochæta pallida* and *spirochæta refringens*. Numerous bacteria stained by Giemsa's solution.

FIG. 4.

Spirochæta refringens. Two examples scraped from a syphilitic hard sore. Stained by Giemsa's solution.

FIG. 5.

Peculiar flagellated body found in film made from fluid expressed from the cut surface of a hard chancre. Stained by Giemsa's solution. $\times$ circ. 1200.

FIG. 6.

Peculiar loops and roundish bodies found not uncommonly in smears from lymph glands. Loops stained by Giemsa's solution (red). Roundish bodies stained by Giemsa's solution (blue).

not typical. Treatment consisted in circumcision and mercurial inunctions. Films made from the excised chancre contained a few spirochætæ pallidæ and a peculiar flagellated body. The films were examined by both Dr. McIntosh and myself. The body stained a brownish colour with Giemsa's stain and in it a clear spot resembling a vacuole was readily distinguished. It had at least six angles and to these whip-like appendages resembling spirochætæ pallida in tinctorial and morphological characters were attached. In the case of two of these the thickness was greatest near the body and gradually lessened as one passed out. The appearance was evidently not an artefact and the staining of the central mass was not like that of red blood corpuscles in the film. As to its nature one can make no suggestion and I have not seen another body like it. It does not altogether look like a red blood corpuscle to which spirochætæ are attached and the hæmoglobin of which is diffusing along their proximal portions. If it is of importance in connexion with spirochætæ pallida it is curious that no other observer has seen an exactly similar

from the macular eruption proved negative. Films from the excised hard chancre contained numerous spirochætæ pallidæ. (Stain: Giemsa.)

CASE 7.—A male, aged 38 years. A hard chancre appeared five weeks before admission. There was a typical hard chancre on the dorsum of the penis. There were typical groin glands and papular syphilide. This was one of the first cases I examined. I made only three films from the chancre and could not find spirochætæ. (Stain: Giemsa.)

CASE 8.—A male, aged 23 years. 13 days after the last connexion an ulcer appeared on the right side of the glans and rapidly enlarged, eating into the glans deeply. There was no glandular enlargement or induration. There was no rash. The ulcer healed rapidly under treatment with lotio nigra. Films from the ulcer contained numerous spirochætæ refringentes. There were no spirochætæ corresponding to spirochætæ pallida. (Stain: Giemsa.)

CASE 9.—A female, aged 32 years. There was a doubtful ulcer at the fourchette. Films made from this contained numerous bacteria and spiral forms most corresponding

FIG. 7.

a-i $\times$ about 1000, *k* $\times$ about 1200. *a*, Spirochætæ pallidæ—*a*₁, in relation to red blood corpuscles; *a*₂ crossing an oblong body (particle of chromatin?) and apparently attached to one of the angles; *a*₃, crossing red blood corpuscles showing apparent thickening. *b*, Spirochætæ refringens. *c*, Fusiform body like bacillus fusiformis, frequently seen in the ulcerated lesions. *d*, Intermediate forms from condyloma latum. *e*, Artefacts, drawn-out nuclei, &c. *f*, A thin filament of fibrin (?) simulating spirochætæ pallida. *g*, Wrinkled margin of deformed red blood corpuscle which might be mistaken for spirochætæ pallida encircling it. *h*, Peculiar loop-shaped bodies staining bright pink with Giemsa's solution found in several of the films from the glands. Their relationship to the spirochætæ though possible is not clear. *i*, Rounded bodies (Winckler's?) in gland fluid. They stain a pale blue with Giemsa's stain. *k*, Peculiar body found once in hard chancre. Note the fusiform thickening seen in the course of the flagella-like extensions.

object. This very fact tends to make one very sceptical as to its importance. Still, it is necessary to put the fact on record and a micro-photograph of it is reproduced. (Figs. 5 and 7.) In the same cases numerous spirochætæ pallidæ were found in the scrapings from a closed papular syphilide. (Stains: Giemsa and modified Pitfield's flagella.)

CASE 5.—A male, aged 22 years. Infection occurred seven weeks before admission. There were a concealed hard chancre and a diffuse and roseolar rash. There were typical indurated groin glands. Treatment consisted in circumcision and mercurial injections. On examination scrapings from the macular eruption proved negative. Films from the excised hard chancre contained numerous spirochætæ pallidæ. (Stains: Giemsa, modified Pitfield's flagella, and carbolic gentian violet.)

CASE 6.—A male, aged 19 years. Infection occurred from five to eight weeks before admission. There were a hard chancre on the prepuce and a macular eruption; the indurated groin glands were typical. Circumcision was performed and mercurial injections were given. Scrapings

morphologically to spirochætæ refringens, but a few delicate with narrow spirals like spirochætæ pallida. On account of this result I asked Dr. McIntosh to watch this patient carefully. About a week later a typical macular rash appeared on the thighs and thorax, the primary sore becoming quite typical. (Stain: Giemsa.)

CASE 10.—A male, aged about 24 years. He had had syphilis seven or eight years ago. Films made from scrapings of the sores contained numerous intermediate forms, some of which were apparently spirochætæ pallidæ and numerous forms corresponding to spirochætæ refringens.

CASE 11.—A female, aged 39 years. There were condyloma, papular rash, and indolent buboes in the groin and behind the sterno-mastoid. Films were brought to me which had been made from the condyloma. No spirochætæ pallidæ were found in six films.

CASE 12.—A male, aged 22 years. The last connexion was 11 days before admission. He was discharged on the ninth day. He had œdema of the prepuce, phimosis, and concealed soft chancre. There were no enlargement of the

glands and no rash. Circumcision was performed, removing the sore. Films showed no spirochætæ like spirochætæ pallida.

CASE 13.—A male, aged 24 years. The patient was infected with gonorrhœa and soft sores about eight months before admission. Now he had induration at the preputial orifice. There was a diffuse macular rash on the body with a few papules on the penis and pubes. Typical indolent buboes were present in the groin and behind the sterno-mastoid. The hard chancre was excised. Films were made from scrapings from the surface and fluid from the cut surface. In eight films no spirochætæ pallidæ were discovered.

CASE 14.—A male, aged 35 years. There was a three and a half months' history. There was a papulo-squamous syphilide on the legs and trunk which was drying up under treatment. He was twice examined with a negative result.

CASE 15.—A male, aged 20 years. There was a three months' history. He had perianal condylomata which were now drying up under local and general treatment. A few spirochætæ refringentes were found but no spirochætæ pallidæ.

CASE 16.—A male, aged 24 years. Infection occurred ten months before admission. A sore formed which never completely healed. Connexion meanwhile was almost fortnightly. Three weeks before admission the sore broke down extensively and became indurated and indolent. The glands were enlarged and hard. There was a papular rash. From fluid from the glands films were made. No spirochætæ pallidæ were found but a few peculiar tailed bodies like spermatozoa. These were possibly drawn-out nuclei of leucocytes. (Fig. 7, e.)

CASE 17.—A male, aged 19 years. Infection occurred four weeks previously to admission. On admission the patient had a concealed hard chancre. Circumcision was performed. Four weeks later (December, 1905) there were a dry parchment sore on the glans which was healing, and indurated groin glands. Scrapings from the sore on the glans showed presence of spirochætæ refringens and some intermediate forms, some of which approximated in form but not in colour to spirochætæ pallida. Some of them had a small dark body attached at one end. Fluid from the groin glands contained no demonstrable spirochætæ but numerous flattened or oval loop-like structures, staining bright pink with Giemsa's solution, were seen (Figs. 6 and 7). Some of these were shaped like the "eye" of the "hook and eye" of the dressmaker. In the same case there were rounded bodies varying in size from 1μ to 4μ , staining a delicate blue with Giemsa, sometimes single, sometimes in clusters, as if they had been contained within a phagocytic cell which had broken up in the process of making the film. These were probably products of the degeneration of cells. They are possibly analogous to the bodies described by Winckler. Some of these can be made out in Fig. 6.

CASE 18.—A male, aged 20 years. There was a two months' history. He had a general rupial eruption over the whole of the body resembling impetigo at parts. Scrapings from the bases of these lesions and fluid expressed from them failed to show the presence of spirochætæ pallida.

CASE 19.—A male, aged 22 years. There was a 13 months' history. Treatment had been irregular. There were peri-anal condylomata. Films were made from the condylomata but no spirochætæ pallidæ could be detected.

CASE 20.—The hard chancre of another male was examined and numerous spirochætæ pallidæ were discovered. No history of the case is now obtainable.

CASE 21.—A male, aged 28 years, suffering from tabes dorsalis. Symptoms began three months after infection with syphilis three years ago. Spinal fluid was removed by lumbar puncture. No spirochætæ were found. On cultivation, in one tube a pure culture of streptococci developed.

Of these 21 cases, two (8 and 12) were not syphilitic; another (21) was a case of tabes dorsalis three years after infection. Excluding these there are left 18 cases showing 25 lesions examined. Out of the 18 cases six showed the presence of undoubted spirochætæ pallida. In three others spirochætæ were found which could not with certainty be placed either with spirochætæ pallida or spirochætæ refringens. One case (7) was examined when I had little experience in looking for spirochætæ pallida. In one case (9) the possibilities are in favour of the bodies seen being spirochætæ pallidæ. In two cases (14 and 15) the lesions were drying up under treatment. The results were quite negative in the remaining five cases.

As regards the lesions, five hard chancres showed positive results; in two the results were doubtful and in three they were negative. Two non-ulcerated papules gave positive results; two dry papulo-squamous syphilides gave negative results, and two roseolar rashes also were negative. Four condylomata were examined; three were negative and one was doubtful. Four glands were examined and in none could undoubted spirochætæ pallidæ be found. One rupioid syphilide was examined with a negative result. Typical spirochætæ pallidæ were found only in hard chancres and in closed papules—i.e., in typical syphilitic lesions.

CONCLUSIONS.

It is astonishing with what unanimity this discovery has been accepted on the continent, an occurrence quite unprecedented in discoveries of similar importance, but there is much to support the relationship of spirochætæ pallida to syphilis. The most important facts in favour are its occurrence in non-ulcerated lesions, such as papules and buboes far away from the primary sore; its occurrence in the circulating blood and splenic blood in acquired syphilis, its occurrence specially in virulent infective lesions, and its absence from non-infective tertiary lesions; its presence in the depths of superficial lesions where it is stated to be unaccompanied by other organisms, and its definite localisation in the corium—if absolutely confirmed; its presence in congenitally syphilitic infants in great numbers and its occurrence—according to most observers—only in primary and secondary syphilitic lesions and in congenital syphilis—never in non-syphilitic diseases. Against it is its frequently scanty numbers, films in very many of the reported successful cases revealing the presence of only one or two spirochætæ, after repeated and prolonged search, the numbers thus frequently bearing no relationship to the severity of the tissue reactive changes, which is, as Siebert points out, without analogy in the domain of pathology; the possibility, which is by no means disproved, that various organisms may invade glands penetrating from a superficial ulcer; the difficulty of diagnosis in the case of the numerous intermediate forms which are found so commonly in such highly infective lesions as the mucous plaque, &c.; the claim, still strongly upheld by some, that forms indistinguishable from spirochætæ pallida are found in non-specific lesions. The last point, of course, even though it were granted, does not necessarily invalidate a causative relationship to syphilis.

In conclusion, one may say that while there is a very considerable probability that spirochætæ pallida bears an etiological relationship to syphilis we require to know more about spirochætæ in general and to have better methods of distinguishing the many forms intermediate between the typical spirochætæ pallida and the typical spirochætæ refringens before we can unreservedly accept this as a completely proved fact in pathology. One can say very little as yet with regard to the peculiar loop-like and rounded bodies found in syphilitic glands and to such bodies as the flagellated structure I have described in a film from a hard chancre.

Bibliography.—[B. K. W. = Berliner Klinische Wochenschrift; D. M. W. = Deutsche Medicinische Wochenschrift; W. K. W. = Wiener Klinische Wochenschrift; M. M. W. = Münchener Medicinische Wochenschrift; B. M. J. = British Medical Journal; and Cpt. Rend. = Comptes Rendus Hebdomadaires de la Société de Biologie.]
¹ Horand: Ref. ³⁷ and Lyon Médical, 1905, 23, 1223. ^{2 3 4} Siegel: M. M. W., 1905, 1321, 1384. ⁵ Schaudinn and Hoffmann: Arbeiten aus dem Kaiserlichen Gesundheitsamte, 1905, xxii., 2, 527. ⁶ Ibid., B. K. W., 1905, 22. ⁷ Ibid., D. M. W., 1905, 18. ⁸ Ibid., D. M. W., 1905, 20. ⁹ Schaudinn: D. M. W., Oct. 19th, 1905. ¹⁰ Hoffmann: B. K. W., 1905, 23. ¹¹ Ibid., B. K. W., 1905, 28. ¹² Ibid., B. K. W., 1905, 42, 1022. ¹³ Ibid., B. K. W., 1905, 46, March. ¹⁴ Ibid., D. M. W., 1905, 51, 1070. ^{15 16} Buschke and Fischer: D. M. W., 1905, 20, 791; 21, 839. ¹⁷ Ibid., B. K. W., 1906, 1, 6. ¹⁸ Metchnikoff and Roux: Annales de l'Institut Pasteur, November, 1904, 761. ¹⁹ Ibid., Bulletin de l'Académie de Médecine, 1905, p. 468. ²⁰ Ibid. (ref.), M. M. W., 1905, 24. ²¹ Ibid. (4th Mem.), Annales de l'Institut Pasteur, November, 1905. ²² Metchnikoff: Archives Générales de Médecine, 1905, 26, 1623. ²³ Raubitschek: W. K. W., 1905, 28, 752. ²⁴ Frosch: ref. ⁹ and ⁵². ²⁵ Reckzeh: ref. ⁵². ²⁶ Weichselmann and Löwenthal: Med. Klin., 1905, 25. ²⁷ Levaditi: La Semaine Médicale, May 24th, 1905. ^{28 29} Ibid., Cpt. Rend., 1905, 243, 845. ³⁰ Levaditi and Petresco: Presse Méd., Par., 1905, 78, 617. ³¹ Salmon: La Semaine Médicale, 1905, 247. ³² Ibid., Cpt. Rend., 1905, 19, 883. ³³ Jacquet and Sevin: ref. ³⁹. ³⁴ Kraus: W. K. W., 1905, 22, 592. ³⁵ Volk: ref. ³⁹. ³⁶ Schüller: Centralblatt für Bakteriologie, Ref., xxxvi., 24, 25, 1905 (gives references to his papers). ³⁷ Vuillemin: Comptes Rendus de l'Académie des Sciences, 1905, cxl., 23, 1567. ³⁸ McWeeney: B. M. J., 1905, i., 1262. ³⁹ Lesourd: Annales de Dermatologie et de Syphilographie, June, 1905. ⁴⁰ C. Fraenkel: M. M. W., 1905, 24, 1129. ⁴¹ V. Jensen: Hospitalstidende, Nr. 25, 1905. ⁴² Bordet and Bayet: Presse Méd. Belge, 1905, 26, 614. ⁴³ Bordet and Gengout: Soc. Roy. de Sc. Brux., 63, 124, 1905. ⁴⁴ Lassar: Dermat. Zeitsch., 1905, 6, 418. ⁴⁵ K. Herxheimer and Hübner: D. M. W., 1905, 26, June 29th. ⁴⁶ Herxheimer and Löser: M. M. W., 1905, 46, 2212. ⁴⁷ Herxheimer: M. M. W., 1905, 39. ⁴⁸ Hübner: Dermat. Zeitsch., xii., ii., Nov., 1905.

⁴⁹ Kiolomenoglou and v. Cube: M. M. W., 1905, 27. ⁵⁰ ⁵¹ Babes and Panea: B. K. W., 1905, 28 and 48. ⁵² Galli-Valerio and Lassueur: Rev. Med. d. l'Suisse Rom., 1905, July 20th. ⁵³ Krzyszalowiec and Siedlecki: Monatschrift für Praktische Dermatologie, 1905, 41, 6, 231. ⁵⁴ Thesing: D. M. W., 1905, 32, August 10th; ⁵⁵ M. M. W., 1905, 28, 1337. ⁵⁶ Borrel: Cpt. Rend., 1905, 16. ⁵⁷ Moritz: Petersburger Medizinische Wochenschrift, 1905, 20; and Deutsches Archiv für Klinische Medizin, Band 84, 1905. ⁵⁸ Rille: M. M. W., 1905, 29, 1377. ⁵⁹ Rille and Vockerdt: M. M. W., 1905, 34. ⁶⁰ Proca and Vasilescu: Cpt. Rend., 1905, 23. ⁶¹ Noeggerath and Staehelin: M. M. W., 1905, 31, 1481. ⁶² Ploeger: M. M. W., 1905, 29, 1381. ⁶³ Jesionek: M. M. W., 1905, 49, 2394. ⁶⁴ R. Kraus and Prantschhoff: W. K. W., 1905, 37, 941. ⁶⁵ Bandler: Prager Medicinische Wochenschrift, 1905, 34, 475. ⁶⁷ Leiner: ref. 68. ⁶⁸ Lannois: ref. 66 and M. M. W., 1905, 31, 1518. ⁶⁹ Almkvist and Jundall: Allgemeine Schwed. Aerztezt., 1905, No. 25, and ref. 66. ⁷⁰ Mulzer: B. K. W., 1905, Sept. 4th (quotes Csillag, Rona, and Menge-Krönig.) ⁷¹ ⁷² Nigris: D. M. W., 1905, 36, 51. ⁷³ Scholtz: D. M. W., 1905, 37, Sept. ⁷⁴ Grouven and Fabry: D. M. W., 1905, 31, 1469. ⁷⁵ Sobernheim and Tomaszewski: M. M. W., 1905, 39, 1857. ⁷⁶ Siebert: D. M. W., 1905, 41. ⁷⁷ Fligel: D. M. W., 1905, 44. ⁷⁸ Brönnum and Ellermann: M. M. W., 1905, 2093, 2241. ⁷⁹ Roscher: B. K. W., 1905, 44, 45, 46. ⁸⁰ De Souza, jun., and Pereira: B. K. W., 1905, 44. ⁸¹ Queyrat: M. M. W., 1905, 29, 1420. ⁸² Queyrat and Joltrain: see 81 and M. M. W., 1905, 31, 1518. ⁸³ Reischauer: D. M. W., 1905, 34, August 24th. ⁸⁴ Neumann: M. M. W., 1905, 38, 1848. ⁸⁵ Oppenheim and Sachs: D. M. W., 1905, 29 and 31. ⁸⁶ Ibid., M. M. W., 1905, 1507, 1517. ⁸⁷ Ibid., W. K. W., 1905, 45, Nov. 9th. ⁸⁸ Bertarelli and Volpino: Centralblatt für Bakteriologie, O., 1905, Nov. 24th. ⁸⁹ Nicolas, Favre, and André: Lyon Médical, cv., 1905, 497. ⁹⁰ Burnet: Annales de Dermatologie et de Syphilographie, 1905, xi., 833. ⁹¹ Spitzer: W. K. W., August 3rd, 1905. ⁹² Bandi and Simonelli: Gazzetta degli Ospedali e delle Cliniche Mil., 1905, 85 and 105. ⁹³ Ibid., Cent. f. Bakt. O., Nov. 24th, 1905. ⁹⁴ Fanoni: Medical News, New York, 1905, 88, 678. ⁹⁵ Taylor and Ballenger: Journal of the American Medical Association, 1905, 20. ⁹⁶ Gordon: Am. Med., July 22nd, 1905. ⁹⁷ Paltauf: ref. 39. ⁹⁸ Lipschutz: D. M. W., 1905, 46. ⁹⁹ Pielicke: ref. 89. ¹⁰⁰ Richards and Hunt: THE LANCET, Sept. 30th, 1905, 962. ¹⁰¹ Dalous: Jl. Mal. Cut. et Syph., 1905, xvii., 481. ¹⁰² Zabolotny: ref. 14, and M. M. W., 1905, 1698. ¹⁰³ Neisser (with Baermann and Halberstädter): D. M. W., 1906, 1, 2, 3. ¹⁰⁴ Thibierge and Ravant: Annales de Dermatologie et de Syphilographie, July, 1905. ¹⁰⁵ Pollio and Fontana: Gazzetta degli Ospedali e delle Cliniche, Mil., 1905, 109, 1143. ¹⁰⁶ Rizzo and Cipollina: Riforma Medica, 1905, 34, 938. ¹⁰⁷ Pascalis: Il Policlinico, Rome, 1905, 12, 861. ¹⁰⁸ Veillon and Girard: Presse Médicale, Paris, 1905, 104, 839. ¹⁰⁹ Moncorvo fils: ref. Presse Médicale, Paris, 1905, 104, 840. ¹¹⁰ Bodin: Presse Médicale, Paris, 1905, 102, 823. ¹¹¹ ^{111a} Paschen: M. M. W., 1905, 932, 2394. ¹¹² Harvey and Bousfield: Journal of the Royal Army Medical Corps, 1905, ii., 263. ¹¹³ Yaws.—¹¹³ Castellani: Journal of Tropical Medicine, August, 1905. ¹¹⁴ Ibid., B M J., 1905, Nov. 16th. ¹¹⁵ Ibid., D. M. W., 1906, 4, 132. ¹¹⁶ Wellmann: Journal of Tropical Medicine, 1905, Dec. 4th. ¹¹⁷ Hutchinson: Sydenham Society's Atlas. *Staining*.—See 7 45 60 85 87 8 39 90 12 113. ¹¹⁷ Giemsa: D. M. W., 1905. 26 (with references). ¹¹⁸ Ibid., D. M. W., 1905, 32. ¹¹⁹ Reitmann: D. M. W., 1905, 25, 997. ¹²⁰ Hirschberg: Journal of the American Medical Association, 1905, 45, 1086. ¹²¹ Weitlaue: M. M. W., 1905, 47, 2293. ¹²² Dudgeon: THE LANCET, August 19th, 1905, 522. ¹²³ Davidsohn: B. K. W., 1905, 31. ¹²⁴ Harvey: Journal of the Royal Army Medical Corps, 1905, iii., 409. *Spirochaete in other conditions*.—*Vaccinia*.—¹²⁵ Bonhoff: B. K. W., 1905, Sept. 4th. ¹²⁶ Süpfle: M. M. W., 1905, 38, 1848. *Gangrene, &c.*—¹²⁷ Pollard: W. K. W., 1905, Nov. 23rd. ¹²⁸ Buday: Ziegler's Beiträge, October, 1905. *Cytorrhyses Luis*.—See 2 3 4 and also ¹²⁹ Schülze: B. K. W., 1905, 32, 653. ¹³⁰ Freund: M. M. W., 1905, 1819. ¹³¹ Merk: W. K. W., 1905, 926.

Edinburgh.

RUPTURE OF THE UTERUS WITH ESCAPE OF THE CHILD INTO THE ABDOMINAL CAVITY; DELIVERY PER VIAS NATURALES; RECOVERY.

BY THOMAS HUNTER, M.B. GLASG.,
CAPTAIN, I.M.S.; CIVIL SURGEON.

THE patient was a Hindu of a labouring caste, aged about 24 years, a well-built healthy woman, with a history of four previous normal labours. From such patients a correct detailed history can rarely be obtained of what has occurred before the arrival of skilled observation, but apparently in this case labour came on at full term about 11 P.M. on Nov. 27th, 1905, and the pains were ordinary and the case usual until about 2 or 3 A.M. on the 28th. About that hour the waters burst and bearing-down pains began. These lasted a very short time and after not more than four bearing-down pains the pain completely ceased and about ten minutes after the cessation the placenta was delivered. With the birth of the placenta there was very profuse bleeding and as this continued the friends took the woman to the hospital. No history of injury at any time could be elicited.

On admission to hospital at 7 A.M. on Nov. 28th the patient was pale and collapsed and the pulse at the wrist could not be felt. Her temperature, which was at first 97° F., rose in the evening to 101.4°, and the pulse-rate was found to be 150 in the course of the day. She seemed

to be suffering some pain and there was extreme tenderness over the abdomen, palpation of which at once revealed the fact that the foetus seemed to be just under the abdominal wall. The placenta was lying outside the vagina attached to the cord. A hypodermic injection of strychnine, digitalin, and ergotin was given at once and the external genitals were thoroughly cleaned. Examination per vaginam was then made and a large irregular tear in the left posterior region of the uterus, and beginning apparently at the external os, was detected with the uterus firmly contracted behind the symphysis and the child in the abdominal cavity. A hot douche of sterilised normal saline solution was then given and the woman was put under chloroform. Examination then showed that the rent was large enough to allow the hand to pass freely into the abdominal cavity and to feel the child lying with the limbs near the rent.

The present-day opinion that the best treatment in such circumstances is laparotomy, cleaning of the abdominal cavity, and removal of the uterus, was borne in mind; but the collapsed condition of the patient was such that to have practised this, it was concluded, would inevitably have resulted in death. On the other hand, the natural passages were roomy and the tear was judged extensive enough to allow a normal child to pass with little further damage. At the same time it was remembered that, with no history of injury, abnormality in the child was almost certainly the cause of the rupture. The head of the child then was felt for and when it was reached the abnormality was found to be hydrocephalus. Perforation of the head and delivery per vias naturales were decided on. The feet were drawn together through the tear followed by the trunk with the upper limbs. The head, steadied over the tear by a hand in the vagina and two hands over the abdominal wall, was then pierced by a perforator. A rush of fluid followed and the child was easily delivered. After the delivery the abdominal cavity was thoroughly washed out with sterilised normal salt solution at a temperature of 100°. To do this a hand was passed in through the tear and the nozzle of the douche guided by it and with this hand in the abdominal cavity the abdominal contents were well washed. A piece of omentum which presented at the mouth of the vagina was replaced by the nozzle of the douche. Finally, a gauze drain was passed into the abdominal cavity through the tear, a pad was put on over the vagina, and a tight binder was applied.

The patient's condition was critical for the first two days, with a temperature rising to 103.4° F. and a pulse-rate of about 150. The gauze drain was changed and a hot douche of sterilised saline solution was given twice a day and the patient was kept on ten minim doses of liquor strychninae hypodermically every four hours. On the morning of Nov. 30th her condition was much better and her temperature was 100°. The saline douche was replaced by an izar douche. On the evening of the same day her temperature rose to 101° and tympanites developed but rapidly yielded to turpentine stupes. On Dec. 1st her temperature was 99.6° in the morning and 102° in the evening, the pulse-rate being about 120. On the 2nd diarrhoea supervened, remaining troublesome for two days but disappearing on treatment with five grains of Dover's powder and ten grains of subnitrate of bismuth every six hours. Her temperature was 99.2° in the morning and 100.2° in the evening. On the morning of the 3rd the temperature touched normal for the first time after delivery but there was still an evening rise—namely, to 99.0° on the 3rd, 4th, and 5th; to 101.0° on the 6th; to 99.8° on the 7th; and to 99.0° on the 8th. The temperature remained normal from the 9th and the progress of the case from that date was uneventful. The patient left hospital in good health on the 20th. She returned for examination on Jan. 10th, 1906, and the uterus was then found to be normal in size and position and the only remnant of the tear that could be detected was a laceration through the whole thickness of the cervix in the left posterior quadrant extending as far as the fornix.

As an interesting indication of the mental activities of the class to which the patient belonged it may be mentioned that when the child, which had a cleft palate and a double harelip, was seen by the relatives its deformities and its large head inspired them with awe and with the belief that it was a demon that had been delivered from the patient and the subsequent occurrence of tympanites caused great consternation as they believed that the demon had re-entered into possession of the mother.

Rae Bareli, Oudh.