tarded his recovery for some time; but on Sept. 3rd he was so much improved as to be able to leave the hospital for the infirmary at Margate. Complete union had taken place, but the knee was puffy and swollen, and there still remained both on the inner and outer side of the joint several small ulcerated patches, which were healing over. The shortening of the limb did not exceed one inch.

April 18th, 1863.—To-day the boy came to the hospital. His health has very much improved, and he is able to walk about all day. The left lower limb is quite straight, and about three-quarters of an inch shorter than the sound one. There is no pain about the joint, and all the ulcers have healed up.

LONG-STANDING DISEASE OF THE KNEE-JOINT; EXCISION;
FATAL RESULT.

(Under the care of Mr. FERGUSON.)

Jane D—, aged twenty-four, was admitted March 18th, with long-standing disease of the knee. She was a corpulent and hysterical-looking young woman. Before her knee became affected she was in general service. All her relations are healthy. Nine years ago she had some swelling about the left knee, which subsided after the application of iodine paint. Three years afterwards a large abscess formed, and pointed on the inner side of the joint, over the head of the tibia. The abscess was opened, and discharged a quantity of pus. This was followed by very severe and constant pain in the joint, which gradually became stiff and contracted, so that when the abscess had healed, and she was able to get up, she found that she could not put her foot to the ground. A year ago she was a patient in the Northampton Infirmary, where an attempt was made to straighten the limb, but with very little success. The knee had increased in size during the last three months, and the pain had been so severe as to keep her in bed.

Upon admission, the right knee was found to be much swollen and slightly bent; there was partial ankylosis; the patella was immovable, and thrown upon the outer condyle. At the upper part of the leg, below the head of the tibia, were the orifices of three sinuses, which extended upwards into the joint and communicated with bare bone. With the exception of these the skin was healthy, not red nor congested. Any attempt to move the joint gave her great pain. The muscles of the left calf were very much wasted, and the whole limb was somewhat smaller and less developed than the right one. The patient's health seemed very fair; her appetite was good; and she complained of very little save the occasional starting of the limb, which broke her rest at night. The limb was ordered to be kept at perfect rest, and to be fomented; the patient to take tincture of the sesquichloride of iron thrice a day, and cod-liver oil every morning.

March 28th.—The pain daily getting more intolerable, and exhausting her strength, Mr. Fergusson determined upon excising the joint, as the long duration of the disease left little to be hoped for from rest, or from any active treatment short of an operation. To-day, having been put under the influence of chloroform, she was brought from the ward into the theatre. Mr. Fergusson first made a single transverse incision over the patella; the flaps of skin being then dissected back, and the limb forcibly bent up at a very acute angle, the patella was removed, and the whole of the interior of the joint was exposed. About an inch and a half of the lower end of the femur was sawn off, and a thin slice from the head of the tibia. The soft tissues were found to be infiltrated with a dark, thick, gelatinous material, particularly at the posterior part of the joint, surrounding the popliteal vessels. The limb was then put straight, and bandaged down upon an excision splint, the ends of the bones coming very well into apposition. The margins of the incision were brought together by sutures, and water dressing applied. Two small arteries required ligatures during the operation.

The excised parts, when examined after the operation, were found to be most extensively diseased. There was a thickened, pulpy state of the synovial membrane, and the cartilage had nearly wholly disappeared from the femur and head of the tibia, leaving a surface of carious bone. The cartilage of the inner surface of the patella was ulcerated here and there, and large patches of bone were exposed. From the lower end of the femur there had sprung two large outgrowths of bone-tissue.

30th.—Was very restless after the operation, and suffered a great deal from pain and starting of the limb. Yesterday she felt very comfortable, and at night slept very well. This morning she looks well; pulse 126; tongue clean. The knee was dressed; the wound looking very healthy, and union taking

place along nearly the whole line of incision. To take six ounces of brandy daily.

31st.—Passed a restless night, and had slight rigors. Feels better this morning. Wound looking well; water-dressing applied.

April 2nd.—When seen to-day about noon was not so well. She is very restless, moaning in an hysterical manner, and complaining of great thirst; pulse 130; tongue red and dry. About six P.M. had a very severe attack of shivering, which was followed by constant vomiting. The knee when examined was found to be quite dry, and there was no discharge in the dressing. A large linseed-meal poultice was ordered to be applied directly, and to be changed every three hours. The patient to suck ice, and to have dilute hydrocyanic acid, with chloric ether, every four hours.

3rd.—The vomiting continued the whole of the night, and nothing could be kept on the stomach. About six in the morning the house-surgeon was called, and found her in a state of collapse. She rallied after some brandy had been administered, and complained of great pain in the abdomen, which was tympanitic. For the relief of this a large linseed-meal poultice was applied. In the evening she had slightly improved, but the sickness still continued.

4th.—Is better this morning; the sickness has nearly ceased, and she is able to take beef-tea with her brandy. Is at times delirious, and complains of severe pain in the left shoulder-joint. Tympanitis is much less, and there is some little discharge from the wound in the leg. Ordered to have ten ounces of brandy daily.

6th.—Much better. The knee at times is in great pain; there is no sickness, tympanitis, or pain in the abdomen.

8th.—Going on well up to last night, when she was very restless and delirious. This morning she had an hysterical fit. Complains of great pain in the shoulder, and seems in a very low condition.

9th.—To day she is rapidly sinking; has a feeble, quick pulse; complains of great pain in the head, and is at times delirious. About noon sickness and hiccough came on, with tympanitis. She gradually sank, and died at ten P.M.

Autopsy, sixty-five hours after death.—Putrefaction taking place rapidly. The body well nourished, and on cutting through the abdomen the subcutaneous layers of fat exceed two inches in thickness. The intestines, kidneys, and spleen were examined, and found healthy. The liver was of the normal size and weight, but its structure was very pale and fatty; the lower half of the right lobe was softened down, and the tissue infiltrated with a dark-coloured fluid. The uterus and ovaries were very much congested. On opening the chest, old pleuritic adhesions were found on both sides. At the apex of the left lung was an extensive deposit of tubercle, which in some parts was softening down. The heart was examined and found healthy. On the left shoulder-joint the synovial membrane was found congested, and about a tablespoonful of pus was let out of the joint. The head was not opened.

ST. MARY'S HOSPITAL.

(OPHTHALMIC DEPARTMENT.)

CASES OF MYDRIASIS TREATED BY THE SOLUTION OF OLD
CALABAR BEAN, A NEW OPHTHALMIC AGENT;
WITH CLINICAL REMARKS.

(Under the care of Mr. ERNEST HART.)

In calling the attention of the students to this new agent for obtaining contraction of the pupil, Mr. Hart said that the need for such an agent had long been felt in ophthalmic surgery, and that he had been experimenting for some time with various substances of which he had hoped that the local application might produce contraction of the circular fibres of the iris. With this view he had employed solutions of various alkaloids in glycerine, including morphine, strychnine, digitaline, and ergotine; but the results were not satisfactory. Dr. Argyll Robertson, of Edinburgh, however, had lately laid before the profession an account of the operation of solutions of the Calabar bean on the pupil, which showed it to possess the desired quality. Dr. Robertson's paper on this subject had been read before the Edinburgh Medico-Chirurgical Society on the 4th of February, 1863. It had been published subsequently in the *Edinburgh Medical Journal* for March. The original investigator of the properties of the bean was Dr. Christison, who had read an account of those properties before the Royal Society

of Edinburgh on Feb. 5th, 1855, in an important paper reprinted in the *Pharmaceutical Journal* in 1855; and subsequently Dr. Fraser had treated of it in his inaugural thesis. It was one of the most powerful poisons, possessing a direct control over the action of the heart, but not paralysing volition. Its systemic effects somewhat resembled those of aconite. The active principle had not yet been isolated; but Mr. Hart stated that he believed Dr. Christison was engaged in an effort to procure it; the difficulty at present was due to the scanty supply of the bean. It was held in great veneration by the natives of Calabar as an ordeal poison, and was not yet to be obtained in commerce.

Mr. Hart showed the effects of the solution prepared of different degrees of strength. They had been prepared as a watery solution of the alcoholic extract of the bean according to the direction of Dr. Argyll Robertson; that in which one minim corresponded to three grains of the bean answered best. It counteracted the effects of a solution of atropia, of three grains to the ounce, in dilating the pupil—inducing recontraction. Hence it is an agent of great utility in removing the inconvenience of dilatation of the pupil for the purpose of ophthalmoscopic study. Dr. Argyll Robertson had informed Mr. Hart in a letter of the tendency of the watery solution of the alcoholic extract to decompose. This decomposition occurred very rapidly, the solution becoming in a few days pink in colour (instead of being colourless) and fetid. It was arrested by the addition of glycerine to the watery solution. But Mr. Hills—of Messrs. Jacob Bell and Co.'s—had found that the alcoholic extract was soluble in glycerine, or apparently so; and Mr. Hart was now using the glycerine solution. This might prove to be a chemical solution, while the other certainly was not.

The action of the Calabar bean in counteracting mydriasis was well shown in several cases: one of partial mydriasis from paralysis of the third nerve, and one of the same condition from excessive debility after fever and accompanying asthenopia. Its power as an antagonist of atropine is equally shown by the firm contraction of the pupil and accommodative changes which it produces in the normal eye. Its action is, however, less durable than that of atropine; and if it be desired to make use of its curative effect in cases of mydriasis, Mr. Hart thought it would be found desirable to repeat the drop every four hours. Sometimes its instillation caused slight irritation; but there was often none at all.

In conclusion, Mr. Hart said he had been the first who had employed this agent in London, but that he should abstain from publishing at length his observations on the physiological and therapeutic effects of the drug, considering that Dr. Argyll Robertson, who had introduced this agent, and had with great care and sagacity at once investigated its powers and foretold its uses, should be allowed to gather the first fruits in this interesting field of observation which was his own by right of discovery. He would say that his observations completely confirmed those which that gentleman had already published.

UNIVERSITY COLLEGE HOSPITAL.

CASE OF SUDDEN DEATH OF A YOUNG WOMAN FROM FATTY DEGENERATION OF THE HEART.

(Under the care of Dr. HARLEY.)

SARAH M—, aged twenty-eight, an enormously stout, florid-complexioned, and puffy-looking woman, came to the hospital on the 14th of April, complaining of symptoms somewhat resembling hysteria. She described herself as being very nervous, and subject to fits of gasping for want of breath. She said that at times she felt as if her strength suddenly left her, on which occasions she was obliged to sit down for fear of falling. She frequently vomited after food. A mixture containing quassia and iron was prescribed. On the 21st she returned, saying she felt no better, and that in addition to her previous symptoms she had during the last week suffered from pain between her shoulders. The medicine was then changed to gentian and sulphuric acid; and after taking this for a few days, she said she felt much better and stronger. On the 28th, however, while attending to her usual household duties, she was suddenly seized with loss of power and a choking sensation. She partially rallied twice from the attack; but the third time she fell back in her chair, and instantly died.

At Dr. Harley's request Dr. Embling examined the body, and the following remarks are from his report:—

Autopsy, three days after death.—Decomposition far advanced; neck much discoloured; vessels evidently distended.

The brain normal in appearance, rather pale if anything; the lungs healthy; a few traces of old adhesions in the pleuræ; the heart much hypertrophied, pale, soft, and flabby, almost entirely enveloped in fat, and containing black fluid blood. When the muscular tissue of the heart was examined with the microscope, the transverse striæ of the fibres were found to have almost entirely disappeared, and their position to be occupied by minute oil globules. The oil globules existed abundantly, and were very regularly distributed in nearly all the muscular fibres. The abdominal organs were all healthy.

ST. GEORGE'S HOSPITAL.

MALIGNANT ULCERATION OF THE PHARYNX.

(Under the care of the late Mr. JOHNSON.)

SAMUEL H—, aged sixty, was admitted Sept. 3rd, 1862, in a state of extreme emaciation. He stated that for some time he had been unable to swallow solid food on account of "sore-throat." On examining the throat, there was found to be considerable ulceration of the soft palate and fauces, with constant discharge of an intensely offensive character. He was quite unable to take solid food, and it was only with the greatest difficulty that a spoonful of liquid could be swallowed. He was given all the nourishment he could take; but he died exhausted ten days after admission.

Autopsy twenty-three hours after death.—The body was extremely emaciated. The heart was blood-stained; the right ventricle contained a clot; the mitral and aortic valves were thickened, and their orifices contracted; the aorta was much affected with atheroma; there were old pleuritic adhesions. The pharynx and soft palate were considerably ulcerated, with deposit in the submucous areolar tissue, which appeared to be malignant to the eye, but under the microscope there was no evidence of cancer cells, the structure appearing to be irregularly fibrous. There was one ulcer just at the commencement of the oesophagus. The glands in the neighbourhood of the pharynx were simply enlarged. The liver was small, and affected with chronic cirrhosis; weight, two pounds ten ounces. Supra-renal capsules contained a quantity of calcareous or chalky matter, but were otherwise healthy; so also were the kidneys. The peritoneum contained much turbid serum.

GUY'S HOSPITAL.

IDIOPATHIC ANÆMIA; FATTY DEGENERATION OF THE HEART.

(Under the care of Dr. HABERSHON.)

ELIZABETH W—, aged forty, was admitted into Lydia ward on 6th October, 1862. She had been in the hospital several months before in the same ward. Her morbid symptoms were those due to anæmia. She was quite white and very feeble, so that she was obliged to keep her bed. She often had sickness, and thus a suspicion arose as to the existence of stomach disease, but none could be detected. She died on December 3rd.

Autopsy, twenty hours after death.—The legs were slightly anasarcaous, and the body remarkably white. The brain was pale and watery; increased serum on the surface and in the ventricles. Several ounces of clear serum were seen in the chest. The lungs were pale and watery. About two ounces of serum were found in the pericardium. The heart was rather small. Excess of fat seen on the front surface; this was observed to encroach on the muscular fibre, and to penetrate even to the interior, the walls being pale and soft. On the interior, especially in left ventricle, well-marked fatty degeneration was seen, with zig-zag striæ on the muscular columns and other parts. Tissue soft. All the other viscera were healthy. The kidneys and their capsules, supra and circumferential, were healthy.

THE ABORIGINES OF AUSTRALIA.—James Morrill, a native of Maldon, in Essex, who was wrecked in the *Peruvian* in 1846 on the eastern coast of Australia, and has just escaped from a seventeen years' captivity amongst the aborigines, testifies to the absence of epidemic diseases among the natives removed from intercourse with the whites.—*Sydney Morning Herald*.