nesser of the former must cool more readily, and enjoy greater alternations of heat and cold. The former part of this doctrine is founded on the experiments and deductions of Sir Everard Home, as detailed in his paper in the Philosophical Transactions. A number of experiments detailed in the paper on the vesicatory powers of differently coloured substances, under the concentrated rays of the sun, contradicted the deductions of Sir E. Home; and hence arose the necessity of looking to the vital properties of the skin of the negro, and the mode in which it is likely to be affected by the radiating and absorbing power of the pigment with which he is provided. Blumenbach and Winterbottom state, that the negro perspires more readily and freely than the European; and Davy says, "In the inhabitants of the tropics, the exhalant arteries of the skin seem unusually expanded, and the whole apparatus peculiar to this secretion unusually developed; and I believe that the blood itself is less viscid, more fluid, and flows more readily through the vessels, so as to promote perspiration, and by that means contributing to the cooling of the surface. And being cooled itself, it contributes again, when it flows back upon the heart, to the reduction of the temperature of the internal parts." Were the inhabitant of the tropics not possessed of this peculiar organization, his system could not respond to the stimulus of heat, by a determination of fluid towards the surface. Doubtless, the excessive absorption of heat by his skin, is useful in promoting this effect; but in the system qualified to respond to the stimulus of heat, and not in the organization of the skin alone, must an explanation be sought of the capability of the negro to withstand the heat of tropical regions.—*Athenæum*.

3. *Experimental Investigation of the Functions of the eight pair of Nerves.*—Dr. JOHN REID communicated to the Medical Section of the British Association the principal results which he has obtained in prosecuting this inquiry since the last meeting of the association. Dr. Reid chiefly confined himself to the Pulmonary and Gastric branches of the Nervus Vagus. In a great number of experiments upon dogs, in which the nervi vagi and recurrens were divided, Dr. Reid satisfied himself that, when a sufficient quantity of air reached the lungs, the respirations were at first performed with ease. The only immediate and constant effect observed, was a great diminution in the number of the respiratory movements, which at the same time became slower and more heaving. When the number of respirations were from sixteen to twenty in a minute, they, in general, instantly fell to from six to eight. In one animal, in which they were from twenty-four to twenty-eight, they immediately fell to fourteen; and, in another animal, in which they were twenty-four, they fell to eight. At a longer or shorter period after section of the nerves, the inspirations became more prolonged and heaving, while the expirations continued to be comparatively short and rapid, and attended by a sound caused by the sudden expulsion of the air. After a time, the inspirations become more heaving and prolonged, the blood is less perfectly arterialized in the lungs, and the arteries circulate blood gradually approaching the venous character; the animal becomes dull and stupid, and dies asphyxiated. In tracing the morbid changes upon the lungs, which follow section of the vagi, Dr. Reid has satisfied himself, from the dissection of animals killed at various periods after section of the vagi, that the bloodvessels of the lungs become gradually loaded with blood, and that this, in a few cases, is the only change which can be observed after death. More frequently, however, this congestion of the lungs gives rise to effusion of serum, which in the bronchial tubes, is rendered frothy by the passage of air through it; to condensation of portions of the lung without any apparent effusion of lymph; and more rarely to true pneumonia. The respiratory murmur is little, if at all, changed for a short time, but it afterwards becomes bronchial; and when the effusion of serum takes place, the true *râle crepitant* may be heard. It appears, however, that these morbid changes in the lungs do not necessarily take place after section of both vagi and recurrens. In a dog, which was killed in the presence of Dr. Alison, twelve days after section of the vagi and recurrens, and which had apparently recovered from the effects of the operation, the lungs were found perfectly healthy, though the cut ends of both vagi were rather more than one

inch distant from each other. Dr. Reid believes that the diminished frequency of the respiratory movements is sufficient to explain all the morbid changes in the lungs, which result from section of the nervi vagi. He related several experiments to prove that the respiratory movements are not arrested by division of the vagi and recurrens after removal of the cerebrum and cerebellum. In these experiments, it was, however, observed, that though section of the vagi did not arrest the respiratory muscular movements, they were much diminished in frequency. While these experiments illustrate the great importance of the par vagum as exciters of respiration, they also prove that there are other nerves which can transmit those impressions to the medulla oblongata, which excite the respiratory movements; and Dr. Reid adduced some facts to show, that one of the most important of these was the larger root of the fifth pair distributed upon the face. Dr. Reid then related four experiments, which he considered to afford satisfactory evidences of the continuance of digestion after section of both vagi, with loss of substance. In three of these, the lacteals and thoracic duct were found full of chyle. From these experiments, he concluded, that, though by division of the vagi a deleterious influence is propagated downward to the stomach, disturbing the functions of the digestive organs; yet, if the animal live long enough, this goes off, and digestion proceeds as before. He related some experiments to prove, that alcohol, opium, and prussic acid exert their deleterious effects as rapidly when injected into the stomach after division of the vagi, as when these nerves are left entire. He then detailed the result of five comparative experiments upon the effects of fatal doses of arsenic upon the watery and mucous secretions from the inner surface of the stomach and intestines. These results were completely at variance with the statement of Sir B. Brodie, that section of the vagi arrests the usual mucous and watery secretions of the stomach and intestines observed after poisoning with arsenic. He stated, that he had of late carefully watched the effects of division of the sympathetic nerve on the eye, and that he had satisfied himself that the contracted state of the pupil, and the partially-closed eyelids, which accompany the inflammation of the conjunctiva, consequent upon section of the sympathetic, take place previously to the inflammation of the conjunctiva, and are independent of it.

Prof. Owen requested to know, if albumen were discovered in the thoracic duct after the division of the vagi nerves. Dr. Reid replied, that no chemical analysis was made of the contents, but true digestion was inferred by him from the return of health and activity to the animal operated on.—Dr. Golding Bird pressed the same objection as Professor Owen, and wished that an experiment on the subject should be followed up by such a chemical investigation as would be decisive.—*Ibid.*

4. *On the Origin and Subsequent Developement of the Human Teeth.*—Dr. Gooden has observed dentition commence by the formation of what he denominates the primitive dental groove, on the floor of which the rudiments of the pulps of the milk teeth appear as globular or conical papillæ; septa afterwards pass from the outer to the inner side of the groove, between the papillæ, and thus each of the latter becomes situated in an open mouthed follicle, which is the primitive condition of the future sac. After the formation of the milk follicles, the lips of the groove still remain prominent; and when in this condition he denominates it the secondary groove. The rudiments of the ten anterior permanent teeth appear as little depressions in the secondary groove, internal to the mouths of the milk follicles. The papillæ of the milk teeth now begin to be moulded into the form of pulps, a change which is synchronous with the closure of the mouths of the follicles by two or more laminae, which agree in number, shape, and position with the cutting edges and tubercles of the future teeth. The lips and walls of the secondary groove now adhere, except in the situations of the ten depressions for the permanent teeth, and for a small extent posteriorly on each side, where a portion of the primitive dental groove remains in its original condition. In this portion the papillæ and follicle of the first large molar tooth appear, and, after it closes over, the lips of the secondary groove above it adhere, but not the walls; so that there is in this situation a