divided between ligatures; and that tumors and broken down tissues were left as irremovable; or that the growth was not disturbed, as it was most probably malignant, etc. With revised methods and improved skill such practice has been replaced by careful, thorough and bold enucleation. Now, by means of forcible pressure, hot water and gauze packing, hæmorrhage from severed adhesions is controlled without exposing the patient to the shock of prolonged operation under anæsthesia. With these improved methods the number of incomplete operations has diminished, and as individual skill improves and operative methods are perfected, the proportion will be still further lessened.

There is another influence which is destined to be very potent in improving results now impaired by delayed and incomplete operations. As knowledge is diffused and diagnostic skill improved, the family physician will recognize serious pelvic disease at an earlier period, and advise operation. With improved results, both the profession and the public will cease to regard peritoneal operations with the fear of the olden and mortal era, and thus delayed operations will be lessened. This should be the aim of every conscientious worker, and it is the purpose of this paper to show how essential are *promptness* and *thoroughness* in applying the advanced principles of modern pelvic surgery.

Discussion.

Dr. G. Betton Massey, Philadelphia, thought it unfortunate that these two papers were read together, as their conclusions are very different. The paper of Dr. Ashton was the result of careful work, as I well know. The paper of Dr. McMurtry seemed to be a collection of dogmatic statements based solely on his own opinion. He urged a complete operation, and said that we could not always say a woman died on account of the incompleteness of the operation. It is a little behind the times to advise the practitioner to send all his cases to the abdominal surgeon every time they have a pain in the belly. I venture the prediction that in the future serious interference with the abdominal cavity will be limited to those cases in which pus has been demonstrated to exist in the pelvis. He wished to call attention to the ease with which a powerful electric light could be enclosed in a series of test tubes, one inside of the other according to the amount of heat. This can be inserted into the vagina and will light up the pelvic cavity. This light will penetrate a cyst, but will not penetrate a collection of pus, which is a means of diagnosis of the presence of pus. This light will penetrate living bone even better. We can notice light shining through our fingers on holding them up before a bright light.

Dr. R. B. Hall, Cincinnati, thought it impossible to determine the presence of pus in all cases, and that the prediction of Dr. Massey would not hold, and that this would be dangerous doctrine to promulgate. The doctor reported a case in point.

Dr. Joseph Tabor Johnson, Washington, D. C., thought the papers read, and others which have been recently published, teach us that we are not thorough enough in our operating.

Dr. Franklin H. Martin, Chicago, thought the paper of Dr. Ashton taught us how to avoid trouble. We should give much care to the treatment of raw surfaces in the intestine. These surfaces will readily take on adhesions. These should be covered with peritoneum if possible, otherwise covered with an omental graft. Stumps should be covered with peritoneum, which will also prevent attachments. Saline treatment is very important. In the Woman's Hospital at Chicago we endeavor to get a passage of feces or gases during the first twelve hours.

Dr. Edwin Ricketts, Cincinnati, has had much benefit from the administration of strychnia in large doses in intestinal paresis. A case which he recently had looked very much like peritonitis, but after the administration of strychnia a large quantity of gas was discharged. The speaker could not agree with Dr. Martin as to the danger of adhesions resulting from the stump. Such had not been his experience.

Dr. W. E. B. Davis, Birmingham, Ala., did not think abra-

sions had such an influence on the causation of adhesions as claimed. Purgation twenty-four or thirty-six hours after operation will not break up adhesions. It should be done earlier. He considered it very important that the bowels be returned within the abdomen during the operation. In fibroids it is the adhesions which give symptoms, and it is unjust to blame these on the electrician.

Dr. E. P. Davis, Philadelphia, said the pathology of the peritoneum was a very important feature. Different portions of peritoneum have different powers of resistance. Iodoform seems to be least irritating to the peritoneum, next boracic acid. It is known that the peritoneum of dogs is different from that of the human being.

Dr. W. E. Ashton, Philadelphia: Obstructions are not always due to faulty technique; sometimes to inflammation which could not be prevented. Taking a graft from the omentum only made a point where the gut might pass through. Strychnia has been followed with very good results. I give one-fifteenth to one-tenth grain three or four days before and after section. It increases peristaltic action of bowels, clears them of feces and lessens the danger of intestinal paresis.

Dr. L. S. McMurtry, Louisville, in reply to criticism, said that the observations and deductions made in his paper were based upon his personal experience in pelvic surgery, and were fortified by illustrative cases. He deprecated resort to operation in doubtful cases, and advocated most careful discrimination in deciding the question of operation in all cases; but he believed it equally an error to delay operation in clearly defined conditions requiring operation until the patient's chances of recovery are reduced or lost. Of course, there must be a restriction on early operation in fibroid tumors in accordance with the rapidity of growth and amount of hæmorrhage. When these conditions clearly point to the necessity for operative interference, the sooner it is done the better. When resorted to early, the appendages may be removed and the growth, as a rule, will be arrested and hæmorrhage cease. If hysterectomy is rendered necessary, the sooner it is done the safer it is for the patient. It has been demonstrated that electricity is a failure in the treatment of uterine myomata; and while these tumors sometimes disappear at the menopause, or sooner, without treatment, they often grow in spite of the menopause and treatment, and unless removed destroy the patient's life.

COLPO-PERINEORRHAPHY.

Read in the Section of Obstetrics and Diseases of Women, at the forty-third Annual Meeting of the American Medical Association, held at Detroit, June, 1892.

BY EDWARD W. JENKS, M.D.,
OF DETROIT.

INTRODUCTION.

In 1877, I read a paper before the Michigan State Medical Society in which were mentioned "some new procedures in the operation for laceration of the perineum" and Jan. 8, 1879, I read a paper before the Cincinnati Obstetrical Society, which was published in the *American Journal of Obstetrics*, for April, 1879, describing the same procedure as in the first mentioned article, the latter being entitled "Perineorrhaphy and a Description of a New Mode of Operating." This paper advocated a new mode of submucous dissection of the tissues or what is now designated flap-splitting.

This mode of operating was favorably commented upon by the leading medical journals of the day and the leading text books on gynecology published in this country continued for many years to sanction it. In that paper the cutting away of the flap instead of preserving it was advised. Subsequently I preserved the flap, and believing for a time my mode of operating as well as the preservation of the flap was original, so stated when asked regarding it by some of my friends, notably the president of this Association (Dr. Marcy), who mentioned the matter in one of his publications. But learning that others had also described saving the flap no claim has been made as to

its originality with myself. The submucous dissection I described, or flap-splitting operation as now designated, was made several hundreds of times by me in public clinics, as well as by many others who had adopted it, some years before the flap-splitting operation of Tait or others had become known.

However, I am not strenuous in claiming priority in the mode of operating I described so long ago, as it is of little importance, for I am only one of the great multitude working in the field of gynecological surgery.

I am prompted to this historical fact as an introduction to this paper as the surgical procedures I made known fifteen years ago are occasionally referred to at the present time; and a late edition of a well known text book that had previously commended it by means of lengthy quotations and illustrations now dismisses it by a brief allusion as "too bloody."

Although repeatedly asked to write an additional paper I have had other fields to cultivate, and what seemed more important in surgical matters to engross my time and attention, but of late there seems to have been a revival of interest in plastic gynecological surgery.

DESCRIPTION.

The operation I make to-day and which is described in this brief paper is, I believe, a great improvement on my former method. It would hardly be possible for one to make hundreds of operations of any kind without finding a chance for some improvement either in the mechanical portion or in results obtained.

Of the surgical devices before the medical public known under the general title of perineorrhaphy the name is legion, consequently the merits or demerits of other operations than the one here described will not be considered.

It is not my purpose to discuss laceration of the perineum or the necessity for repair in detail. It is well known to every careful observer that rather extensive laceration of the perineum and vaginal walls occur in childbirth from which no bad results follow. Again it is equally well known that many discomforts and divers reflex symptoms often ensue from lacerations that at the time of their occurrence seem too insignificant to demand a passing thought or even a single suture.

It is to the consideration of secondary operations alone that your attention is called. I will state as a general proposition that operations are not demanded because of laceration *per se*, but when there are unmistakable discomforts that can be plainly traced to them, and health and comfort can only be recovered by restoring the torn parts to their normal relations.

The portion of the recto-vaginal septum known as the perineum supports the lower portion of the posterior vaginal wall, which in turn supports a corresponding part of the anterior vaginal wall. The lower portion of the rectum is sustained and the proper performance of its functions aided by the perineum. Four muscles, the levator ani, sphincter ani, transversus perinei and bulbo-cavernosus are here united; and it is the severance of these from their fellows on the opposite side together with the separation of the perineal fascia which produces the mischief, therefore laceration of the perineum and a portion of the

posterior vaginal wall, whether partial or complete, may cause a variety of conditions, such as loss of vulvar integrity and impairment of the functions of the rectum, partial or complete incontinence of the rectum and bladder, increased and irritating secretions of the vagina and rectum and recurring prolapse of the rectum after operation for prolapsus recti, descent of the recto-vaginal septum or rectocele, a similar condition of the anterior vaginal wall and bladder or cystocele, and descent of the rectum. There are also various neurotic and sympathetic disorders which it is needless at this time to dwell upon.

I have seen quite a number of patients that had been operated upon with the result of having, to all external appearance, a perfect perineum but with a rectocele above the line of the dissection or flap-splitting that formed an excellent pocket for the accumulation of uterine and vaginal secretions, in which the neck of the prolapsed uterus was constantly macerating. In many of the cases where only a portion of the redundancy is remedied the only benefit seems to be, to provide a better support for some form of pessary to sustain the uterus and vaginal walls.

My own observation and experience, which I presume agrees with others, is that there are, at least in a general sense, four important ends to be attained in repairing the class of injuries under consideration: 1. To restore the loss of power and function to the lower portion of the rectum and vagina. 2. To restore the normal sustaining quality of the posterior vaginal wall for the anterior vaginal wall and bladder. 3. To provide as much support for the uterus as the perineum naturally gives. 4. To cure the many distressing nervous accompaniments.

Any surgical procedure which does not obtain such results to a great degree is not in a strict sense successful.

A perineum may be operated upon and, as far as external appearances are concerned, is successful, but if above the point of dissection there still remains a redundancy of the vaginal walls, or the restoration is not sufficient to support the anterior vaginal wall the operation is but partially successful. This is true whether the uterus is held up to the health line or not. There are so many causes operating to produce uterine displacement, that it is not as a general rule just to gauge the success of an operation by the measure of uterine support it secures. For the reason that these partial operations are often insufficient and are not followed by the anticipated beneficial results, I have chosen to designate the surgical procedure I have been making for a number of years, colpo-perineorrhaphy. To accomplish the best permanent results it is essential that dissection of the flap extend as high within the recto-vaginal septum as there are signs of slack or redundancy of the posterior vaginal wall.

My mode of procedure is as follows: I first nick with the scissors each labium to mark either termination of the anterior margin of the flap and then, having introduced two fingers into the rectum and assistants making the parts taut, I insert the sharp pointed scissors near the juncture of the integument and mucous membrane in the median line or sometimes on one of the nicked lips and proceed to dissect a flap up the septum as far as redundancy of the walls can be observed. (Fig. 3.) It is important for the sake of making a more rapid and neat operation that the dissection be made in its entirety without withdraw-

ing the scissors. When I first described my mode of operating in 1877, my friend, the late Dr. Albert H. Smith, devised a knife to take the place of scissors and subsequently I caused a knife to be made larger and more flexible than his. (See Fig. 1.)

The scissors which I have used for the past five years are slightly curved with blades completely overlapping each other, and both inner and outer edges ground equally sharp. I have found this form

ent directions; still I do not claim any originality in this matter and wish the credit to be given to whom it is due.

No better results have ever been obtained than by silver sutures, but on account of their stiffness they cause more pain, therefore, in extremely sensitive patients I substitute the silk-worm gut, which possesses the principal advantages of silver wire and is not so unyielding. Kangaroo tendon is also a safe

Figure 3.

of blade is less liable to become entangled in submucous dissections than the ordinary shaped blade. (See Fig. 2.) These scissors can be used alternately in their proper capacity or with blades closed as a knife to dissect the flap more perfectly, as their sharpened outer edges and points render this an easy task, and there is a great advantage in their use in cases where there is much bleeding, as there is less hemorrhage than with an ordinary knife.

Figure 4.

and useful suture, as its time of continuity is from fifteen to twenty days.

The needles are a straight, flat non-cutting needle about two inches in length and a slightly curved Peaslee needle.

The latter is provided with a carrying thread which is number 4 or 5 braided silk about eighteen inches long. The former is used only for the short and superficial sutures which are threaded directly into the needle.

One objection recently made to this operation is that it is bloody and sometimes one or more arteries have to be tied.

To my mind, if that is the case, the surgeon of the present day is not intimidated by a few small spouting vessels and if such a calamity occurs it is quickly and easily remedied.

In many hundreds of operations I have not been compelled to ligate vessels, all told, more than six times.

The next important step in the operation after the dissection of the flap is the insertion and adjustment of the sutures. I was led many years ago to adopt the method of the late Dr. Agnew of Philadelphia, in placing sutures in the perineum. Since then I have carried them deeper, higher and in somewhat differ-

In the majority of methods of operating for incomplete laceration, the first and frequently the second sutures are shorter and of far less importance than the third or fourth, or fourth and fifth, as the case may be. But in my operation matters are reversed, for the first two sutures are the longest and most important. Indeed, for want of a better term I often call them *the parent stitches*. (Fig. 6.)

The first assistant lifts up the flap by means of a tenaculum hooked into the edge at the center. (Fig. 4.) Introducing two fingers of the left hand into the rectum to guard against wounding it, I start the needle in at the distance of one-third to one-half inch back from the denuded surface, and turning the point well toward the left buttock and the handle correspondingly as far in the direction of right buttock, I

push it rather deeply into the tissue of the anterior ischio-rectal space, then upwards and finally inwards along recto-vaginal wall until it has been carried just above the highest point of dissection in the center, at which location or as near to it as possible the needle-point is brought out. When the point is about to come through, if counter-pressure is made with a blunt hook, the needle can be pushed through more easily, with less strain upon the septum and with less likelihood of pricking the anterior wall by the

Figure 5.

sudden emergence of the needle than without its use. As soon as the eye appears a loop of the carrying thread is pulled out by a tenaculum and the suture passed through it. When the needle is withdrawn one half of the first stitch will be *in situ*.

The needle is then introduced in the same manner in the opposite side, the upper end of the suture threaded into the loop and the other half of the stitch carried to place.

For the second stitch the needle is started in about a third of an inch above the first and its point directed at first outward in the same manner as in the introduction of the first suture. Not quite so much lateral tissue is taken up this time, that is, the needle does not make quite so wide a side sweep for the second suture but passes more directly up along the recto-vaginal septum and when it has reached the upper third of its course, crosses the first suture and comes out on the vaginal mucous membrane about one-half or two-thirds of an inch above the central highest point of dissection. After drawing the first half of the suture into place the needle is introduced in the same manner on the opposite side for the second half of this stitch.

For the third stitch, a third of an inch above the second, the needle passes along the denuded surface till it reaches the line of junction of the septum and the flap, when it enters the latter at about its upper fourth, burrows across to the opposite side and down the denuded surface to the outside.

This stitch can sometimes be introduced in one continuous circuit, in other cases one-half at a time.

The fourth and fifth stitches are buried under the denuded surface as far up as the junction of the septum and flap where they pass under the flap without burrowing in it to the opposite side. With each of these sutures it is usually more convenient, although not necessary, to put in one-half and then introduce the needle on the opposite side in the same manner for the other half.

The ends of the sutures are now placed together and traction enough made upon them to determine whether the denuded parts are coming properly into apposition. At this time the flap will emerge more or less from the introitus and will frequently have the appearance of being much too long; the operator will consequently be tempted to pare off a quarter of an inch or more from the anterior edge, but this

should not be done except in rare cases of great redundancy. This slack is disposed of by the gradual retraction of the flap during the process of healing and settling into normal relations. The sutures are now loosened again and the sixth stitch introduced which is designed to purse up the anterior side of the flap and also bring together the last of the denuded surface. For this purpose the straight thin needle previously mentioned is used. The needle is passed under the portion of denuded surface contig-

Figure 6.

uous to the edge of the flap and thence into and through the latter to the opposite side. All the sutures are now picked up and slackened enough to allow the flap to be raised in order to clear out by the douche any clots that may have collected.

The sutures are then all drawn up ready for fastening.

If silver wire has been used each suture is carefully shouldered and twisted. If silkworm gut has been employed the sutures may be either tied or secured by perforated shot.

Care is required in adjusting the first two long sutures not to draw them too tightly, otherwise they will cut in a little, and owing to their including so much tissue adjacent to the rectum, will be painful. The third, fourth and fifth sutures can be drawn more tightly. The sixth, again, should be but moderately tight, as the pursed up edge of the flap will not bear too much constriction.

Usually two or three superficial sutures of fine silk, horse hair or small silkworm gut will be required to coapt any raw edges turned out by the puckering up of the flap.

In case there seems to be any liability of hæmorrhage beneath the flap I place a strong silk suture by means of a Peaslee needle outside of the adjusted sutures and over the flap, which I retain for about twenty-four hours and then remove. (Fig. 5, a.)

These last are not absolutely essential but they give a neat appearance to the operation as well as dispose of surfaces for absorption or granulation. If silver wire has been used the ends may be massed together and inserted into half an inch of small rubber tubing to prevent them from pricking the patient.

In any surgical work, but especially in the plastic operations of gynecology, it is extremely convenient and helpful to an operator if he is ambidextrous. However, but few possess this accomplishment to a full degree and it is not really a *sine qui non* to skillful operating. Many expert operators do not interchange much the work of the two hands in the use of the knife, scissors and needle.

In this operation or almost any other for perineorrhaphy, the long stitch or stitches which pass from

the outside up to the highest point of dissection in the vagina should always be put in one half at a time. If the dissection is made even approximately as high as it should be, a slightly curved needle cannot make the whole circuit at once, except in a patient with lax tissues and a broad space between the tubera ischiorum.

But even when it can be done it is accomplished at the expense of considerable strain upon the parts operated upon and the whole vaginal column.

I have constantly mentioned six as the number of sutures employed, but only because that is the number most commonly required. Occasionally five are sufficient, and sometimes seven or eight are needed.

As there are no exposed raw surfaces either externally or internally, but little dressing of any kind, and but few vaginal douches are demanded. Three or four are the usual number; one at the end of thirty-six hours, another on the fourth or fifth day, and another on the morning of the day the stitches are removed, usually the seventh. The external parts on the other hand, require about the same attention as in other perineal operations. Night and morning, and each time after urinating the soft parts adjacent to the line of union, and also the buttocks are carefully separated and the wound and the surrounding parts gently irrigated with sterilized water or a one to six thousand solution of bichloride.

The surgical procedure which I have here described under the name of colpo-perineorrhaphy, cannot commend itself on account of the consummate ease or rapidity in which it can be made. It is not as easily or quickly done as ordinary perineorrhaphy, nor even as the flap-splitting operation of which so much has been written of late.

But after essaying different operations from Baker Brown's to many of the present day, I have settled upon the method I have here briefly outlined as the best one I can make for the great majority of cases that present themselves to me for treatment.

In conclusion there are a few points to which I wish to direct attention.

1. Any single mode of operating is not adapted to every case of laceration of the perineum.

2. All other qualifications being equal that surgeon will be the most successful in this class of operations, who, instead of following hard and fast rules, possesses a mechanical skill which he can adapt to the peculiarities of each individual case.

3. The subsequent comfort of patients is not facilitated by superabundance of cicatricial tissue within the vagina; therefore the anterior wall, instead of being subjected to any surgical procedure for redundancy should be sustained by a restoration of the normal posterior wall.

4. The surgical operation here advocated has for its object, a restoration of the torn posterior vaginal wall and perineum to its normal condition, whereby there is afforded (a) support for the uterus to the full extent provided for in the vaginal walls; (b) support for the anterior vaginal wall and bladder; (c) support for the lower end of the rectum.

84 Lafayette Avenue.

THE American Microscopical Association will meet in Rochester, N. Y., August 9, and not August 10, as heretofore announced.

OVARİOTOMY IN THE PRESENCE OF PREGNANCY.

Read before the Section of Gynecology and Obstetrics, at the Forty-third Annual Meeting of the American Medical Association, at Detroit, Mich., June, 1892.

BY WILLIAM H. MYERS, M.D.,
OF FORT WAYNE, IND.

Seventy-six (76) years have elapsed since Recamier presented to the medical world the uterine speculum, and the dark age of uterine pathology was passed and a new era begun.

As in history, we progress by alternation, by conflict and by revolution, so were these the steps in medicine and surgery by which we have attained the advanced thought of to-day.

From 1816 to the present, a wide range of material has been submitted; brilliant at the time, but existing now as the residue, as merely a trace of a previous pathology.

Recamier taught that the primary element of uterine disease was inflammation and ulceration; Bennet, ulceration of the cervix; Tyler Smith, hyper-secretion of the mucus glands of the cervix; Meigs, retro-version; Velpeau, flexion; Lisfranc, congestion and engorgement of the cervix; Madame Bovine, uterine catarrh; Grailly Hewitt believes that all the ills of woman are due to uterine displacement; Tilt, ovarian peritonitis. I might in this connection refer to the labors of Goupil and Bernitz in pelvis peritonitis; for they were the first to teach us a correct pathology of this condition, and the secondary importance of cellulitis. We of the present accept with the greatest facility the doctrine that the peritoneum and the ovaries are the structures underlying nearly all of the phenomena presented in uterine disease, and we relegate cellulitis. The truth of this doctrine opened the field of intra-peritoneal operations, and ovariectomy, according to a recent writer, passed through three successive periods, a period of groping, ending with the successful work of W. L. Atlee, Baker Brown and Sir Spencer Wells; second, a short period of excessive specialization; and third, a period of popular adoption deriving its impulse from the introduction of antiseptics.

The treatment of ovarian cysts has been settled, but the presence of an ovarian cyst during pregnancy presents questions for our consideration. The doctrine advanced may be thus enumerated:

1. Non-interference.
2. Puncture of the cyst.
3. Induction of premature labor.
4. Porro's operation.
5. Ovariectomy.

In regard to the question of non-interference, in the light of modern surgery, we dare not entertain it, for the dangers of axial rotation of the tumor, rupture of the cyst, of impeded labor, rupture of the vagina during labor, far outweigh the dangers of the removal of an ovarian tumor. The exceptional cases reported, where pregnancy has coexisted with an ovarian tumor, and the woman has passed safely through confinement, are too few to be accepted as a precedent; for in a large proportion of cases, if the tumor be large, abortion will occur; if not, a lingering labor and still-born child will be the result, and the danger of sudden death be ever impending. In a paper by Doctor Playfair, published in the ninth volume of the Obstetrical Transactions, he reports