was folded over the tumor, giving it the complete peritoneal covering. At no point was the tumor fused with the structures to which it was attached by peritoneal adhesions. The transverse colon was attached along the entire anterior inferior border of the tumor, but the gut walls were nowhere invaded by the tumor growth. When the tumor was cut open it was found to contain in its interior a cavity 13 by 11 by 10 cm. The cavity was subdivided into two compartments by a septum formed by the tumor tissue. The roof of the lower compartment and the internal surface toward the left side were perfectly smooth. Most of the right internal surface and most of the floor were uneven and covered with a ragged grumous mass. Here the tumor tissue had broken into the smooth internal lining of the cyst. The upper compartment of the tumor cavity had evidently not originated as a true cyst, like the lower compartment. It was an irregular pyramidal cavity with a very ragged and irregular internal surface. The cavity was partly filled with broken-down tumor tissue. The tumor walls surrounding the central cavity in varied thickness from 5 to 10 cm. The tumor tissue on section appeared divided by connective tissue trabeculae into roundish and oval nodules varying in size from a fraction to several centimeters in diameter. Near the outer surface the tumor tissue was grayish-white and quite solid; more toward the interior it assumed a light dirty brown, was spongy and honeycombed. The transverse colon, as stated before, was firmly attached to the anterior lower aspect of the tumor. When cut open the gut showed an intact mucous membrane, the tumor not having involved the walls of the gut.

The microscopic examination showed that when the tumor had not undergone any retrograde or degenerative changes it consisted of large spindle cells. These cells took their origin from the adventitia of vessels. The cell nuclei were generally fusiform, some decidedly rod-shaped, others oval and even round. They generally contained one or two nucleoli and a moderate amount of finely granular chromatin. The cell protoplasm was quite abundant and stained deeply with eosin.

The tumor cells formed bundles and masses which interlaced each other freely in every direction. Thin-walled blood-vessels were abundant. At some distance from the surface the tissue exhibited a tendency to retrograde changes. Here the cells showed coagulation necrosis, and the nucleus became indistinct, likewise the cell body. Deeper still the nuclei disappeared and a rarefied tissue was formed which looked like myxoid tissue. In such necrotic places extravasated blood was found free between the degenerating tumor tissue.

The grumous material found in the interior of the tumor consisted of necrotic cells mixed with free decomposing blood, fibrin and hematoidin granules. In some places the necrotic tissues showed numerous polynuclear leucocytes, some exhibiting nuclear fragmentation. One of the pieces of tumor examined showed where the tumor had been attached to the wall of the stomach. The viscous wall was thin and atrophic.

The capsule of the tumor consisted of coarse connective tissue fibers with slender, fusiform, deeply-staining nuclei. No endothelial cells were seen on the capsule.

The kidney showed marked congestion of all the vessels. The epithelium lining the uriniferous tubules showed cloudy swelling. Many of the tubules were filled with a granular material and some contained hyaline casts. The interlobular connective tissue of the liver was moderately increased. The peripheral cells of the lobule were fairly normal, but in the central zone they were densely filled with bile-coloring matter. Both in the peripheral and central zone a number of liver cells showed cloudy swelling. The spleen pulp was densely crowded with erythrocytes. In such places hemosiderin was found in large amounts. In other places the pulp cells showed a tendency to become fusiform. The myocardium was normal.

From the microscopic examination it was evident that we were dealing with a malignant connective tissue tumor—a sarcoma. The neoplasm, as appears from the history, grew rapidly, and speedily led to a profound cachexia. While the tumor formed adhesions to several of the neighboring organs it had not broken through any of them, nor had it given rise to metastases. The

absorption of the ptomains had produced a chronic splenitis and changes in the liver, partly parenchymatous, and partly indicative of early interstitial cirrhotic changes.

I here express my gratitude to Dr. Herzog for his assistance in preparing this report.

BIBLIOGRAPHY.

1. Deutsche Med. Woch., Oct. 22, 1889.
2. Senn's Surgery of Pancreas, 1886.
3. Mayo's Outlines of Human Pathology.
4. Contribution à anatom. path. de pancreas; Gaz. Méd. de Paris, 1874, p. 624.
5. Proc. of N. Y. Anatomical and Surgical Society, Vol. ii, p. 495, 1880.
6. Progreso Med. e Farm., Madrid, Vol. i, p. 77.
7. A. Blind: Sarcoma of Tail of Pancreas, Bull. de la Soc. anatomique de Paris, Séance du 21 Février, 1894.
8. Jour. Am. Med. Assn., Vol. xxvii, p. 1240.
9. Ziegler: Beiträge path. Anat., Bd. 22, S. 105-131.
10. Trans. Path. Soc., London, 1898, p. 178.
11. Neve: The Lancet, p. 659, 1891.
12. Schueler: Fall von Sarcoma pancreas haemorrhag.; Dissertation, 1894.
13. Sarcoma Encephaloide de cabica de pancreas; Correio Med. de Lisboa, Vol. xii, p. 61, 1883.
14. Aldor: Beiträge zur Casuistik der Pancreas Geschwulst. Gyogyasat, 1895.
15. Ueber ein Sarkom des Pankreas, inaugural dissertation, Gies-sen, 1897.
16. Ehrmann: Transact. Chic. Path. Soc., Vol. ii, 1897.
17. E. Italia: Policlinic, Roma, 1900, Vol. vi, p. 239.
18. Primarius Sarcom in Kopf des Pancreas; Allg. Med. Centr. Ztg., Berlin, 1868, Vol. 37, p. 781.
19. Briggs: St. Louis Med. and Surg. Jour., Vol. lviii, p. 154, 1890.
20. E. T. Bruen: Boston Med. and Surg. Jour., Vol. cviii, p. 110, 1883.
21. Virchow: Hirsch Archiv, Vol. ii, p. 211.
22. Pepper: Proc. Phila. Path. Soc., Vol. iii, p. 182, 1871.
23. Dieterich: Archiv für Klin. Med., Vol. xxxv, No. 2.

A CASE OF EPITHELIOMA DEVELOPED ON THE BASIS OF A HEALED LUPUS VULGARIS TREATED BY X-RAYS.*

DAVID LIEBERTHAL, M.D.

CHICAGO.

The patient, a tradesman, a native of Poland, is 53 years old, and does not remember ever having been afflicted with any disease, with the exception of that herein described. His parents, who are dead, enjoyed good health, and neither they nor any other of his relatives were subject to skin affections. He has eight children, and two grandchildren, all enjoying good health. At the age of about 2 years there developed behind his left ear a growth which soon exulcerated, and in the course of a few years his face, especially the nose and mouth, became affected. The skin thus diseased repeatedly broke down and rehealed. The mouth gradually became so closed that he could only insert his little finger.

At this stage the nose, which meantime had repeatedly broken and healed, began to break down until its tip was destroyed at about the age of 12 years. He was then brought to the University Clinic of Koenigsberg. There the ulcerating surfaces were regularly treated with the thermocautery under chloroform anesthesia. The ulcers healed, but the mouth grew smaller by cicatrization, and the nose was closed up. During his stay of a year in Koenigsberg he took cod-liver oil, a tablespoonful three times daily. He was then sent to the clinic in Berlin, where he underwent an operation consisting of wide incisions at the angles of the mouth and transplantation of mucous membrane from the cheeks upon the wound surfaces. The residue of the nose, which had healed up during the treatment in Koenigsberg, had no opening, and inasmuch as the patient did not consent to the formation of a new nose, it was decided to render it possible for him to breathe through the organ. To this end there were made two

* Presented to the Chicago Medical Society, Jan. 2, 1901.

symmetrical openings into which were placed silver tubes fastened with bandages. But the septum soon sloughed, and a single opening resulted, which remained permeable and permitted him to breathe freely. He stayed in Berlin about two years, and during that time continued the use of cod-liver oil, a tablespoonful three times a day. He considered himself cured, and soon after immigrated to Chicago, some thirty-eight years ago. Until about a year ago his condition gave him no concern. He then felt a painful nodule behind his left ear. About eight months ago a wart the size of a bean developed on the left side of the lower jaw, and slowly enlarged. It was itchy and soon showed openings out of which exuded whitish, worm-like masses on pressure. The tumor continued to increase in size, and during the last eight weeks developed more rapidly to its present condition. Pain is felt at times, and then on the lower border of the tumor only. He is of strong build, and of good health. On the neck just behind the left ear a scar is visible at the seat of the first affection of the skin. On the left side of the forehead is another scar. On the cheeks and chin, especially around the

grayish-white, ill-smelling exudate is visible, which increases in quantity by pressure. At times there is pain in the lower and posterior part. At the level of the posterior angle of the left lower jaw is found a bean-sized, round, hard, retramaxillary gland. I have to add that the respiration through the nose, as well as the sense of smell, is unimpaired.

Judging from the history, and more especially from the sequelæ of the skin disease which he acquired in early childhood, there remains but little doubt that it was a case of lupus vulgaris. The tumor on his face is clinically an epithelioma, which opinion is confirmed by the microscopic examination.

In regard to the prognosis, it is a well-established fact that epithelioma developed on a scar of lupus vulgaris gives but little prospect.

Drs. Anthony and Harris, who saw the case with me, suggested the application of the x-ray, and Dr. Haiselden of the x-ray department of the Chicago Polyclinic commenced the exposures.

Since the presentation of this case, the treatment has been continued. Beginning December 26, the exposures

nose and mouth, there are extensive scars. The tip of the nose is absent, and on its site there is cicatricial tissue, with a hole in the center. The red border of the lips is uneven, owing to cicatricial contraction, and for the same reason we find deep fissures at the angles of the mouth and on the mucosa of the cheeks. Over the left zygoma we see infiltrated skin, with a red, glossy, somewhat scaly surface. The patient states that he noticed it since he wore a bandage for the fixation of the silver tubes. On the left side of the face and neck we see a bluish-red, lobulated, cauliflower-like tumor, raised about half an inch above the level of the surrounding surface. Around the upper margins of its base scars are visible. It has the shape of a triangle, with its apex in the middle of the cheek, and its base about $1\frac{1}{2}$ inches beneath and parallel to the margin of the lower jaw. It is movable with the skin, has a soft consistence, and overlaps its base. At the junction of the tumor with the seemingly healthy surrounding tissue, deep fissures are to be seen. On its surface, a

were made twice a week. After the third exposure January 7, the skin in the vicinity of the tumor reddened, whereupon they were discontinued for one week, and after the disappearance of the irritation were repeated twice a week during January and February, during the first three weeks of March every other day and during the last week of March and first of April, every day except Sundays. Since then no exposures have been made. The primary current used for the induction of the secondary was between 8 and 12 volts and $1\frac{1}{2}$ amperes; interruptions averaged 800 per minute, the distance from the exposed surface varied from 6 to 3 inches, and the time of exposure from ten to twenty-five minutes.

After the third exposure there was noticeable diminution of the secretion and at times the tumor appeared perfectly dry. It had also flattened to a slight degree, but had spread downward and toward the chin about an inch. Seeing that the growth had not been checked, the use of the x-ray will be discontinued. I believe

that in incipient cases good results may be obtained by the x-ray. I would refer here to cases thus treated and cured by Drs. W. Johnson and W. H. Merrill.¹

S. Steenbeck reported at the International Medical Congress, Paris, August 1900, two cases of cancer of the nose, one of which had developed upon lupus vulgaris, both cured by the x-ray. All these cases were of very small size.

In conclusion, I take great pleasure in expressing my gratitude to Dr. H. J. Haiselden for his painstaking application of the x-ray.

103 State Street.

URETERAL IMPLANTATION INTO THE BOWEL FOR DIVERSION OF THE URINE.*

AN EXPERIMENTAL RESEARCH.

JACOB FRANK, M.D.

Surgeon to the German Hospital; Consulting Surgeon to St. Elizabeth's Hospital and Jewish Orphans' Home; Member of the American Medical Association; Fellow of the Chicago Academy of Medicine, Etc.

CHICAGO.

I fear that I have chosen for my paper a subject which may seem old and of comparatively little interest to some, and have consulted therein rather my own tastes and inclinations. The diversity of surgical conditions, brought about in the course of some pelvic or abdominal operation, to the ureters is sufficiently large and the number of cases met with great enough to make the subject of interest to every surgeon.

Again, the numerous pathologic conditions to which the urinary bladder is heir, which must sooner or later terminate fatally unless some means of relief can be afforded, have been sufficient incentive for experimenting in search of some surgical means that would furnish relief to those unfortunate sufferers.

The establishment of an artificial communication between the ureters and the intestines by means of surgical procedure should not be one of choice, but of necessity. It is applicable only in those cases where other means for the preservation of the kidneys are impossible or contraindicated. In cases of uretero-vaginal fistula the operations of Schede, Simon, Landau, Bandl and Pozzi should first be tried. If these attempts fail, intestinal implantation is much preferable to nephrectomy, which should only be done as a last resort. In carcinoma of the bladder, where the outlets of the ureters are encroached upon and entire resection of the bladder is indicated, ureteral implantation is all that can be done. The same holds true in severe cases of tuberculosis, but the renal lesions which are so apt to exist at the same time render permanent benefit unlikely. Unilateral anastomosis is indicated in those cases where the ureter is wounded higher up, as may happen during the course of an intra-abdominal operation, or where trauma or morbid processes have brought about the same result. Loss of substance of the ureter, if the ureteral wall is in such a condition that it can not be united, as after the removal of a calculus, may be treated by intestinal implantation. In exstrophy of the bladder I believe that vesico-rectal anastomosis will give the best results for urinary diversion. Much has already been done in this field of surgery, and I will give a little semi-history of ureteral implantation into the intestine. The union

of the ureters with the intestinal tract may be accomplished by means of a mechanical device or suture, in three ways, viz.: 1. The severed ureters may be anastomosed in the region of the rectum, with some portion of the ascending or descending colon, or even into the small intestine. 2. An anastomosis may be accomplished by implanting the vesical trigone in the intestine. 3. A permanent fistula may be established between the bladder and the rectum.

The first efforts made to direct the course of the urine into the intestine were in cases of exstrophy of the bladder. Simon first attempted this after the suggestion by Roux in 1851, upon a human patient, by passing a loop of thread through the walls of both the ureter and rectum, and tying them tightly together.

Figure 1.

Necrosis occurred at the point of ligation, and a fistula resulted. The patient died of pyelonephritis. Thomas Smith, in 1879, sutured successively the two ureters into the colon, but death quickly followed the second operation. The autopsy showed that on the left side, the first one subjected to operation, the point of exit of the ureter was obliterated, causing hydronephrosis. On the right side pyelonephritis and inflammation of the ureter were found. These fruitless attempts found few imitators, and the question seems to have been held in abeyance for a few years. In 1881 Glücke and Zoller published their unsuccessful attempts at implantation of the ureters into the rectum, and empha-

1. "The X-ray in the Treatment of Carcinoma." Philadelphia Medical Journal, Dec. 8 and 15, 1901.

* Read before the Chicago Academy of Medicine, Jan. 11, 1901.