pended and the attacks recurred. It was renewed, and the attacks became much less severe, and diminished to two in a month. It does not at all follow that there is much local pelvic disease to produce this form of epilepsy. In some cases the attacks occur only at certain menstrual periods at first, and after a few years they come on more frequently and with more severity, and between the menstrual periods. This often occurs in patients who are irregular in menstruating. It is difficult to say just what pathological lesions are present in these cases. Many of these cases occur in persons who are illy developed as regards their nervous systems. They need improved hygienic surroundings, good food, and evenness of life. They should not be idle, but the occupation should not be arduous. If young, great attention should be given to proper schooling of the mind, so that it may be better developed, when it is less liable to disturbances. Mental shock, moral disturbances, indigestion, worms, ovulation, etc., may disturb the nervous system sufficiently to cause these attacks. It is probable that this patient masturbates. Patients who have migraine often escape epilepsy, and *vice versa*. If you can carry the patient over one attack it is a great gain for the nervous system, as every attack weakens, and increases the liability to others. Finally, for these cases the ovaries may be removed, but this should be done only as a *der-rier ressort*.

ORIGINAL ARTICLES.

BORACINE.

BY W. THORNTON PARKER, M.D.,

OF NEWPORT, R. I.

The earliest of medical writers knew of the antiseptic qualities of boric acid, and its healing and purifying properties have been recognized for centuries. It has been recommended for use in solution, in powder and in ointment for the treatment of wounds, burns, and sores of all kinds, and yet when the dangerous properties of carbolic acid, iodoform, and the bichloride solution had been proved over and over again, and no satisfactory substitute discovered, borax still remained in great obscurity. Although its value was known so well, yet it was not considered worthy of rank as an antiseptic, and few believed that it possessed germicide properties.

The writer first became interested in the subject of antiseptics while attending the surgical clinics of Prof. Lister, in Edinburgh, in 1870, and subsequently in the Continental hospitals opportunities were afforded for witnessing the effects of carbolic acid, salicylic acid, iodoform, bichloride solution, etc. In the hospitals of Bonn, Leipzig, Vienna, Freiburg, Tübingen, Munich, Heidelberg, Würzburg, Paris, London and Dublin, I watched the development of the antiseptic method for surgery which Lister had so brilliantly brought forward. Each hospital had its individual procedure for the protection of wounds. The Charing Cross Hospital, London, in 1882, had developed the value of boric acid in combination with glycerine,

and since that time to the present the writer has found no preparation more satisfactory for antiseptic purposes than that now manufactured by Theodore Metcalf & Co., of Boston, viz.: boracine.

Boracine is composed of boric acid, glycerine, menthyl salicylate, menthol, thymol and eucalyptol. For surgical purposes it is best prescribed in solution, one part of boracine to twenty or thirty of water. It can be used, however, from 10 per cent. to 50 per cent. in strength as may be indicated. Under its use wounds of all kinds heal as rapidly and as satisfactorily as under carbolic acid dressings; but with boracine there is no danger from poisoning, and the wound and the surrounding tissues are not inflamed or injured. It is, in fact, entirely innocuous when applied to a wounded surface. For minor surgery it is useful as an ointment combined with cosmoline, vaseline, or lanolin, the new basis for ointments.

R.	Boracine.....	3iss.
	Ad ps.....	5ij.
	Lanolin.....	5iv.
	M.	

Boracine is useful as an application in chronic otorrhœa, in diseases of the nasal passages, chronic catarrh, etc., and for diseases of the rectum. I have found it very useful as an application for chronic cases of pruritus ani, as well as in general cases of prurigo. In such cases it will often afford perfect relief where other remedies have failed. For the treatment of ulcers of the leg nothing could be more satisfactory. It should be used as a thick paste completely covering the sore. I direct the patient to carefully and thoroughly wash the ulcer and the surrounding skin with the best Castile soap, then apply the pure boracine, and then to bandage the leg with the solid rubber bandage.

For injection in cystitis it is to be highly recommended. For use in diseases of the eye it is excellent. For the treatment of diphtheria in 25 per cent. solution for the steam atomizer I have found it excellent, and altogether preferable to any other preparation known to me. For an injection in the puerperal state nothing could be better than a 25 per cent. solution of boracine. This can be used with safety, while other preparations like carbolic acid have often proved injurious and even dangerous. In gynecology it is especially useful in solution as an injection for leucorrhœa, vaginitis, etc. It is an excellent application for ulcerated cervix, spread thickly on the diseased surface. It can also be spread on absorbent cotton and introduced and applied in that manner. After considerable experience with the various preparations used for local treatment in these disorders, I find none yielding such satisfactory results as the boracine. The cleanliness and the gentleness of this remedy, and the steady improvement resulting from its use, will be very satisfactory to both patient and physician.

Messrs. Metcalf & Co. have, at my suggestion, made vaginal and rectal suppositories of boracine in combination with gelatin, and pencils of the same for intra-uterine and other uses. These contain 25 per cent. of boracine. The suppositories are of three sizes—Nos. 1, 2, 3. No. 1, the smaller, is a useful suppository in various rectal disorders, particularly

in piles, and I have also used them with great success in the treatment of "thread-worms." No. 2 is of medium size, and can be used either for the rectum or for the vagina. Where a larger amount of boracine is required, in cases of chronic leucorrhœa, vaginitis, or uterine disease, No. 3 suppository is indicated. The larger suppositories should not be used more than three nights in succession, and experience has taught me that the best results are obtained usually by using them every other night.

In chronic eczema of the scalp I have found the action of boracine remarkably good. Cases which have refused to improve under ordinary treatment respond quickly to the use of boracine ointment. I direct the head to be shaved and carefully washed with Castile soap, and boracine ointment, 15 to 20 per cent., applied. The washing with Castile soap and careful drying is to be attended to night and morning, and then the ointment is applied with the most satisfactory results. In many obstinate diseases of the skin boracine will be found very valuable.

So many new antiseptics have come into notice of late that boracine will be viewed with some doubt, but the writer feels confident that the experience he has had with the new antiseptic justifies him in praising its merits and in cordially recommending it for use by the medical profession. To say the least, we have no antiseptic better at hand at present, and, in the writer's opinion, nothing which can equal it in general usefulness.

Newport, R. I., April, 1886.

MEDICAL PROGRESS.

THE ANTISEPTIC TREATMENT OF WOUNDS.—At the close of an interesting article on this subject SURGEON-MAJOR W. GRAY calls attention to the fact that in a long series of cases the latter half were much more successful, and this greater success appears to coincide substantially with the disuse of the waterproof covering, and with the adoption of dryer and thicker absorbent dressing. Accurate adaptation of the flaps I regard as of great importance. I now always mark them out with a scalpel, in order to insure evenly cut edges, and when the skin has contracted they are completed by transfixion in the usual way. My experience has taught me, too, that antiseptic applications direct to the wound are in many cases unnecessary. I allude especially to those cases where the tissues are perfectly healthy. Still they do no harm, and, by way of making "assurance doubly sure," may be employed in every instance. All I mean to convey, however, is that we can in a certain class of cases obtain as good results without as with them, provided we attend carefully to other precautions. I never under any circumstances use the spray. Moisture in any shape or form I look upon as one of the most fertile sources of failure. Our whole efforts should be directed to preserve both the wound and the dressing in the driest condition possible. The essentials, then, of my present procedure are cleanliness, infrequent dry dressing, free drainage, accurate adap-

tation of flaps, and rest. Under the last mentioned, pressure by means of bandages may be included. To prevent putrefaction of bloody and serous discharge I prefer to have the dressing medicated with some unirritating stable material. Of all I have tried, boracic acid or iodoform is decidedly the best. The latter is more expensive, however; hence I have found boric cotton more convenient to use. In order to press the flaps close together, and thus lessen or avoid exudation, I draw the bandage as tight as possible; the thick elastic pad of cotton and oakum underneath will always prevent it being drawn too tight. So effectually does this answer its purpose that I have on more than one occasion found the original dressing, on first exposing the stump, almost perfectly dry and clean. Oakum, besides being more or less antiseptic, keeps the cotton together, and enables us to save the use of the latter to a certain extent as well. Wood-wool, although a most excellent absorbent dressing in suitable cases, is somewhat unmanageable in amputations. Made into pads, it does not fit closely enough; and employed loose, it is difficult to apply, and is inclined to get lumpy or become displaced. To ensure complete rest of the parts while the healing process goes on, the limb, if it be a lower one, is fixed in a heavy box splint and well packed round with cotton. Should the upper limb be operated on, the stump is securely fastened to the side of the trunk, with plenty of cotton interposed and placed around. The cotton packing, as well as the thick dry dressing, preserves the normal heat of the parts, a matter which I believe is essential to rapid and perfect union. Lowering of the temperature by wet dressing, exposure, or cold applications tends, I am convinced, to delay or prevent union by first intention. During the progress of the case the thermometer is used morning and evening; if the temperature remains normal, or does not rise beyond 99.5° F. occasionally, all is going on well, and the original dressing need not be disturbed sooner than the ninth or tenth day. When removed the flaps will be found firmly united. The drainage-tube may now be withdrawn, its track gently syringed out with carbolic or sublimate lotion, a little iodoform introduced, and the stump redressed with oiled lint or with boric cotton, taking care to sprinkle either with iodoform. Some of the unsoiled boric cotton from the original dressing will do quite well enough. A pad of oakum and a bandage complete the second dressing. This may remain on for four or five days longer, and when removed the track of the drainage-tube will have closed. The sutures may now be taken out, the stump being practically healed. Should the temperature, during the course of the second or third day, rise to 100° or beyond, and remain so, the antiseptic dressing has probably failed and must be removed, the stump being treated in the ordinary way, or, better still, with wood-wool and iodoform. Before doing this I always administer quinine, as the fever may be of malarial origin, and not traumatic. In more than one instance this has proved to be the case. The precaution I note here is, however, scarcely necessary, except in a climate where malaria prevails.