which the ordinary treatment was adopted—in the second, with complete success, in the first, with a fatal result. I think there cannot be a doubt that the immediate cause of death was the excitement and exertion caused by her husband's imprudence in going to see her; added to this, she had, for some time past, suffered considerable deprivations. It is probable that a more favourable result might have obtained, but for these two circumstances; for during the whole of the time the symptoms were not so urgent as in the third case.

In case three, the placenta was not over the os, but close to its edge. As the hæmorrhage continued after the discharge of the "waters," and as the os was so high up as only just to be reached, the ergot seemed indicated, and it produced a favourable progress till eleven A.M., on the 17th, up to which time I trusted she would be delivered by the natural process, as the head descended slowly, and the hæmorrhage gradually ceased. At this time symptoms recorded induced me to repeat my examination, when I found the placenta almost plugging up the os. This, and the other symptoms, proved that there could now be no doubt that immediate delivery was imperatively called for; but the firm (hour-glass?) contraction of the uterus on the child, offered a serious obstacle to an easy and speedy delivery; and as the hæmorrhage still continued, I determined to detach the placenta as I proceeded. With the detachment of the placenta, the hæmorrhage immediately and entirely ceased, and did not recur. Might delivery have been left to nature, after the detachment of the placenta? From subsequent reflection, I think that very probably it might; for the bleeding never recurred, and I conceive that delivery by the natural process would not have made a more serious demand upon her strength than the painful and tedious operation of turning. At the time, however, and under the urgent circumstances, I feared to leave the woman undelivered longer than I could help; as, at the time, the loss of a very small quantity of blood might have turned the scale against her.

I consider the detachment of the placenta to have been a most important step in this case, as it put a complete stop to the hæmorrhage by which life began to be threatened; and I am disposed to think that had delivery been effected in the ordinary way, and had the hæmorrhage continued, as it most probably would have done, during the operation, that death might have resulted. I, for one, feel thankful to Drs. Redford and Simpson for having called our attention to the propriety of detaching the placenta; and I believe that in placental presentations, where turning would be found impracticable, or very difficult, or in which the balance is so nearly drawn as to render immediate arrest of hæmorrhage important, the detachment of the placenta will be found to be of great service.

Dudley, December, 1845.

THE TREATMENT OF UTERINE HÆMORRHAGE.

By JOHN CRAIG, Esq., M.D., Paisley.

IN THE LANCET for Nov. 22, 1845, p. 558, two cases of supposed pregnancy at the third month, accompanied with hæmorrhage, are reported as having been attended by Dr. Hall Davis. I have little to say regarding the mode of treatment pursued by Dr. Davis, in the management of these two cases; indeed, in the first of them there was some reason to be alarmed respecting the safety of the patient, and in my opinion the treatment was prompt and judicious as far as the mother's safety was concerned. It gives me much pleasure to observe that so experienced and active a physician as Dr. Davis should, along with Professor Murphy, take the lead, in the metropolis, of administering opium in uterine hæmorrhage—a medicine of the greatest utility in such cases. The successful issue of those cases that are treated by the employment of suitable doses of opium, compared with others in which it is not used, will, I entertain no doubt, compel those practitioners whose minds are still prejudiced against the employment of this salutary drug in uterine hæmorrhage, to administer it in order to preserve their own reputation.

From the efficient manner in which Dr. Davis advises the application of the plug in hæmorrhage, accompanied with symptoms of abortion, there can be little doubt of its efficacy; but so far as my experience extends, it is never necessary in the early stages of pregnancy; and from the Doctor's own statements, it must be injurious in some cases, for he maintains that its operation not only arrests the discharge, but excites the action of the uterus to throw off its contents. In Dr. Davis's two reported cases of supposed pregnancy, threatening miscarriage, he informs us that he plugged the vagina, and gave ergot of rye, with the view of bringing off the contents of the uterus. In the early part of my experience in the

management of such cases, when considerable discharge had taken place previously to my being called to the patient, and especially when the discharge was accompanied with considerable pain of back and lower portion of belly, abortion, in almost every instance, was the result.

Latterly, however, several cases of a similar description have come under my care, and two of them very lately. The first of the two latter cases had flooded partially for several days before I was called to see the woman, but the day on which I made my first visit the hæmorrhage was alarmingly profuse, accompanied by severe pain at the lower part of the abdomen; the pulse could scarcely be felt; the skin was cold, and the countenance shrunk. The woman was in the fifth month of her pregnancy, and the motions of the child were distinctly felt. On examination, the os uteri was not at all dilated; but the uterus, above the pubis, felt like a round very hard ball. I enjoined that the patient should be kept perfectly quiet in the recumbent position, that free ventilation be allowed in her room, and that her drinks should be cold and acidulated with diluted sulphuric acid, and that cold wet cloths should be applied to the lower part of bowels, and over the genitals. At the same time, sixty drops of a solution of the muriate of morphia were given to the patient, and the same quantity was ordered to be repeated if required. By this mode of proceeding, the pains and hæmorrhage ceased, the patient became easy, and the natural heat returned to the skin. From this time, the patient had not a bad symptom. This woman continued in a tolerably good state of health until about the eighth, or a little after the eighth month of utero-gestation, when she was safely delivered of a fine healthy boy. Now, had plugging in this case, and ergot of rye been used, abortion, in all likelihood, would have been the result.

The second case was somewhat similar to the above; but the patient was more robust, and had not lost quite so much blood, yet the pain and discharge had continued about twenty-four hours before I was called to the patient. As the pulse was still pretty full and strong, twelve ounces of blood were taken from the arm, and, in every other respect, she was treated as the former case. She likewise went to her full time, and bore a healthy child.

This is a simple and safe mode of procedure in such cases, and is certainly fraught with great advantages over the method by which floodings are generally managed. To arrest the expulsion of the child at a time when it is not viable, and to protract the delivery to near the full period of utero-gestation, so that a living and healthy child is born, often creates a consummation of great interest to those more immediately concerned.

In THE LANCET for December 13th, 1845, p. 644, a case is reported, headed "On the Treatment of Placenta Prævia. By John L. P'ou, M.R.C.S.E., Lambeth."

In this case, uterine hæmorrhage came on about the sixth month of pregnancy, and seems to have been arrested for a month, when it again recurred to a considerable extent; yet the os uteri had dilated only so far as to barely admit the introduction of the index finger of the right hand. The os uteri was rigid and unyielding, and there did not appear the least disposition to dilatation. In such circumstances the plug was used; still no expulsive efforts of the uterus occurred for five hours afterwards. About this period the pains were considerable, but on removing the plug they were again suspended, and did not return until two doses of ergot of rye were exhibited, showing, to some extent at least, that the presence of the plug was the cause of the pains previous to those which succeeded to the use of the ergot, by which the expulsive efforts were produced. After the removal of the plug there was no return of the hæmorrhage, until expulsive pains came on after the ergot had been administered. The seventh month of gestation had been completed; the child was considered to have been dead three or four hours previous to its birth. Mr. P'ou states, in the first column, "The preservation of the life of the fœtus does not appear to me a question for consideration; for as very few females arrive at the full period of utero-gestation in these cases, (premature labour being almost an invariable consequence,) in the majority the life of the fœtus is almost certain to be forfeited." This statement regarding such cases is quite at variance with my experience; for in almost every case of this description which I have seen, the patient, before it was considered necessary to deliver, had advanced to the eighth, or near the eighth month of gestation—a period at which, in such cases, the half of the children have been born that I have attended, not only alive, but were reared, at least as long as they remained within the range of my knowledge. My experience, then, enables me to state my opinion pretty confidently, that if women under such circum-

stances be confined to the recumbent posture, kept quiet, supported by mild nourishment, have cold acidulated drinks, cold applications to the lower part of the abdomen, and over the genital organs, with repeated full doses of laudanum, as necessity may require, so as to subdue pain or uneasiness, and arrest the discharge, the recurrence of each attack of hæmorrhage may be so completely checked as to enable the woman to go to about the eighth month of utero-gestation, when, generally, both she and her child may, by artificial delivery, be preserved alive. I have had only one case of the death of the mother in labours of this description, when there was anything like a fair prospect of saving both the mother and the child. This woman's mind was fixedly prepossessed with the belief, that if she were delivered artificially she would assuredly die in consequence of it; and, previously to delivery, she took leave of her husband under this impression. Whether such an appalling state of mind could have any effect in producing the mournful result, is not for me to decide. She was not by any means too much exhausted by previous loss of blood, and delivery was effected with great facility. The child, also, was strong and active at birth, yet the mother seemed quite indifferent at the result. The after-birth lay loose in the vagina, and only required removal.

The patient now seemed so well that the medical friend who kindly assisted me left the house, and a short time afterwards, I left the room for a few minutes, when, on my return she was sunk and ghastly. The discharge, to a considerable extent, had suddenly come on, and though it was at once subdued, it seemed, from all the previous circumstances, to be more than she could bear, and she died about two hours afterwards.

It will now be very obvious, that if the mode of treatment pursued by me be necessary to the completion of utero-gestation, so far at least as to give a fair chance to the safe delivery of the mother of a viable child, the method adopted by Mr. P'on precludes the probability of effecting this very desirable result; for he clearly, as it were, compels the uterus to expel its contents at an early period of pregnancy, and under the impression, "that the preservation of the life of the fœtus does not appear to him a question for consideration."

On the subject which has lately excited so much controversy,—namely, the delivery of the placenta before that of the child,—I am not prepared to give any decided opinion, never having seen a case when the placenta was expelled before the expulsion of the child. I may here remark, that in such presentations no particular difficulty ever occurred to me in turning and delivering the child, for if the attacks of floodings be properly managed until the period at which the woman must be delivered, the os uteri, from the relaxed state of the parts, will be pretty easily and safely dilated by cautious perseverance in the introduction of the hand.

I do think, however, that Dr. Simpson is at least entitled to the grateful acknowledgments of the profession, for calling their attention so ably and truthfully to the subject. He only advocates the occasional employment of the plan he espouses, and from the various facts advanced, it does appear that it may be sometimes useful. At all events, in urgent cases of difficulty and danger, it is of the utmost importance that we should be acquainted with every suitable kind of information; and any idea that adds to our stock of useful knowledge should be received, not with frivolous objections, but with a candour becoming a sound philosophic mind.

Paisley, December, 1846.

A REPORT OF A CASE OF OVARIAN DROPSY SUCCESSFULLY TREATED ON MR. BROWN'S PLAN OF CURE, UNDER UNFAVOURABLE CIRCUMSTANCES.

By THOMAS HUNT, Esq., Surgeon, Herne Bay.

MRS. F—, aged thirty, the wife of a bricklayer in this place, was delivered of her fourth child in the month of January, 1841. She was then residing at Ingatestone, and Mr. Cornelius Butler, who attended her on that occasion, afterwards gave her the following memorandum in writing—viz.,

"In Mrs. F—'s last confinement, the progress of the head of the child was much impeded by a tumour in the recto-vaginal septum. After some difficulty I delivered her with the forceps. I merely write this to sanction the puncture of the tumour, if it be necessary, at any future time, as it has increased considerably, by the testimony of the gentleman who attended her previously, and with whom I am acquainted.

"Ingatestone, Jan. 1st, 1841."

"C. H. BUTLER.

The patient subsequently observed an enlargement on the left side of the abdomen. In the autumn of 1843 proving again pregnant, she consulted Dr. Blundell, who is said to have advised her then medical attendant to have recourse to

premature delivery. This was accomplished in the sixth month of gestation. From this time the abdominal tumour increased more rapidly.

In the autumn of 1845 she placed herself under my care.

October 15, 1845.—The abdomen measures thirty-four inches, parietes very thin, umbilicus prominent, fluctuation very distinct. The health is moderately good, and the menstruation regular. The os uteri is slightly tumid, and has rather more sensibility than usual, but presents no indication of disease. She has the appearance of a woman in the seventh month of pregnancy.

Having read the cases detailed in *THE LANCET*, by Mr. Brown, of Paddington, in which he succeeded in curing encysted dropsy, by pressure, mercurial treatment and diuretics, and the age of the patient and other circumstances appearing to justify the experiment, I requested my patient to wait upon Mr. Brown, who reported that he considered his mode of treatment to be highly appropriate in the case, and requested me to commence the treatment without delay.

22nd.—The abdomen was tightly bound with a flannel bandage, which was renewed night and morning. One drachm of the strong mercurial ointment was directed to be rubbed into the abdomen every night, a diuretic mixture ordered to be taken twice a day, and three grains of blue pill night and morning.

24th.—The diuretic having produced vomiting, the medicine was administered in the effervescing form. She has great uneasiness in the chest, from the tightness of the bandage.

28th.—She complains of vomiting and pain in the jaws. The gums and throat are very sore, and a copious salivation has set in. The quantity of urine passed for the last three days exceeds that of the fluid swallowed. The abdomen is decreased in size about two inches. Discontinue the mercury.

30th.—The mouth continues excessively sore; the soft palate is extensively, but superficially, ulcerated; the tongue can scarcely be protruded; and the salivation is unabated. Flow of urine still copious. The stomach rejects all medicine, except an occasional dose of aperient pills. The menses have appeared at the end of a fortnight from the last period. The abdomen measures less by three inches than at the beginning of the treatment. Milk diet ordered.

November 4th.—The soreness of the mouth is beginning to amend slowly. Urine smaller in quantity since the diuretics have been rejected, which has happened in every form of their administration.

11th.—Improving in health; is able to take more nourishment. Secretion of the kidneys diminished; bowels costive. The abdomen, having previously measured less by three inches and a half, is now somewhat increased in size. The bandage has been constantly applied, but cannot be borne very tight. Ordered a dose of aperient pills.

12th.—The cyst was tapped by Mr. Brown in my presence, and nearly nine pints of fluid were drawn off. It was almost as clear as water, and as colourless. When treated with nitric acid at a high temperature, it exhibited a slight albuminous cloud, scarcely visible at the time, but distinctly precipitated on the following morning. She bore the operation without fainting, and was put to bed, tightly bound with a flannel bandage, a book, enveloped in napkins, being applied to the umbilical and pubic regions, as a compress.

13th.—She has had a tolerable night, but complains of headach. She has passed five pints and four ounces of urine since the cyst was tapped, and has taken only a pint and a half of fluid. The bowels being costive, a dose of aperient pills was ordered, and some gin-and-water.

14th.—The pills excited vomiting, but have acted on the bowels. Complains of the corners of the book hurting the ribs and hips. A large toilet pin-cushion was substituted, and the bandage re-applied. She has nearly lost her appetite, feels weak, and is distressed with flatulency. Ordered a glass or two of wine.

15th, 16th, and 17th.—Has been constantly harassed with flatus. When the bandage is withdrawn, she eructates an immense quantity of gas. Urine about equal to the fluids taken.

18th.—Appetite entirely gone. She has slept but little, is much depressed in spirits. Pulse 120; skin hot; abdomen somewhat tender; urine scanty; bowels costive. A cathartic pill was ordered, and an effervescing draught every fourth hour.

19th.—Better to-day. Skin cool; pulse weak, and slower. Complains of noises in the head; has drank about a quart of fluid, and passed less than that quantity of urine. The bowels have been freely relieved. An attempt was now made to resume the mercurials and diuretics, but it was found impossible to persevere. The stomach became excessively irritable.

24th.—It is proper to mention that, about three weeks ago, one of her children returned from a visit to town with slight