cent., had mitral regurgitation, functional or organic. Of these, 496 were treated dietetically to reduce their weight. In 58 the murmur disappeared together with all signs of cardiac distress and usually with a marked drop in blood pressure. Most obese patients have a more or less elevated blood pressure which promptly drops to approximately normal, under treatment, provided there is no other cause for the hypertension.

Insurance companies do not reject fat people because of sentiment, as per the old adage that "Nobody loves a fat man," but because they are poor insurance risks, on account of their proneness to develop disease of the heart and arteries.

I mention this class of cases because so little has been written about them, and because in my opinion, they are all *potential* cases of myocardial disease, unless appropriate treatment is instituted.

REFERENCES.

1. Emerson, Haven: The Prevention of Heart Disease—A New Practical Problem. Boston Med. & Surg. Jour., June 9, 1921.
2. Mackenzie, James: Diseases of the Heart, 1910.
3. Cabot, R. C.: The Four Common Types of Heart Disease. Jour. Amer. Med. Assoc., Oct. 24, 1914, p. 1461.
4. White, Paul D.: Discussion on Cardiac Diagnosis before Cambridge Medical Improvement Society, April 25, 1921. (Not published.)
5. Christian, Henry A.: Chronic Myocarditis. Jour. Amer. Med. Assoc., June 22, 1918, p. 1909.
6. Smith, Wm. H.: Chronic Adherent Pericarditis. Jour. Amer. Med. Assoc., Sept. 6, 1913, p. 739.
7. Lewis, Thomas: The Mechanism and Graphic Registration of the Heart Beat, 1920.
8. White, Paul D.: Diagnosis of Chronic Valvular Disease. Jour. Amer. Med. Assoc., April 30, 1921, p. 1207.
9. Christian, Henry A.: The Significance of Arrhythmias and Systolic Murmurs, etc. Jour. Amer. Med. Assoc., Nov. 29, 1917, p. 750.
10. Reid, Wm. D.: Jour. Amer. Med. Assoc., Feb. 12, 1921, p. 432.
11. Strumpel: Textbook of Medicine, 1921, Vol. I, p. 481.
12. Monographic Medicine, 1916, Vol. 5, p. 433.

ANKYLOSIS OF THE JAW.

By R. H. GILPATRICK, M.D., BOSTON.

[From the Surgical Department of the Boston Dispensary.]

A. N., a Holland boy of fifteen years, was brought to the clinic on May 10, 1921, with the following history: At the age of one year he had scarlet fever, complicated by infection of both mastoids, for which some six or seven operations were performed at the time and subsequently up to the age of six, when he migrated to Canada with his family. During the following three years there would appear to have been performed twelve more operations of varying sorts, the last by Dr. Starr at the Children's Hospital in Toronto. This last was a radical mastoid. The notes of the hospital at that time describe an almost complete ankylosis of the inferior maxilla, and the history from the parents would indicate that some marked degree of limitation of motion was evident from the time of the first operations during the second year of life, or fourteen years before coming under our observation.

It is evident from superficial examination at this time that very extensive suppurative processes have been endured, for there are scars over both mastoids, in front of both ears, on the sides of the neck and cheeks, one forward below the angle of the mouth. There is a complete right-sided facial paralysis, evidently as a result of suppuration of the middle ear and in front of the external auditory meatus and possibly from the numerous incisions in that area which have from time to time been made for drainage. The inferior maxilla is markedly undeveloped, as shown in the photographs and the skiagrams. The angles are of the infantile type. There is an unerupted molar on each side, both inclined at a considerable angle with the crowns forward and impinging upon the next teeth in front. There is complete ankylosis with not the slightest degree of mobility either up and down or from side to side. This is said to have been so for the past six years, or since soon after the last operation. There is considerable impairment of hearing on both sides, more marked on the right. The skiagrams of the temporo-mandibular joints are unsatisfactory, as they are likely to be. They indicate, however, a bony union on the right side and a fibrous ankylosis on the left.

How the boy has been able to sufficiently nourish himself, particularly these past six years, is difficult to understand, for he is generally very well developed for his age and is nearly six feet in height. He has had several teeth removed for the purpose of obtaining an entrance for food, but the under jaw being so short, the lower incisors and all teeth save the last molars on each side below fail to meet those above, and the result is that the lower jaw shuts past the upper, the lower incisors being firmly indented into the hard palate about one cm. back of the upper incisors. This has very seriously narrowed the buccal cavity, so that he has very little room for manipulation of the tongue and still less for the reception of food. His method has been to very thoroughly macerate all food, confining himself to such as could be so treated, form it into balls or small masses, introduce these between the lips in front of the gap left by removal of the teeth, then painstakingly force the food through as well as possible by manipulation with thumb or finger, and then carry it about with the tongue to be almost immediately swallowed. It has been a slow and tedious process, as testified to by his mother, and what it must have meant to one with the appetite of an actively growing boy of his age can scarcely be comprehended. He is not at present in any apparent danger from actual starvation, but the handicaps of his present condition are very obvious. Never to have known the satisfaction of a meal, never to have been able to eat other than foods adaptable to complete maceration in the plate, never to have been able to express himself save through tightly clenched and overlapping teeth, without room enough in the mouth even for active use of the tongue, such a combination of distress has made

Before operation, showing complete ankylosis, with lower teeth impressed into palate inside upper teeth.

Before operation, showing deformity from non-development of lower jaw, caused by disuse.

Before operation, showing deformity

Before operation, showing deformity.

Ten days after operation. Extent of voluntary movement.

Ten days after operation, showing scars.

him quite heedless of his facial palsy. But in spite of it all he has evidently persisted in a life of activity, both mentally and physically, and is above the average in intelligence.

After some study of the case, it appeared probable that we would have to deal with an extensive union of the condyle of the lower jaw to the glenoid, some considerable involvement of the coronoid insertion of the temporal muscle in the osseous overgrowth all on the right side, and a firm, fibrous ankylosis on the left. Realizing the likelihood of the fibrous ankylosis offering very serious difficulties, we yet decided that the right side would better be attacked first. Some unusual freedom of movement during operation was offered on the right by reason of the seventh nerve, already destroyed, not requiring such careful attention and avoidance. It was determined, however, to proceed, so far as possible, as if the nerve were normal. It was hoped that division of the union about the joint would be sufficient to free that side, and that the left side could then be attacked at the same sitting.

In May, 1921, gas and ether were given by Dr. Temple, at first by the inhaler and later through a tracheal tube, and throughout the rather long operation there was at no time the least limitation of the operative procedure from the anesthesia. An incision one and a half inches in length was made in front of the right ear from a point one-half inch below the zygoma upward and just sufficiently forward to avoid the temporal artery and auriculo-temporal nerve. This was carried down to the bone and all bleeding secured. All further dissection was carried out, so far as possible, with the idea of elevating the parotid and continuing the work beneath it. It was soon demonstrated that the hypertrophic process was much more extensive than had been realized. The whole space beneath the zygoma and extending forward to completely envelop the coronoid process was filled with firm, new bone. The temporo-mandibular articulation was surrounded by a dense mass of the same. A second short incision over the anterior extremity of the zygoma, through which the zygoma could be there divided with the motor saw, made it possible to elevate the zygoma and secure added space. The mass of new bone was gradually peeled and burred out, the condyle of the jaw amputated and the space freed, but still the jaw was as firmly fixed as ever. It became evident that the temporal muscle must be sacrificed, and the coronoid process was therefore removed, leaving the lower jaw without support by either temporal or masseter. When this was thoroughly done, it was discovered that the mouth could be opened. A tonsil gag of the common type was easily inserted and the jaw forced to extreme depression. This demonstrated that the left side was uninvolved and required no treatment. The space above the stump of the condyle of the inferior maxilla and between it and the floor of what had once been the glenoid was further en-

larged and the pterygoids thus brought clearly into view were seen to be apparently normal.

The next step was the preparation of a flap of fat and fascia for the new articulation. This was secured from beneath the skin anteriorly to the original incision, and having as wide a base as possible and the base downward with the belief that by so cutting a better blood supply could be expected. A flap of quite sufficient proportions was thus freed, carried into the cavity over the stump of the condyle and sutured to the pterygoid muscle with fine plain gut. Movement of the jaw was now quite free, so the wounds were closed with a small tissue drain in one. The middle section of the zygoma had been completely freed and was removed, after at first planning to secure it in place by drilling and suture. It was deemed wiser not to prolong the operation or to assume the danger of sepsis being increased.

The immediate surgical convalescence was surprisingly easy. The patient was able to walk to an ambulance on the afternoon of the day of operation, for transfer to another hospital where he had of necessity to go. He was able to move the lower jaw voluntarily and without serious pain from the start. There was considerable œdema of the face for two days. The drain was removed after twenty-four hours, and healing was by first intention. By the fifth day he could masticate soft material, and he left the hospital on the eighth day. On two occasions during the first few days a gag was inserted and the jaw carried fully open to guard against the formation of limiting adhesions. From the time of his hospital discharge the usefulness of the new joint has steadily increased. The photographs indicate the degree of motion ten days after operation. He now chews everything, and save for the handicap of the facial palsy is not seriously inconvenienced.

It is believed that some degree of development in this jaw is still possible with normal use. There will, of course, always be the deformity of the receding chin and the facial palsy, but his ability to masticate food and to speak distinctly has been restored to him, and he has been restored to the society of his fellows.

It is evident from the notes of the Children's Hospital of Toronto that it was expected that this boy would return to them for arthroplasty after his recovery from the radical mastoid, but in the meantime his parents again migrated to Massachusetts. The inspiration for undertaking this case came to us from the late John B. Murphy, as so much has come to many for a long time. In his *Surgical Clinics*, Vol. I, No. 6, December, 1912, he reports a case which in many respects was so nearly identical with this one as to be remarkable. The age, sex and duration of the disease are the same in both. In his case there was osseous ankylosis on the right and fibrous on the left. The cause was undoubtedly very similar in both.

Middle ear suppuratation, long continuing and

spreading forward, as is so much more likely to be the case in early life than at a later time when the cranial bones have closed up and hardened, entered the temporo-mandibular articulation and after a chronic course resulted in this chronic hypertrophic arthritis. As time went on and the infection persisted in one spot or another, the osteoarthritis extended and evidently resulted in complete ankylosis at about the age of nine years, or six years before our observation. It is worthy of remark that the masseter, temporal and pterygoids on the left side could remain in disuse for so long without permanent impairment of function. In Dr. Murphy's case it was necessary to perform the arthroplasty on the one side and to very extensively free the fibrous ankylosis on the other; not until masseter, temporal and pterygoid insertions had been severed could the mouth be opened. In case of osseous union on both sides it seems doubtful how useful a double arthroplasty of this sort would prove to be, chiefly because of the probability that with free motion up and down a very annoying freedom from side to side would be the result. But so long as one side is normal, or showing but a fibrous ankylosis, the usefulness of the operation is manifest at once. Restoration of function in any ankylosed joint is a desirable object of the greatest importance, the fundamental importance varying with the importance of the involved joint. The comparative rarity of complete temporo-mandibular ankylosis among similar joint lesions has offered comparatively little material for the fat-fascia flap method of arthroplasty in that joint, but save for the difficulty of the needed dissection and the necessity of securing great freedom of movement in a space where free access is limited by the proximity of structures which allow of no inaccurate work, the operation offers a probability of a result completely satisfactory in a disability most distressing, if not actually threatening to life.

THE PUBLIC HEALTH STATUS OF AMOEBIC DYSENTERY IN THE UNITED STATES AS POTENTIALLY INFLUENCED BY THE WORLD WAR.*

By C. W. STILES, WASHINGTON, D. C.

Professor of Zoology, U. S. Public Health Service.

One of the results of the hookworm campaign has been a more general adoption of routine examinations in clinical laboratories, not only in this country but abroad also. Since any examination for intestinal worms, properly carried out by a competent microscopist, is likely to uncover infections with various protozoa, it is not unnatural that a more general interest has been awakened in the subject of the protozoan infections of the intestine of man. Equally

*Cutter Lecture in Preventive Medicine, Harvard University, Jan. 17, 1922.