similar in all, I shall take the first case. Previous to my seeing the patients, the nurses had applied mustard cataplasms over the stomach and bowels, with friction to the limbs, and a draught of hot tea, with a spoonful of brandy, which was quickly rejected. I immediately ordered 15 grs. of calomel, with half a gr. of opium, to be formed into a pill, given directly and repeated, every quarter of an hour till four doses were taken, after which a pill, containing $1\frac{1}{2}$ gr. of calomel was directed to be given every quarter of an hour, with a tablespoonful of a mixture containing about two drops of laudanum, and the same of the sp. am. arom. after each pill. This plan was continued for 24 hours without intermission, at the end of which period, by gradual steps the purging ceased, the pulse returned, the livid colour disappeared, a general warmth and universal perspiration were produced, the voice grew stronger, and the improvement was so decided, that I ordered one pill only to be taken every hour, after twelve hours this was discontinued, the patient having no material ailment except debility; it was now necessary to give an aperient, the bowels being in a constipated state. He was discharged in four days well; no salivation ensued; the only troublesome symptom during the cure was excessive thirst; this made him incessantly call for drink, of which, if the nurses inadvertently gave too much, vomiting was excited; I found very small effervescing draughts the most efficient in removing it.

In this case then 219 grains of mercury were taken without any appearance of ptyalism being produced, or any unpleasant symptom.

The other two cases were treated exactly similar and with the same results, but being somewhat milder in their symptoms, a less quantity of mercury was necessary. The first case was considered by those who saw it as a hopeless one.

I am perfectly aware that it may be said the favourable results of three cases are as nothing in deciding a point of such magnitude; I admit it, but I conceive the above cases tend strongly to support the plan of Dr. Ayre and others, and strengthen the opinion of those who consider the mercurial method of treatment in this dreadful scourge and pestilence the best and most eligible.

August 29, 1832.

COLD WATER IN CHOLERA.

To the Editor of THE LANCET.

SIR,—I have been much pleased by the perusal of Dr. Hardwicke Shute's paper, detailing the important results of his treatment of the collapsed stage of malignant cholera, by the unrestrained use of cold water.

In the treatment of the patients in the cholera hospital of Warsaw, in the summer and autumn of last year, I had frequent opportunities of witnessing the distressing sufferings occasioned by a restricted use of liquids of any kind—a small cup (about two tablespoonfuls) of *salop* being all that was allowed, and that only at intervals of half an hour or an hour. The consequence was, that the thirst became almost insupportable, and the hospital continually resounded with the cry of “water”—sounds more heart-rending than even the sight of the most violent symptoms of the disease. The alleged reason of this restriction, on the part of the medical officer, being, the supposed injurious effects of distending the stomach, on the operation of the remedial agents administered, which were chiefly large doses of calomel. I then had my doubts of the propriety of depriving the unfortunate patient of what nature so urgently demanded. This opinion was subsequently confirmed, when I found that the treatment adopted at Berlin with the greatest success in the stage of collapse was cold affusion, and allowing, at the same time, the free use of cold water as a beverage. * * *

I am, Sir, yours, &c.,

CHARLES ROBERT BREE.

Ipswich, Sept. 1, 1832.

Observations on the Blood. By W. STEVENS, M.D.

Sur le Sang Humain. Par S. DENIS.

Sur le Sang. Par M. L. R. LECANU.—
(Concluded.)

Experimental Inquiries in Chemical Physiology. (Part I.) *On the Blood; with Remarks on the Nature, &c. of Cholera.* By HORATIO PRATER, M.D.. London: Highley. 1832. 8vo. pp. 304.

SINCE the publication of our first notice of Dr. Stevens' work (page 659), another volume has appeared on the same subject, which we find to be one well worthy to receive our early attention. We accordingly add the title to our former list, and include a notice of it in the present review.