Critical Temperature of Water.

IN answer to Mr. Martin's letter in your last issue (p. 80), Cailletet and Colardeau found the critical temperature of water to be 365°C ., the corresponding pressure being 200.5 atmos. An account of their experiments is given in Preston's "Theory of Heat," p. 384. S. GEOGHEGAN.

Dublin, November 29.

SPECTRUM OF A METEOR.¹

THE photographs of the spectra of the stars taken at the Harvard College Observatory as part of the Henry Draper Memorial differ in two respects from those ordinarily taken elsewhere. Instead of using a spectroscope with a slit, in which but one star is photographed at a time, a large prism is placed over the object-glass of the telescope, and thus spectra of all the bright stars in the field of view are obtained. The number of stars photographed simultaneously is still further increased by substituting for the object-glass a portrait lens like that used by photographers, only larger. The field of view is in this way increased from two degrees square to ten degrees square, and a photograph is obtained of the spectra of all the brighter stars in this large region. Many thousand plates, covering the entire sky, have been taken in this way at the Cambridge and Arequipa Stations of this Observatory. All have been examined by Mrs. Fleming, and, as a result, numerous remarkable objects have been discovered. One of the latest is the spectrum of a meteor which has thus been photographed for the first time. Since it is impossible to foresee when the bright meteors will appear, or what path they will follow, a photograph will be obtained only when one happens to cross the field of the telescope. A number of trails of meteors have been obtained, both here and elsewhere, when charts of the stars were photographed, no prism being used. When the prism was in place no meteor bright enough to leave a noticeable trail has heretofore been photographed on the many thousand plates examined. At about 11 p.m. on June 18, 1897, however, when the 8-inch Bache telescope at Arequipa was directed towards the constellation Telescopium, a bright meteor appeared in right ascension 18h. 19m., declination $-47^{\circ} 10'$, and passed out of the field at right ascension 18h. 29m., declination $-50^{\circ} 30'$. The spectrum consists of six bright lines whose intensity varies in different portions of the photograph, thereby showing that the light of the meteor changed as its image passed across the plate. The approximate wave-lengths of these lines are 3954, 4121, 4195, 4344, 4636, and 4857, and their intensities are estimated as 40, 100, 2, 13, 10, and 10, respectively. The first, second, fourth, and sixth of these lines are probably identical with the hydrogen lines $\text{H}\epsilon$, $\text{H}\delta$, $\text{H}\gamma$, and $\text{H}\beta$, whose wave-lengths are 3970, 4101, 4341, and 4862. The fifth line is probably identical with the band at wave-length 4633, present in spectra of stars of the fifth type and forming the distinctive feature of the third class of these stars. The third line, which is barely visible, is perhaps identical with the band at wave-length 4200, contained in these stars (*Astron. Nach.* 127, p. 1).

It will be noticed that of the four hydrogen lines in the spectrum of the meteor, $\text{H}\delta$ is the most intense. This is also the case in the spectrum of α Ceti and of many other variable stars of long period. In some variables of long period $\text{H}\delta$ and $\text{H}\gamma$ are equally intense, while in others $\text{H}\gamma$ is the more intense. In some stars of the first type in which the hydrogen lines are bright, like γ Cassiopeiæ, the line $\text{H}\beta$ is much more intense in the photographic spectrum than any of the other lines, while in the spectra of stars like P Cygni and η Carinæ, $\text{H}\delta$, $\text{H}\gamma$, and $\text{H}\beta$ are nearly equally bright. These

results show an important resemblance between meteors and stars having bright lines in their spectra, and may aid in determining the conditions of temperature and pressure in these bodies. Since bright meteors sometimes appear during the November meteoric shower, a special effort will be made to obtain photographs of them, both trails and spectra, on November 13.

November 8.

EDWARD C. PICKERING.

GEOLOGY AND SANITARY SCIENCE.

THE address recently delivered by Mr. W. Whitaker, F.R.S., before Section iii. of the Congress of the Sanitary Institute (*Journal of the Sanitary Institute*, vol. xviii., 1897, pp. 304-316), touches upon several matters of special interest at the present time. Section iii. deals with chemistry, meteorology, and geology, and Mr. Whitaker chose for his subject "Water," as being appropriate for such a triple alliance. Most important is his suggestion "that round each work for the public supply of water a certain tract of the water-bearing bed should be saved from the assaults of surface-contamination." The address was delivered before the Maidstone epidemic had occurred, but reference is made to it in a postscript, and no stronger support for Mr. Whitaker's proposal could have been given. He rightly observes that "we are yet without much information as to how far pollution may be able to reach along an underground course," but "it is here that geology comes in, for the nature of the surroundings of waterworks must be taken into account, often for some distance and to some depth." From a sanitary point of view the mapping of all the surface deposits on the six-inch scale by the Geological Survey is referred to as most necessary. A great deal has been done, but very much more remains to be done; and, as Mr. Whitaker observes, "a survey of the drifts in the London area on the six-inch scale is greatly needed."

It is a serious matter when sewage-farms and cemeteries are placed on porous strata, which are perhaps in adjacent tracts utilised for their supply of water; and yet, as Mr. Whitaker remarks, "sewage-farms and cemeteries must exist, until some other methods of disposing of waste material are not only found out, but are generally adopted."

Clearly there is need for greater control over the sites chosen for any of the purposes mentioned, and public sources of water-supply must be carefully safeguarded. The question of federation in the matter of water-supply in many localities is one to which attention is drawn, and this also is becoming urgent.

NOTES.

THE French Government, through its Embassy in London, has presented to Sir Archibald Geikie a handsome vase of Sèvres porcelain in recognition of the services rendered by him to the Geological Survey of France.

WE are pleased to hear of the foundation of the Zoological Society of Western Australia, with the object of establishing a Zoological Garden at South Perth in that colony. Mr. E. A. Le Souef has been appointed director of the new institution.

IT is announced that the Council of the Iron and Steel Institute have accepted an invitation from the Association of Swedish Ironmasters to hold the autumn meeting of the Institute next year at Stockholm. The meeting will be held in August, and, in view of the large quantities of Swedish iron and ores consumed in this country, there is no doubt that it will prove a very popular meeting. Previous autumn meetings of the Institute have been held at Belgium, France, Germany, Austria-Hungary, Spain, and the United States.

¹ Harvard College Observatory. Circular No. 20.