

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: CRTC2 polymorphism as a risk factor for the incidence of metabolic syndrome in patients with solid organ transplantation.

Authors: Quteineh L, Bochud PY, Golshayan D, Crettol S, Venetz JP, Manuel O, Kutalik Z, Treyer A, Lehmann R, Mueller NJ, Binet I, van Delden C, Steiger J, Mohacsi P, Dufour JF, Socal PM, Pascual M, Eap CB, and The Swiss Transplant Cohort Study

Journal: The pharmacogenomics journal

Year: 2015 Dec 8

DOI: 10.1038/tpj.2015.82

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

1 ***CRTC2 polymorphism* as a risk factor for the incidence of metabolic**
2 **syndrome in patients with solid organ transplantation**
3

4 CRTC2 and Post-transplant metabolic syndrome.
5

6 Lina Quteineh, MD, PhD (1), Pierre-Yves Bochud, MD (2), Dela Golshayan, MD, PhD (3),
7 Severine Crettol, PhD (1), Jean-Pierre Venetz, MD (3), Oriol Manuel, MD (2,3), Zoltán
8 Kutalik, PhD (4,5), Andrea Treyer, PharmD (1), Roger Lehmann, MD (6), Nicolas J Mueller,
9 MD (7), Isabelle Binet, MD (8), Christian van Delden, MD (9), Jürg Steiger, MD (10), Paul
10 Mohacsi, MD (11), Jean-francois Dufour, MD, PhD (12), Paola M. Soccac, MD (13), Manuel
11 Pascual, MD (3), Chin B Eap, PhD (1, 14) and the Swiss Transplant Cohort Study.
12

13 ¹ Unit of Pharmacogenetics and Clinical Psychopharmacology, Department of Psychiatry,
14 Lausanne University Hospital, Prilly, Switzerland.

15 ² Service of Infectious Diseases, Lausanne University Hospital, Lausanne, Switzerland

16 ³ Transplant Center, Lausanne University Hospital, Lausanne, Switzerland.

17 ⁴ Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital,
18 Lausanne, Switzerland.

19 ⁵ Swiss Institute of Bioinformatics, Lausanne, Switzerland.

20 ⁶ Service of Endocrinology and Diabetes, University Hospital, Zurich, Switzerland.

21 ⁷ Division of Infectious Diseases and Hospital Epidemiology, University Hospital, Zurich,
22 Switzerland.

23 ⁸ Service of Nephrology and Transplantation Medicine, Kantonsspital, St Gallen, Switzerland.

24 ⁹ Service of Infectious Diseases, University Hospitals, Geneva, Switzerland.

25 ¹⁰ Service of Nephrology, University Hospital, Basel, Switzerland.

26 ¹¹ Swiss Cardiovascular Center Bern, University Hospital, Bern, Switzerland.

27 ¹² Department of Clinical Pharmacology, University Hospital, Bern, Switzerland.

28 ¹³ Service of Pulmonary Medicine, University Hospital, Geneva, Switzerland.

29 ¹⁴ School of Pharmaceutical Sciences, University of Geneva, University of Lausanne, Geneva,
30 Switzerland.

31

32

33

34

35 **For correspondence:**

36 Prof CB. Eap

37 Hôpital de Cery,

38 1008 Prilly – Lausanne, Switzerland

39 Tel: 0041 21 314 26 04

Fax: 0041 21 314 24 44

Email: chin.eap@chuv.ch

40

41

42

43

44

45

46

47

48

49

50 **ABSTRACT**

51 Metabolic syndrome after transplantation is a major concern following solid organ transplantation
52 (SOT). The CREB-regulated transcription coactivator 2 (CRTC2) regulates glucose metabolism. The
53 effect of *CRTC2* polymorphisms on new-onset diabetes after transplantation (NODAT) was
54 investigated in a discovery sample of SOT recipients ($n_1=197$). Positive results were tested for
55 replication in two samples from the Swiss Transplant Cohort Study (STCS, $n_2=1294$ and $n_3=759$).
56 Obesity and other metabolic traits were also tested. Associations with metabolic traits in population-
57 based samples ($n_4=46'186$, $n_5=123'865$, $n_6>100,000$) were finally analyzed. In the discovery sample,
58 *CRTC2 rs8450-AA* genotype was associated with NODAT, fasting blood glucose and BMI
59 ($p_{\text{corrected}}<0.05$). *CRTC2 rs8450-AA* genotype was associated with NODAT in the second STCS
60 replication sample (OR=2.01, $p=0.04$). In the combined STCS replication samples, the effect of
61 *rs8450-AA* genotype on NODAT was observed in patients having received SOT from a deceased
62 donor and treated with tacrolimus ($n=395$, OR=2.08, $p=0.02$) and in non-kidney transplant recipients
63 (OR=2.09, $p=0.02$). Moreover, *rs8450-AA* genotype was associated with overweight or obesity
64 ($n=1215$, OR=1.56, $p=0.02$), new-onset hyperlipidemia ($n=1007$, OR=1.76, $p=0.007$), and lower
65 HDL-cholesterol ($n=1214$, $\beta=-0.08$, $p=0.001$). In the population-based samples, a proxy of
66 *rs8450G>A* was significantly associated with several metabolic abnormalities. *CRTC2 rs8450G>A*
67 appears to play an important role in the high prevalence of metabolic traits observed in patients with
68 SOT. A weak association with metabolic traits was also observed in the population-based samples.

69

70

71 **Key terms:** Transplantation, metabolic syndrome after transplantation, genetic polymorphisms.

72

73

74

75 **Introduction**

76 The introduction of calcineurin inhibitors (CNIs) - cyclosporine (CSA) and tacrolimus (TAC) - in
77 solid organ transplantation (SOT) has reduced the incidence of acute rejection episodes and improved
78 short-term graft survival^{1,2}. However, both drugs are not devoid of metabolic complications, such as
79 glucose intolerance, hypertension, and hyperlipidemia which can be very pronounced and have a
80 detrimental impact on patients' life quality, and increase the mortality risk due to cardiovascular
81 events³. Indeed, cardiovascular disease is responsible for approximately 20-40% of non-graft-related
82 death after the first post-transplantation year⁴⁻⁶.

83 New onset diabetes mellitus after transplantation (NODAT) is a serious complication partly related to
84 the use of CNIs, mainly TAC^{7,8}. It is also associated with increased cardiovascular events, infectious
85 complications, and graft loss^{9,10}. There are several risk factors which increase the risk of NODAT in
86 transplantation such as obesity, increased age, male sex, deceased donor, hepatitis C (HCV) status,
87 acute rejection, and African-American or Hispanic descent^{9,11}. The identification of genetic factors
88 involved in the development of NODAT may greatly benefit from a more detailed understanding of
89 the complex metabolic pathways involved in glucose metabolism. Genome-wide association studies
90 (GWAS) conducted to date explain only 10% of type 2 diabetes heritability and more diabetic
91 susceptibility genes remain to be discovered (reviewed in¹² and¹³). A recent GWAS investigating
92 NODAT identified eight polymorphisms¹⁴. However, only one of these was replicated in another
93 study¹⁵. Whereas GWAS have been extremely valuable, other approaches are needed to further
94 understand the pathophysiology of NODAT.

95 The cAMP-regulated transcriptional coactivator 2 (CRTC2), a transcriptional coactivator that
96 promotes the transcription of genes targeted by the cAMP response element-binding protein¹⁶, is an
97 interesting target protein in glucose metabolism. CRTC2 belongs to the CRTC family which
98 comprises 2 other members, CRTC1 and CRTC3¹⁷. CRTC1 is mainly expressed in the central nervous
99 system and we recently showed an association between *CRTC1* polymorphisms and obesity markers
100 (body mass index (BMI) and fat mass) in psychiatric and population-based samples¹⁸, CRTC2 is

101 highly expressed in the thymus, and present in both T and B-lymphocytes¹⁷. It is also expressed in the
102 liver and it plays a direct role during the fasting state in the induction of gluconeogenic genes¹⁶. It
103 further enhances hepatic insulin signalling by stimulating expression of the *insulin receptor substrate*
104 *2* gene, thus triggering a feedback response that limits glucose output from the liver during fasting¹⁹,
105 Calcineurin, the target of both TAC and CSA, plays an important role in the activation of *CRTC2*^{20, 21}.
106 Overexpression of *CRTC2* induced by a mutation at 2 regulatory sites rendered *CRTC2* constitutively
107 active in an animal model and permitted *CRTC2*-target gene activation even when calcineurin was
108 inhibited by CNIs²². The authors of this study also showed that *CRTC2* is required for β -cell function
109 and proliferation and promoting this pathway could ameliorate symptoms of NODAT²².
110 *CRTC2* single nucleotide polymorphisms (SNPs) were previously investigated in two Asian
111 populations and one coding SNP (*R379C*), with a very low minor allele frequency (MAF), was
112 associated with type II diabetes²³ and with lung cancer²⁴. So far *CRTC2* SNPs have not been
113 investigated in other ethnic populations or with other phenotypes. In this work, we aimed to study the
114 influence of *CRTC2* SNPs on the incidence of NODAT in a sample of Caucasian SOT recipients, and
115 positive results were then tested for replication in the Swiss transplant cohort Study (STCS). We also
116 aimed to extend our analysis to obesity and other variables of the metabolic syndrome (MetS)
117 following SOT. Finally, we aimed to test if the associations with MetS components could be found in
118 general population-based samples (>100,000 subjects).

119

120 **Materials and Methods**

121 **Discovery study sample:**

122 This study aimed initially to investigate the effect of genetic polymorphisms of drug metabolizing
123 enzymes and/or transporters on the incidence of different post-transplant complications and on
124 immunosuppressive doses and blood levels^{25, 26}. For this study, a total of 197 patients were enrolled
125 between 2003 and 2005 from the outpatient clinic of the Transplant Center of the University Hospital
126 of Lausanne, Switzerland. Patients with functional graft for more than 12 months after transplantation
127 were eligible to participate in the study. Data regarding patient's age, gender, BMI, ethnic origin,
128 donor's age, HLA mismatch, duration of graft cold ischemia and delayed graft function were collected

129 retrospectively from patients' medical files. Immunosuppressive regimens, doses and blood levels
130 were obtained retrospectively at different time-points during the first year post-transplantation. The
131 study was approved by the ethics committee of the University of Lausanne. Patients gave their written
132 informed consent to participate in this pharmacogenetic study. Once included, venous blood samples
133 were collected for DNA extraction and further genotyping analysis.

134 For the present study, data were collected between October 2011 and April 2012 as related to the
135 development of NODAT during the first 5 years following transplantation. Data regarding fasting
136 blood glucose (FBG), glycated haemoglobin (HbA1c), 2hrs oral glucose tolerance test (OGTT),
137 insulin and oral anti-diabetic treatment were collected retrospectively from the medical files at the
138 time of transplantation, at 1, 3, 6, 9 and 12 month post-transplantation and at the yearly follow-up until
139 5-year post-transplantation. NODAT was diagnosed if a patient needed anti-diabetic treatment (either
140 insulin or oral anti-diabetic agents) for at least 6 months following transplantation or had several
141 abnormal glucose profiles during the follow-up period that fulfil the criteria given by the WHO and
142 ADA consensuses²⁷, including $FBG \geq 7.0$ mmol/l (in \geq two occasions) or 2 hours plasma glucose ≥ 11.1
143 mmol/l during OGTT²⁷ (even if not treated with anti-diabetic drugs). Patients who had diabetes or
144 prediabetes before transplantation or were transplanted because of diabetic nephropathy were excluded
145 from the present study.

146

147 **The Swiss Transplant Cohort Study (STCS):**

148 The STCS is an ongoing prospective multicenter cohort project (Basel, Bern, Genève, Lausanne, St.
149 Gallen and Zurich), aiming at a nationwide comprehensive and structured data collection in all SOT
150 recipients. All recipients of SOTs in Switzerland are prospectively registered since May 2008.
151 Currently more than 3500 patients are included in the STCS. No particular eligibility or exclusion
152 criteria exist for enrolment. After transplantation, all patients are mandatorily followed in their
153 respective transplant centers. After baseline assessment, STCS follow-up assessments take place at 6-,
154 12 months and yearly thereafter. Biological samples are collected in relation to the case at baseline,
155 and at 6-, and the 12-month visits. Data regarding patient's age, gender, BMI, blood pressure and lipid
156 profiles (total cholesterol, HDL- and LDL-cholesterol) were available for all patients at the different

157 time-points of the STCS follow-up. Full description of this cohort is published elsewhere^{28, 29}. For the
158 first STCS replication sample, patients transplanted from May 2008 to 8th of May 2011 with a
159 functional graft for at least 12 months after transplantation were included in the analyses (n=1294).
160 NODAT was diagnosed if a patient needed anti-diabetic treatments following transplantation or if
161 such new metabolic event was reported in their case report forms. For the second STCS replication
162 sample, patients transplanted from 9th of May 2011 to May 2013 with a functional graft for at least 12
163 months after transplantation were included in the analyses (n=759). For this second STCS replication
164 sample, several random and FBG were available in the database, as well as HbA1c. Therefore, as for
165 the discovery sample, NODAT was diagnosed if a patient needed anti-diabetic treatment (either
166 insulin or oral anti-diabetic agents) or fulfilled the criteria given by the WHO and ADA consensus
167 mentioned earlier. For both STCS samples, patients who were diagnosed with diabetes or were
168 prediabetics before transplantation or were transplanted because of diabetic nephropathy were
169 excluded from the NODAT analyses.

170 For the combined STCS sample, new-onset hypertension and new-onset hyperlipidemia were
171 diagnosed if patients needed anti-hypertensive and hypolipidemic treatment post-SOT. Patients with
172 previous hypertension or hyperlipidemia were excluded from the new-onset hypertension and new-
173 onset hyperlipidemia analyses, respectively. The study was approved by the ethics committee of their
174 respective centers. Patients gave their written informed consent to participate in this pharmacogenetic
175 study.

176 Recipients younger than 18 years old and recipients with multiple organ transplantation were excluded
177 from the whole analyses. If a patient received more than one transplantation during the inclusion
178 period, only data from the first SOT was included in the analyses.

179 For both the discovery and STCS samples, as abdominal obesity (waist circumference) was not
180 available, BMI was used as a marker of obesity.

181

182 **Population based samples:**

183 We aimed to replicate results with MetS traits in several population-based samples: the Meta-Analyses
184 of Glucose and Insulin-related traits Consortium (MAGIC)(n=46'186)³⁰, Genetic Investigation of

185 ANthropometric Traits (GIANT) consortium (n=123'865)³¹⁻³³, and Genome Wide Associations Scans
186 for Total Cholesterol, HDL-C, LDL-C and triglycerides (n>100'000)^{34, 35}. More details are in the
187 Supplementary Material.

188

189 **Polymorphism selection and genotyping:**

190 Two tagging SNPs within *CRTC2* gene (*rs8450G>A* and *rs12117078G>C*) were selected using
191 HapMap Genome Browser by limiting the search to SNPs with a MAF>5% in the Caucasian
192 population and r^2 cutoff of 0.8. Genotyping in the discovery study sample was done using Taqman
193 allelic discrimination assays. Genetic analyses for the STCS replication samples were done using the
194 fluorescence-based competitive allele-specific PCR technology (KASPar)³⁶. More details are in the
195 Supplementary Material.

196

197 **Statistical analysis:**

198 Detailed statistical methods are presented in the Supplementary Material.

199 *Discovery and STCS samples:*

200 We assessed the association of *CRTC2* SNPs with MetS traits 1) by applying a logistic regression for
201 binary outcomes adjusted for recipient's age at transplantation and sex. Other variables identified
202 through the univariate analysis (P<0.10) were also added as covariates in these models, 2) by using
203 linear mixed models for continuous variables, and 3) by fitting a Generalized Additive Mixed Model
204 (GAMM) to include a smooth trend for the response in time (allowing multiple observations for each
205 patient) for non-linear continuous variables (BMI and FBG).

206 For the discovery sample, the p-values of the models were adjusted for multiple comparisons using
207 Bonferroni correction and the SNP that survived this correction was investigated in the STCS.

208 Data for both the discovery and replications samples were analyzed using Stata 12 (StataCorp, College
209 Station TX, USA) and R version 2.13.0 software (<http://www.R-project.org>).

210 *Population-based samples:*

211 We analyzed the associations of *CRTC2* SNP with different MetS traits using multivariate linear
212 regression with allele dosage in which potential confounding factors such as age, sex, and smoking
213 status were added as covariates.

214

215 **Results:**

216 **Discovery and STCS replication samples:**

217 General characteristics of the discovery and STCS replication samples are presented in the
218 Supplementary Material and Supplementary Tables 1 and 2. Supplementary Table 2 represents the
219 global STCS sample without the exclusion of patients with previous history of diabetes or other
220 metabolic abnormalities before transplantation.

221 The distribution of *CRTC2* genotypes in all studies did not deviate from the Hardy-Weinberg
222 equilibrium ($P>0.05$) and the MAFs were similar to those reported in HapMap (Supplementary Table
223 3).

224

225 ***CRTC2* SNPs and NODAT, FBG, and BMI in the discovery study sample:**

226 Carriers of the *CRTC2 rs8450-AA* genotype showed increased risk of NODAT as compared to wild-
227 type *GG* genotype (odd ratio (OR)=6.91, 95% CI:1.52-31.36, $p_{\text{corrected}}=0.02$). An increased risk was
228 also observed by analyzing the *rs8450* SNP in a recessive model (Table 1). Neither cumulative
229 prednisone dosages nor calcineurin inhibitors' doses, trough levels or concentration/dose ratios tested
230 at 2 time periods (at 1 month or at 12 months post-SOT) were associated with NODAT in the
231 univariate analysis or when used in the multivariate model including *CRTC2* SNPs and the other
232 covariables (data not shown).

233 FBG were also available at several time-points after SOT. By applying the GAMM model, increased
234 FBG levels were found in patients with the *CRTC2 rs8450-GA and AA* genotypes (0.24 mmol/L,
235 $p_{\text{corrected}}=0.06$ and 0.47 mmol/L, $p_{\text{corrected}}=0.02$, respectively) as compared to wild-type *GG* genotype.
236 We observed an equal increased of FBG levels by analyzing the *rs8450* SNP in dominant model
237 (Table 1).

238 Patients with the *CRTC2 rs8450-GA* genotype showed a 1.36 kg/m² increased BMI compared to the
239 wild-type genotype ($p_{\text{corrected}}=0.02$). By analyzing the SNP in a dominant model, we observed a
240 significant association between *CRTC2 rs8450* and BMI ($\beta=1.17$, 95%CI:0.17-2.35, $p_{\text{corrected}}=0.04$).
241 We observed no association between *CRTC2 rs12117078G>C* SNP and NODAT, FBG or BMI (Table
242 1).

243 ***CRTC2* SNP in the STCS replication samples:**

244 The *CRTC2 rs8450G>A* SNP that survived corrections for multiple testing in the discovery study
245 sample was investigated in the STCS. Recessive models for this SNP are presented in the text and in
246 Tables 2 and 3 while additive and dominant models regarding NODAT are presented in
247 Supplementary Table 4.

248

249 ***CRTC2* SNP and NODAT:**

250 In the univariate analyses, increased recipient's age ($p=0.0003$), male sex ($p=0.03$), positive HCV
251 status ($p=0.003$), deceased donor ($p=0.04$), treatment with TAC ($p=0.001$) and baseline BMI
252 ($p=0.0002$) were among the non-genetic factors associated with an increased risk of NODAT and were
253 used as covariates in the genetic models. In the first STCS replication samples, we observed a non-
254 significant association between *rs8450G>A* and NODAT (Table 2). However, by analyzing the SNP
255 in the subgroup of patients having received SOT from a deceased donor and treated with TAC, a
256 significant effect of the SNP was observed ($n=281$, OR=2.36, 95%CI:1.16-4.78, $p=0.01$)(Table 2). In
257 the second replication sample ($n=438$), in which the diagnostic criteria for NODAT was the same as
258 for the discovery sample, recipients carrying the *rs8450-AA* genotype showed a significant increased
259 risk of NODAT ($n=438$, OR=2.01, 95%CI:1.03-3.91, $p=0.04$). In the combined STCS sample, the
260 effect of the *CRTC2* genotype was only observed in the subgroup of patients having received SOT
261 from a deceased donor and treated with TAC ($n=395$, OR=2.08, 95%CI:1.15-3.74, $p=0.02$) (Table 2).

262

263

264

265 ***CRTC2* SNP and BMI, new-onset hypertension/ hyperlipidemia:**

266 We investigated the influence of *CRTC2 rs8450G>A* on BMI and on the incidence of new-onset
267 hypertension/hyperlipidemia in the combined STCS sample.

268 Increased recipient's age ($p<0.0001$), male sex ($p<0.0001$) and type of transplanted organ ($p<0.0001$)
269 were among the non-genetic factors associated with increased BMI in the univariate analyses and were
270 used as covariates in the genetic models. BMI values were also dichotomized into normal versus
271 overweight or obese ($BMI\geq 25$ kg/m²). At 12-months post-SOT, overweight or obese patients were
272 more prevalent in the *CRTC2 rs8450-AA* genotype compared to the other genotypes ($n=1215$,
273 $OR=1.56$, $95\%CI:1.08-2.25$, $p=0.02$). We observed the same association but with lower significance
274 for the combined STCS sample at 6-months post-SOT ($n=1389$, $OR=1.41$, $95\%CI:1.01-1.97$, $p=0.04$),
275 and a non-significant association with baseline BMI ($n=1515$, $OR=1.20$, $95\%CI:0.87-1.65$, $p=0.27$),
276 suggesting that time post-SOT modulates the effect of the *CRTC2* genotypes.

277 By applying the GAMM model to test the association between BMI over time and *CRTC2 rs8450G>A*
278 SNP ($n=1625$), patients with the *CRTC2 rs8450-AA* genotype showed a 0.47 kg/m² ($95\%CI:0.11-1.02$,
279 $p=0.01$) increase in BMI compared to the other genotypes.

280 We found no association between the *CRTC2 rs8450G>A* and the incidence of new-onset
281 hypertension (Supplementary Material). On the other hand, carriers of the *CRTC2 rs8450-AA*
282 genotype showed increased risk of new-onset hyperlipidemia as compared to the other genotypes
283 ($n=1007$, $OR=1.76$, $95\%CI:1.16-2.66$, $p=0.007$). Additionally, patients with the *CRTC2 rs8450-AA*
284 genotype also showed a significant decrease of HDL-Cholesterol levels in the combined STCS sample
285 ($n=1214$, $\beta= -0.08$ mmol/L, $p=0.001$). *CRTC2 rs8450G>A* SNP was not associated with total
286 cholesterol or with LDL-cholesterol blood levels in the combined STCS sample (data not shown).
287 More details can be found in the Supplementary Material.

288

289 ***CRTC2* SNP and MetS traits in kidney and non-kidney transplant recipients:**

290 Patients were dichotomized into kidney versus non-kidney transplant recipients (Table 3). The *CRTC2*
291 *rs8450-AA* genotype was associated with significantly higher OR for NODAT in non-kidney

292 transplant recipients (n=409, OR=2.09, 95%CI:1.13-3.86, p=0.02), while the *rs8450 AA*-genotype was
293 only associated with higher risk of new-onset hyperlipidemia in kidney transplant recipients (n=573,
294 OR=1.95, 95%CI:1.14-3.15, p=0.01).

295

296 **Population-based samples:**

297 Several glyceic traits were available in the MAGIC study (Supplementary Table 5); each *rs1572788*
298 *C*-allele (in complete linkage disequilibrium (LD) with *rs8450 A*-allele) increased FBG levels by 0.01
299 mmol/l (n=46'186, p=0.004). Additionally, each *rs1572788 C*-allele decreased HOMA-B by 0.008%
300 (n=46'186, p=0.03). No association was observed between *rs1572788T>C* and HOMA-IR or the 2
301 hour OGTT.

302 Decreased HDL-cholesterol and increased triglycerides were observed for each *rs1572788 C*-allele in
303 the Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides
304 (n=96'908, β =-0.008, p=0.013 and n=93'562, β =0.009, p=0.004, respectively). We observed non-
305 significant associations for *rs1572788T>C* with total and LDL-cholesterol, as well as with obesity
306 traits studied in the GIANT study, however, the direction of the effect is consistent with increased
307 blood lipids and obesity with each *rs1572788 C*-allele (Supplementary Table 5).

308 **Discussions:**

309 The transcriptional co-activator (CRTC) family is implicated with energy expenditure (CRTC1 and
310 CRTC3)^{37,38} and glucose metabolism (CRTC2)¹⁶ in animal models. In the present study, we aimed to
311 study the effect of *CRTC2* SNPs on the development of diabetes in SOT recipients, a population at
312 high risk of developing metabolic abnormalities, and in population-based samples. We also extended
313 our analyses to investigate obesity and other metabolic traits. The *CRTC2 rs8450-AA* genotype was
314 associated with increased risk of NODAT, increased FBG, and BMI in a sample of 156 patients with
315 SOT. Results on NODAT were replicated only in the second STCS replication sample (n=438), while
316 in the first and the combined STCS replication samples, the SNP only showed significant influence in
317 the population at higher risk of developing NODAT, mainly in recipients having received a graft from
318 a deceased donor and treated with TAC. We also observed a significant association for BMI in the

319 STCS replication sample; overweight and obesity were also more frequent in patients carrying the
320 *rs8450-AA* genotype at 12 months post-SOT. *CRTC2 rs8450G>A* was also associated with increased
321 risk of new-onset hyperlipidemia following SOT and with lower HDL-cholesterol blood levels.
322 Furthermore, we also found the increased risk of the *C-allele* of *rs1572788* (in complete LD with
323 *rs8450 A-allele*) on diabetic and lipidemic traits in two large population-based samples (MAGIC and
324 GWAS for Total Cholesterol, HDL-C, LDL-C and triglycerides, respectively), and a consistent
325 direction even though non-significant between *rs1572788T>C* and obesity traits in the GIANT study.

326 *CRTC2 rs8450G>A* was associated with NODAT in the total discovery sample and only in the second
327 STCS replication sample, but not in the total STCS sample in which a significant association was
328 observed only in the subgroup of patients at higher risk of developing NODAT. This could partly be
329 explained by the difference of NODAT diagnostic criteria, which is only based on anti-diabetic
330 treatment in the first STCS replication sample. Another explanation could be the high percentage of
331 patients receiving an organ from a living donor in the first STCS replication sample (28%,
332 Supplementary Table 2) compared to only 11.5% in the discovery sample (Supplementary Table 1).
333 Deceased donor is a common risk factor of NODAT¹¹ and the impact of the SNP seems to be more
334 prominent in the groups at higher risk of developing MetS traits. Additionally, when differentiating
335 kidney and non-kidney transplant recipients (Table 3), we observed a significant association between
336 the *CRTC2 rs8450-AA* genotype and NODAT in non-kidney transplant recipients. The percentage of
337 non-kidney transplant recipients was higher in the second STCS replication sample compared to the
338 first STCS sample (Supplementary Table 2) and it should be noted that kidney transplant recipients
339 had higher percentages of living donors compared to non-kidney transplant recipients (43% vs. 2%
340 respectively). Moreover, kidney transplant recipients usually have obesity and other metabolic
341 problems already before transplantation compared to other solid organ recipients, which could explain
342 the differential effect of the SNP depending on the type of the transplanted organ. Interestingly, in
343 both the discovery and the combined STCS samples, the effect of the SNP was stronger in NODAT
344 cases developed after the first year post-SOT (data not shown), It maybe that the effect of the *CRTC2*
345 SNP on NODAT is more important in the long-term, but this remains to be further studied.

346 In the GWAS literature, *CRTC2* SNPs were not found to be associated with diabetes, obesity or
347 hyperlipidemia. On the other hand, by testing the individual effect of *CRTC2 rs1572788T>C* SNP on
348 diabetic and lipidemic traits in two population-based samples, a significant effect was observed, even
349 though this effect is very weak and the variance explained by the SNP is very small (Supplementary
350 Table 5) and could thus not be detected by GWAS. However, patients with SOT is a population at risk
351 of developing MetS traits, because of the disease itself and/or the immunosuppressive medications,
352 both can act as an important trigger, unmasking different genetic factors. So in this at risk population,
353 the effect of *CRTC2* SNP was more pronounced, even in much smaller sample sizes compared to
354 population-based samples. Additionally, in the GWAS of NODAT¹⁴, *CRTC2* SNPs did not reach
355 genome-wide level of significance. Moreover, most of the SNPs retained in this GWAS were not
356 replicated in another study with NODAT¹⁵. In fact, both GWAS and candidate gene studies are
357 important gene association approaches and both are subject to the same artifacts of spurious
358 association findings. Depending on the methodology of these studies (sample size, definition of the
359 phenotypes, MAF of the SNPs, corrections for multiple testing, etc...), discordances between both
360 approaches could be seen. GWAS relies on indirect association to locate a disease-causing variant
361 while candidate gene approach relies on an a priori hypothesis to identify this disease-causing variant.
362 For all of these reasons, replication of positive finding in either approach is of utmost importance to
363 validate the results.

364 The functional activity of this SNP, in the regulatory 3'UTR, is unknown and data from 1000
365 Genomes³⁹ showed that 78 SNPs in the *CRTC2* gene region are in high LD (r^2 threshold ≥ 0.8) with
366 *rs8450G>A*. We used the RegulomeDB database that annotates SNPs with known and predicted
367 regulatory elements in the intergenic regions of the *H. sapiens* genome⁴⁰. This database reveals known
368 and predicted regulatory DNA elements including regions of DNAase hypersensitivity, binding sites
369 of transcription factors, and promoter regions that have been biochemically characterized to regulation
370 transcription. Source of these data include public datasets from GEO, the ENCODE project, and
371 published literature. Regarding *CRTC2 rs8450G>A*, several of these proxy SNPs (e.g. *rs6680140* with
372 $r^2=1$) show cis-eQTL effect on the expression of *CRTC2* and fall into the autoimmune regulatory

373 (AIRE) motif, influence the binding of several proteins (*JUN*, *CREBBP* and *ELF1*), and has several
374 histone marks (H3k09me3, H4k20me1, etc.). Altogether, these data suggest that the SNP might have a
375 regulatory function.

376 Interestingly, among the analyzed clinical factors, treatment with corticosteroids was not associated
377 with NODAT, neither in the discovery cohort nor in the STCS sample. However, in our samples most
378 of the patients were treated with corticosteroids so it is difficult to draw major conclusions. On the
379 other hand, the cumulative corticosteroids dosages and the duration of therapy rather than the simple
380 corticosteroids administration could be the most influencing factors. In our discovery sample, the
381 cumulative dosages of corticosteroids at the first year post-transplantation were not associated with
382 NODAT outcome. In the literature, several studies did not find any influence of cumulative
383 corticosteroid dosages on the appearance of NODAT⁴¹⁻⁴⁵. The current lower dosages of corticosteroids
384 used in different immunosuppressive regimens could contribute to the negative association with
385 NODAT appearance in our study.

386 This study has several limitations and strengths. Random and FBG levels were only available in the
387 second STCS replication sample and the diagnosis of NODAT in the first STCS replication sample
388 was only based on anti-diabetic treatments post-SOT. The genotype of the *rs8450G>A* SNP of the
389 liver transplant donors is unknown, therefore the significant effect of the SNP on NODAT found in
390 non-kidney transplant recipients, in which nearly half of them are liver transplant recipients, should be
391 interpreted with caution. However, *CRTC2* is expressed in different tissues and is implicated in
392 glucose regulation through different mechanisms (gluconeogenesis, insulin signaling and beta-cell
393 survival^{16, 19, 22}) which could highlight the importance of recipient *CRTC2* genotype even in liver
394 transplant recipients. Additionally, by excluding patients with liver transplantation from the non-
395 kidney transplant recipients' analyses, a significant association between the *CRTC2* SNP and NODAT
396 is still observed (data not shown). The present results do not allow the determination of whether the
397 *rs8450G>A* SNP is the causative variant or merely a proxy of one or more yet unidentified variants.
398 Despite the possible regulatory functions proposed by the RegulomeDB database, further studies are
399 needed to elucidate which precise mechanisms underlie the observed association. This study included

400 people of Caucasian origin and results cannot be generalized to other ethnic groups. On the other hand,
401 the fact that the results were replicated in two independent samples with SOT and in large population-
402 based samples, the latter used as a proof of concept of the effect of the polymorphism, strengthens the
403 validity of our data.

404 In conclusion, our results suggest that variations in *CRTC2* play an important role in the high
405 prevalence of MetS complications observed in patients with SOT. Besides, by studying *CRTC2* SNPs
406 in population-based samples, we still observed a weak yet significant association, thereby suggesting
407 that the effect of *CRTC2* variations is more important in population at risk of developing different
408 MetS traits than in the general population. This is the first study showing the importance of *CRTC2*
409 variations with NODAT and other MetS traits in patients with SOT. The assessment of the major risk
410 gene variants would allow predicting vulnerability for developing Mets phenotypes and thus adapt the
411 immunosuppressive treatment from the beginning by genotyping the patients before transplantation. In
412 the long-term, these patients could benefit from individualized immunosuppressive regimens adapted
413 to their genetics and environment.

414

415

416

417

418

419

420

421

422

423

424

425

426

427 **Funding**

428 This work has been funded in part by the Swiss National Science Foundation (CBE: 324730_144064).
429 LQ and CBE received research support from the Roche Organ Transplantation Research Foundation
430 (#152358701) and the Swiss Transplant Cohort Study in the past 3 years. ZK was funded by the Swiss
431 National Science Foundation (31003A-143914) and the Leenaards Foundation.

432 This study has been conducted in the framework of the Swiss Transplant Cohort Study, supported by
433 the Swiss National Science Foundation and the Swiss University Hospitals (G15) and transplant
434 centers.

435

436 **Acknowledgements**

437 Data on glycaemic traits have been contributed by MAGIC investigators and have been downloaded
438 from www.magicinvestigators.org

439 CBE takes full responsibility for the work as a whole, including the study design, access to data, and
440 the decision to submit and publish the manuscript.

441

442 **Declaration of interests:**

443 CBE received honoraria for conferences or teaching CME courses from Advisis, Astra Zeneca,
444 Lundbeck, MSD, Sandoz, Servier and Vifor-Pharma in the past 3 years. He received an unrestricted
445 educational grant from Takeda in the past 3 years.

446 JFD Advisory committees: Bayer, BMS, Gilead Science, Janssen Cilag, Jennerex, Merck, Novartis,
447 Roche. Speaking and teaching: Bayer, Boehringer-Ingelheim, Novartis, Roche

448 SC received honoraria for teaching CME courses from Astra Zeneca and Lundbeck.

449 The authors of this manuscript have no conflicts of interest to disclose.

450

451 **Footnote**

452 The members of the Swiss Transplant Cohort Study are: Rita Achermann, John-David Aubert,
453 Philippe Baumann, Guido Beldi, Christian Benden, Christoph Berger, Isabelle Binet, Pierre-Yves
454 Bochud, Elsa Boely (Head of local data management), Heiner Bucher, Leo Bühler, Thierry Carell,
455 Emmanuelle Catana, Yves Chalandon, Sabina de Geest, Olivier de Rougemont, Michael Dickenmann,
456 Michel Duchosal, Thomas Fehr, Sylvie Ferrari-Lacraz, Christian Garzoni, Yvan Gasche, Paola Gasche
457 Socal, Emiliano Giostra, Déla Golshayan, Daniel Good, Karine Hadaya, Christoph Hess, Sven
458 Hillinger, Hans Hirsch, Günther Hofbauer, Uyen Huynh-Do, Franz Immer, Richard Klaghofer,
459 Michael Koller (Head of the data center), Thomas Kuntzen, Bettina Laesser, Roger Lehmann,
460 Christian Lovis, Oriol Manuel, Hans-Peter Marti, Pierre Yves Martin, Pascal Meylan (Head,
461 Biological samples management group), Paul Mohacsi, Isabelle Morard, Philippe Morel, Ulrike
462 Mueller, Nicolas Mueller (Chairman Scientific Committee), Helen Mueller-McKenna, Thomas
463 Müller, Beat Müllhaupt, David Nadal, Gayathri Nair, Manuel Pascual (Executive office), Jakob
464 Passweg, Chantal Piot Ziegler, Juliane Rick, Eddy Roosnek, Anne Rosselet, Silvia Rothlin, Frank
465 Ruschitzka, Urs Schanz, Stefan Schaub, Christian Seiler, Nasser Semmo, Susanne Stampf, Jürg
466 Steiger (Head, Executive Office), Christian Toso, Dimitri Tsinalis, Christian Van Delden (Executive
467 office), Jean-Pierre Venetz, Jean Villard, Madeleine Wick (STCS coordinator), Markus Wilhelm,
468 Patrick Yerly.

469

470 **Supplementary data:**

471 Supplementary information is available at journal's website

472

473

474

475

476 **References:**

- 477 1. Mayer AD, Dmitrewski J, Squifflet JP, Besse T, Grabensee B, Klein B, *et al.* Multicenter
478 randomized trial comparing tacrolimus (FK506) and cyclosporine in the prevention of renal
479 allograft rejection: a report of the European Tacrolimus Multicenter Renal Study Group.
480 *Transplantation* 1997; **64**(3): 436-443.
- 481
482 2. Hariharan S, Johnson CP, Bresnahan BA, Taranto SE, McIntosh MJ, Stablein D. Improved graft
483 survival after renal transplantation in the United States, 1988 to 1996. *The New England*
484 *journal of medicine* 2000; **342**(9): 605-612.
- 485
486 3. Ojo AO, Hanson JA, Wolfe RA, Leichtman AB, Agodoa LY, Port FK. Long-term survival in renal
487 transplant recipients with graft function. *Kidney international* 2000; **57**(1): 307-313.
- 488
489 4. Collins AJ, Foley RN, Herzog C, Chavers B, Gilbertson D, Ishani A, *et al.* United States Renal
490 Data System 2008 Annual Data Report. *American journal of kidney diseases : the official*
491 *journal of the National Kidney Foundation* 2009; **53**(1 Suppl): S1-374.
- 492
493 5. Johnston SD, Morris JK, Cramb R, Gunson BK, Neuberger J. Cardiovascular morbidity and
494 mortality after orthotopic liver transplantation. *Transplantation* 2002; **73**(6): 901-906.
- 495
496 6. Vogt DP, Henderson JM, Carey WD, Barnes D. The long-term survival and causes of death in
497 patients who survive at least 1 year after liver transplantation. *Surgery* 2002; **132**(4): 775-
498 780; discussion 780.
- 499
500 7. Opelz G, Dohler B, Collaborative Transplant S. Influence of immunosuppressive regimens on
501 graft survival and secondary outcomes after kidney transplantation. *Transplantation* 2009;
502 **87**(6): 795-802.
- 503
504 8. First MR, Dhadda S, Croy R, Holman J, Fitzsimmons WE. New-onset diabetes after
505 transplantation (NODAT): an evaluation of definitions in clinical trials. *Transplantation* 2013;
506 **96**(1): 58-64.
- 507
508 9. Kasiske BL, Snyder JJ, Gilbertson D, Matas AJ. Diabetes mellitus after kidney transplantation
509 in the United States. *American journal of transplantation : official journal of the American*
510 *Society of Transplantation and the American Society of Transplant Surgeons* 2003; **3**(2): 178-
511 185.
- 512
513 10. Israni AK, Snyder JJ, Skeans MA, Kasiske BL, Investigators P. Clinical diagnosis of metabolic
514 syndrome: predicting new-onset diabetes, coronary heart disease, and allograft failure late
515 after kidney transplant. *Transplant international : official journal of the European Society for*
516 *Organ Transplantation* 2012; **25**(7): 748-757.

517

- 518 11. Pham PT, Pham PM, Pham SV, Pham PA, Pham PC. New onset diabetes after transplantation
519 (NODAT): an overview. *Diabetes, metabolic syndrome and obesity : targets and therapy*
520 2011; **4**: 175-186.
- 521
522 12. Billings LK, Florez JC. The genetics of type 2 diabetes: what have we learned from GWAS? *Ann*
523 *N Y Acad Sci* 2010; **1212**: 59-77.
- 524
525 13. Torres JM, Cox NJ, Philipson LH. Genome wide association studies for diabetes: perspective
526 on results and challenges. *Pediatr Diabetes* 2013; **14**(2): 90-96.
- 527
528 14. McCaughan JA, McKnight AJ, Maxwell AP. Genetics of new-onset diabetes after
529 transplantation. *J Am Soc Nephrol* 2014; **25**(5): 1037-1049.
- 530
531 15. Chand S, Shabir S, Chan W, McCaughan JA, McKnight AJ, Maxwell AP, *et al.* beta cell
532 glucotoxic-associated single nucleotide polymorphisms in impaired glucose tolerance and
533 new-onset diabetes after transplantation. *Transplantation* 2014; **98**(3): e19-20.
- 534
535 16. Koo SH, Flechner L, Qi L, Zhang X, Sreaton RA, Jeffries S, *et al.* The CREB coactivator TORC2 is
536 a key regulator of fasting glucose metabolism. *Nature* 2005; **437**(7062): 1109-1111.
- 537
538 17. Conkright MD, Canettieri G, Sreaton R, Guzman E, Miraglia L, Hogenesch JB, *et al.* TORCs:
539 transducers of regulated CREB activity. *Mol Cell* 2003; **12**(2): 413-423.
- 540
541 18. Choong E, Quteineh L, Cardinaux JR, Gholam-Rezaee M, Vandenberghe F, Dobrinas M, *et al.*
542 Influence of CRT1 Polymorphisms on Body Mass Index and Fat Mass in Psychiatric Patients
543 and the General Adult Population. *JAMA psychiatry* 2013; **70**(10): 1011-1019.
- 544
545 19. Canettieri G, Koo SH, Berdeaux R, Heredia J, Hedrick S, Zhang X, *et al.* Dual role of the
546 coactivator TORC2 in modulating hepatic glucose output and insulin signaling. *Cell*
547 *metabolism* 2005; **2**(5): 331-338.
- 548
549 20. Sreaton RA, Conkright MD, Katoh Y, Best JL, Canettieri G, Jeffries S, *et al.* The CREB
550 coactivator TORC2 functions as a calcium- and cAMP-sensitive coincidence detector. *Cell*
551 2004; **119**(1): 61-74.
- 552
553 21. Bittinger MA, McWhinnie E, Meltzer J, Iourgenko V, Latario B, Liu X, *et al.* Activation of cAMP
554 response element-mediated gene expression by regulated nuclear transport of TORC
555 proteins. *Current biology : CB* 2004; **14**(23): 2156-2161.
- 556
557 22. Eberhard CE, Fu A, Reeks C, Sreaton RA. CRT2 is required for beta-cell function and
558 proliferation. *Endocrinology* 2013; **154**(7): 2308-2317.

559

- 560 23. Keshavarz P, Inoue H, Nakamura N, Yoshikawa T, Tanahashi T, Itakura M. Single nucleotide
561 polymorphisms in genes encoding LKB1 (STK11), TORC2 (CRTC2) and AMPK alpha2-subunit
562 (PRKAA2) and risk of type 2 diabetes. *Molecular genetics and metabolism* 2008; **93**(2): 200-
563 209.
- 564
565 24. He Y, Li Y, Qiu Z, Zhou B, Shi S, Zhang K, *et al.* Identification and validation of PROM1 and
566 CRTC2 mutations in lung cancer patients. *Molecular cancer* 2014; **13**(1): 19.
- 567
568 25. Crettol S, Venetz JP, Fontana M, Aubert JD, Pascual M, Eap CB. CYP3A7, CYP3A5, CYP3A4, and
569 ABCB1 genetic polymorphisms, cyclosporine concentration, and dose requirement in
570 transplant recipients. *Therapeutic drug monitoring* 2008; **30**(6): 689-699.
- 571
572 26. Crettol S, Venetz JP, Fontana M, Aubert JD, Ansermot N, Fathi M, *et al.* Influence of ABCB1
573 genetic polymorphisms on cyclosporine intracellular concentration in transplant recipients.
574 *Pharmacogenetics and genomics* 2008; **18**(4): 307-315.
- 575
576 27. Davidson J, Wilkinson A, Dantal J, Dotta F, Haller H, Hernandez D, *et al.* New-onset diabetes
577 after transplantation: 2003 International consensus guidelines. Proceedings of an
578 international expert panel meeting. Barcelona, Spain, 19 February 2003. *Transplantation*
579 2003; **75**(10 Suppl): SS3-24.
- 580
581 28. Koller MT, van Delden C, Muller NJ, Baumann P, Lovis C, Marti HP, *et al.* Design and
582 methodology of the Swiss Transplant Cohort Study (STCS): a comprehensive prospective
583 nationwide long-term follow-up cohort. *European journal of epidemiology* 2013; **28**(4): 347-
584 355.
- 585
586 29. Manuel O, Kralidis G, Mueller NJ, Hirsch HH, Garzoni C, van Delden C, *et al.* Impact of antiviral
587 preventive strategies on the incidence and outcomes of cytomegalovirus disease in solid
588 organ transplant recipients. *American journal of transplantation : official journal of the*
589 *American Society of Transplantation and the American Society of Transplant Surgeons* 2013;
590 **13**(9): 2402-2410.
- 591
592 30. Prokopenko I, Langenberg C, Florez JC, Saxena R, Soranzo N, Thorleifsson G, *et al.* Variants in
593 MTNR1B influence fasting glucose levels. *Nature genetics* 2009; **41**(1): 77-81.
- 594
595 31. Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, *et al.* Hundreds of
596 variants clustered in genomic loci and biological pathways affect human height. *Nature* 2010;
597 **467**(7317): 832-838.
- 598
599 32. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, *et al.* Association
600 analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature*
601 *genetics* 2010; **42**(11): 937-948.
- 602

- 603 33. Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, Steinthorsdottir V, *et al.* Meta-analysis
604 identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the
605 genetic basis of fat distribution. *Nature genetics* 2010; **42**(11): 949-960.
- 606
607 34. Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides.
- 608
609 35. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, *et al.*
610 Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 2010;
611 **466**(7307): 707-713.
- 612
613 36. He C, Holme J, Anthony J. SNP genotyping: the KASP assay. *Methods in molecular biology*
614 2014; **1145**: 75-86.
- 615
616 37. Altarejos JY, Goebel N, Conkright MD, Inoue H, Xie J, Arias CM, *et al.* The Creb1 coactivator
617 Crtc1 is required for energy balance and fertility. *Nature medicine* 2008; **14**(10): 1112-1117.
- 618
619 38. Song Y, Altarejos J, Goodarzi MO, Inoue H, Guo X, Berdeaux R, *et al.* CRT3 links
620 catecholamine signalling to energy balance. *Nature* 2010; **468**(7326): 933-939.
- 621
622 39. Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PI. SNAP: a web-
623 based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics*
624 2008; **24**(24): 2938-2939.
- 625
626 40. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, *et al.* Annotation of
627 functional variation in personal genomes using RegulomeDB. *Genome research* 2012; **22**(9):
628 1790-1797.
- 629
630 41. Rodrigo E, Fernandez-Fresnedo G, Valero R, Ruiz JC, Pinera C, Palomar R, *et al.* New-onset
631 diabetes after kidney transplantation: risk factors. *Journal of the American Society of*
632 *Nephrology : JASN* 2006; **17**(12 Suppl 3): S291-295.
- 633
634 42. Numakura K, Satoh S, Tsuchiya N, Horikawa Y, Inoue T, Kakinuma H, *et al.* Clinical and genetic
635 risk factors for posttransplant diabetes mellitus in adult renal transplant recipients treated
636 with tacrolimus. *Transplantation* 2005; **80**(10): 1419-1424.
- 637
638 43. Gourishankar S, Jhangri GS, Tonelli M, Wales LH, Cockfield SM. Development of diabetes
639 mellitus following kidney transplantation: a Canadian experience. *American journal of*
640 *transplantation : official journal of the American Society of Transplantation and the American*
641 *Society of Transplant Surgeons* 2004; **4**(11): 1876-1882.
- 642
643 44. Sulanc E, Lane JT, Puumala SE, Groggel GC, Wrenshall LE, Stevens RB. New-onset diabetes
644 after kidney transplantation: an application of 2003 International Guidelines. *Transplantation*
645 2005; **80**(7): 945-952.

646
647 45. Maes BD, Kuypers D, Messiaen T, Evenepoel P, Mathieu C, Coosemans W, *et al.*
648 Posttransplantation diabetes mellitus in FK-506-treated renal transplant recipients: analysis
649 of incidence and risk factors. *Transplantation* 2001; **72**(10): 1655-1661.

650

651

652

653

654

655

656

657

658

Table 1 Association between *CRTC2* SNPs and NODAT, blood glucose levels and BMI in patients with solid organ transplantation from the discovery study sample:

SNP	NODAT (n=156)		FBG (n=156)		BMI (n=156)	
	OR (95% CI)	P corrected	Difference (mmol/L) compared to wild-type (ref) (95% CI)	P corrected	Difference (kg/m ²) compared to wild-type (ref) (95% CI)	P corrected
<i>CRTC2 rs8450</i>						
GG	ref		ref		ref	
GA	1.81 (0.71 - 4.61)	0.46	0.24 (-0.00 - 0.49)	0.06	1.36 (0.28- 2.42)	0.02
AA	6.91 (1.52 - 31.36)	0.02	0.47 (0.07 - 0.86)	0.02	0.37 (-1.38 - 2.06)	0.76
Dominant	2.32 (0.96 - 5.63)	0.12	0.28 (0.05 - 0.51)	0.02	1.17 (0.17 - 2.35)	0.04
Recessive	5.11 (1.24 - 21.03)	0.04	0.36 (-0.03 - 0.74)	0.06	-0.24 (-2.33 - 1.68)	0.80
<i>CRTC2 rs12117078</i>						
GG	ref		ref		ref	
GC & CC	0.36 (0.11 - 1.16)	0.18	-0.07 (-0.40 - 0.24)	0.70	0.24 (-1.41 - 1.62)	0.78

NODAT: New-onset diabetes after transplantation, FBG: Fasting blood glucose, OR: odd ratio, CI: confidence interval, BMI: body mass index.

Regarding NODAT and blood glucose levels analyses: P values were adjusted for age of recipient at transplantation, sex of the recipient and baseline BMI and type of calcineurin inhibitor.

Regarding BMI analyses: P values were adjusted for age of recipient at transplantation, sex of the recipient and type of calcineurin inhibitor.

Table 2: Association between *CRTC2 rs8450* SNP in a recessive model and NODAT in the first, second and combined population of STCS and in different subgroups at risk of NODAT:

CRTC2 rs8450	GG&GA (n)		AA	P value
	N total		NODAT OR (95% CI)	
First STCS replication sample[§]:				
Total	711	Ref (n=623)	1.02 (0.62 - 1.70)	0.93
Treatment with TAC & deceased donor	281	Ref (n=244)	2.36 (1.16 - 4.78)	0.01
Second STCS replication sample[§]:				
Total	438	Ref (n=385)	2.01 (1.03 - 3.91)	0.04
Combined STCS replication sample[§]:				
Total	1149	Ref (n=1008)	1.25 (0.84 - 1.87)	0.26
Treatment with TAC & deceased donor	395	Ref (n=340)	2.08 (1.15 - 3.74)	0.02

[§]Excluding patients with previous history of diabetes before transplantation

NODAT: New-onset diabetes after transplantation, TAC: tacrolimus, OR: odd ratio, CI: confidence interval

P values were adjusted (when appropriate) for age of recipient at transplantation, sex of the recipient, hepatitis C status, baseline BMI, type of calcineurin inhibitor and type of donor.

Table 3: Association between CRTC2 rs8450 SNP and MetS traits in all kidney and non-kidney transplant recipients in the combined STCS sample

	Kidney transplant recipients			Non-kidney transplant recipients		
	n	OR (95% CI) for rs8450-AA	p-value	n	OR (95% CI) for rs8450-AA	p-value
NODAT	740	0.87 (0.50 - 1.52)	0.62	409	2.09 (1.13 - 3.86)	0.02
BMI\geq25 kg/m² at 12months post SOT	761	1.59 (0.98 - 2.58)	0.06	454	1.52 (0.84 - 2.75)	0.17
New-onset hyperlipidemia	573	1.95 (1.14 - 3.15)	0.01	434	1.73 (0.84 - 3.56)	0.14
New-onset hypertension	146	1.64 (0.43 - 6.30)	0.47	345	1.44 (0.75 - 2.78)	0.27

OR: odd ratio, CI: confidence interval

NODAT: New-onset diabetes after transplantation, BMI: body mass index

For NODAT analyses: P values were adjusted for age of recipient at transplantation, sex of the recipient, baseline BMI, type of calcineurin inhibitor and type of donor.

For BMI analyses: P values were adjusted for age of recipient at transplantation and sex of the recipient.

For new-onset hyperlipidemia analyses: P values were adjusted for age of recipient at transplantation, sex of the recipient, baseline BMI, the type of calcineurin inhibitor and treatment with corticosteroids.

For new-onset hypertension analyses: P values were adjusted for age of recipient at transplantation and sex of the recipient, treatment with corticosteroids, the type of calcineurin inhibitor and type of donor.

1 **Supplementary data:**

2 **Materials and Methods**

3 - **Population based samples:**

4 ➤ **The Meta-Analyses of Glucose and Insulin-related traits Consortium (MAGIC)**

5 ➤ **Genetic Investigation of ANthropometric Traits (GIANT) consortium**

6 ➤ **Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and**
7 **triglycerides**

8 - **Polymorphism selection and genotyping**

9 - **Statistical analysis**

10

11 **Results:**

12 - **General characteristics of the discovery sample**

13 - **General characteristics of the replication sample (STCS):**

14

15 - ***CRTC2* SNP and new-onset hypertension in the STCS**

16

17 - ***CRTC2* SNP and new-onset hyperlipidemia in the STCS**

18

19 **Supplementary Tables 1-5**

20

21

22

23

24

25

26

27 **Materials and Methods**

28 **Population based samples:**

29 **The Meta-Analyses of Glucose and Insulin-related traits Consortium (MAGIC)**

30 MAGIC is a large-scale meta-analyses of genome-wide data for continuous diabetes-related traits in
31 participants without diabetes ¹. Meta-analyses of ~2.5 million directly genotyped or imputed
32 autosomal single nucleotide polymorphisms (SNPs) were performed from genome-wide association
33 studies (GWAS). These cohorts include up to 46'186 non-diabetic participants of European descent
34 informative for FBG, the surrogate estimates of beta-cell function (HOMA-B) and insulin resistance
35 (HOMA-IR) derived from fasting variables by homeostasis model assessment and up to 15'234 non-
36 diabetic individuals informative for 2 hour oral glucose tolerance test (OGTT) ².

37

38 **Genetic Investigation of ANthropometric Traits (GIANT) consortium**

39 The GIANT consortium performed a meta-analysis of GWAS data with a discovery set of 123'865
40 individuals of European ancestry from 46 studies for height ³, BMI⁴ and waist-to hip ratio ⁵.

41

42 **Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides**

43 Data on lipid traits have been downloaded from “Genome Wide Associations Scans for Total
44 Cholesterol, HDL-C, LDL-C and triglycerides” website ^{6, 7} which is a meta-analysis of 46 lipid
45 GWASs. These studies together comprise >100,000 individuals of European descent (maximum
46 sample size 100,184 for Total Cholesterol, 95,454 for LDL-C, 99,900 for HDL-C and 96,598 for
47 triglycerides), ascertained in the United States, Europe or Australia.

48

49

50

51

52

53

54

55 **Polymorphism selection and genotyping:**

56 Polymorphisms within *CRTC2* were selected using HapMap Genome Browser (release 28)⁸. Two
57 tagging SNPs were found by limiting the search to SNPs within *CRTC2* gene \pm 1kb with a minor
58 allele frequency $>$ 5% in the Caucasian population and r^2 cutoff of 0.8: *rs8450G>A* within the 3'UTR
59 and the intronic *rs12117078G>C* SNP.

60 Genotyping of these 2 SNPS for the discovery study sample was done using Taqman allelic
61 discrimination assays (ABI PRISM 7000 Sequence Detection System; Applied Biosystems, Rotkreuz,
62 Switzerland) after genomic DNA extraction from whole blood. Life technologies Taqman genotyping
63 assays C__8722376_10 and C__30997814_10 were used for *CRTC2 rs8450G>A* and
64 *rs12117078G>C* SNPs, respectively. Genotyping were performed in the Unit of Pharmacogenetics
65 and Clinical Psychopharmacology, in Lausanne, Switzerland.

66 Genetic analyses for the first STCS replication sample were performed by KBioscience Institute in
67 United Kingdom using the novel fluorescence-based competitive allele-specific PCR technology
68 (KASPTM). Details about this technology are available at
69 <http://www.lgcgenomics.com/genotyping/kasp-genotyping-chemistry/>.

70 Genetic analyses for the second STCS replication sample were also performed using the KASPTM
71 technology on ABI 7500 Fast real-time thermocycler (Applied Biosystems). Genotyping were
72 performed in the service of infectious disease in Lausanne University Hospital, Lausanne,
73 Switzerland, according to manufacturer's protocols (LGC Genomics, UK). The KASP primers were
74 designed by KrakenTM assay design and workflow management software and further validated by
75 manufacturer (LGC Genomics, UK). Automated allele calling was performed using SDS software
76 according to standard protocols (Applied Biosystems). The following primers were used for *CRTC2*
77 *rs8450G>A* genotyping in this sample: primer allele FAM: ACCCCTAGGCATCCGGAAAAG,
78 primer allele HEX: CACCCCTAGGCATCCGGAAAAA and common primer
79 GGGTAGAGGGGAGCCCTGGAA.

80

81

82

83 **Statistical analysis:**

84 Quantitative data are presented as median and range unless otherwise mentioned, while qualitative
85 data are presented as frequency and percentage. For association studies, the chi-square (Chi^2) test or
86 the Fisher exact test for binomial variables were used. Differences in allele and genotype frequencies
87 as well as deviation from Hardy-Weinberg equilibrium were assessed using Chi^2 test.

88

89 **Discovery study sample:**

90 For NODAT analyses, logistic regression was applied adjusted for recipient's age at transplantation,
91 sex, BMI at baseline and type of CNIs. Due to nonlinearity of our models and the absence of any
92 linear transformation, the association between *CRTC2* SNPs with BMI and FBG levels was assessed
93 by fitting a Generalized Additive Mixed Model (GAMM)^{9, 10} to allow a smooth trend for the response
94 in time based on multiple observations for each patient (using a thin plate regression spline basis)
95 adjusting for recipient's age at transplantation, sex, and type of CNIs (antidiabetic treatments for FBG
96 analyses). GAMMs were fitted using the mgcv package of R (settings were fixed at package defaults).
97 In order to be more conservative, the uncertainty of estimated parameters were assessed by 1000
98 bootstraps¹¹ at the subject level and results were similar with those gained by 10'000 bootstraps.

99 Due to the small number of subjects being homozygous for the variant allele of *CRTC2*
100 *rs12117078G>C*, this SNP was analyzed in a dominant model.

101

102 **Replication samples (STCS):**

103 For the statistical analyses of NODAT, new-onset hypertension, new-onset hyperlipidemia, and
104 overweight or obese, logistic regression models were applied adjusted for recipient's age at
105 transplantation and sex. Other variables identified through the univariate analysis ($P<0.10$) were also
106 added as covariates in these models. GAMM models were applied for BMI and HDL-cholesterol
107 analyses in the STCS sample, as both models were not linear. Linear mixed models were used for
108 systolic and diastolic blood pressure analyses and for blood total cholesterol and LDL-cholesterol
109 analyses. In addition to recipient's age at transplantation, sex, BMI at baseline and type of CNIs, blood

110 pressure models and lipid models were also adjusted for antihypertensive and hypolipidemic drugs
111 intake, respectively.

112

113 **Population-based samples:**

114 The associations of *CRTC2* SNP with different MetS traits were analyzed using multivariate linear
115 regression with allele dosage in which potential confounding factors such as age, sex and smoking
116 status were added as covariates. FBG, OGTT, the surrogate estimates of beta-cell function (HOMA-B)
117 and insulin resistance (HOMA-IR) were analyzed in the MAGIC study. BMI, waist circumference and
118 waist-to hip ratio were the only adiposity traits analyzed by the GIANT Consortium. Triglycerides,
119 total cholesterol, HDL-cholesterol and LDL-cholesterol were analyzed in the “Genome Wide
120 Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides” study.

121

122 **Results:**

123 **General characteristics of the discovery sample**

124 Among the 197 patients included in the initial study, 17 patients were excluded because they were
125 diagnosed with diabetes before transplantation, 11 patients had no follow-up data in their medical
126 files, 2 patients did not sign consents for MetS-related genetic analysis, 4 patients dropped-out from
127 the study and 7 were non-Caucasians. Finally, 156 Caucasian patients were included in the current
128 study, 65% were kidney transplant recipients. The mean age at transplantation was 47.8 ± 11 years,
129 61% were male and the mean BMI pre-transplant was 24 ± 4 kg/m². During the five years of follow-up
130 post-transplantation, NODAT occurred in 45 (28.9%) patients. Most NODAT cases (67%) occurred
131 during the 1st year after transplantation. Most of the patients were treated with CSA (n=102, 65%),
132 while 35% (n=54) were treated with TAC. As expected, NODAT was higher in the TAC-treated group
133 compared to CSA¹² (46.3 vs. 19.6 % respectively). Full characteristics of our population and their
134 distribution between NODAT and non-NODAT patients are presented in Supplementary Table 1.
135 Recipient age (p=0.002), male sex (p=0.05), donor age (p=0.04), baseline BMI (p=0.0003) and the
136 type of CNIs (p<0.001) were significantly different between patients with and without NODAT
137 (Supplementary Table 1).

138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

General characteristics of the replication sample (STCS):

Overall, 1294 patients from the first STCS replication sample received a SOT from May 2008 until May 2011, among them 1219 patients had both the clinical data and the genetic material available and were included in the current study. By excluding patients younger than 18 years old and patients with multiple organ transplantation, 958 patients remained and included in the analysis. The mean age at transplantation was 52.3 ± 13 years, 66% were male and pre-transplant BMI was 24.6 kg/m^2 (range:13.7-41.2). During the follow-up period post-transplantation, NODAT, new-onset HTN and new-onset hyperlipidemia occurred in 27.4%, 67.6% and 33.1% of the patients, respectively. Among the patients included in the second STCS replication sample (n=759) with both clinical and genetic data and by excluding patients younger than 18 years old and patients with multiple organ transplantation, 667 remained and were included in the analysis. Lower incidence of NODAT, new-onset HTN and new-onset hyperlipidemia were observed in this sample, which could be explained by the shorter follow-up duration compared to the first STCS replication sample (Supplementary Table 2). Full description of the first, second and combined STCS samples are presented in Supplementary Table 2. All patients from both STCS samples were of Caucasian origin.

***CRTC2* SNP and new-onset hypertension in the STCS:**

Among the non-genetic factors, type of calcineurin inhibitor ($p < 0.0001$), type of transplanted organ ($p < 0.0001$), treatment with corticosteroids ($p < 0.0001$) and type of donor ($p = 0.02$) were associated with increased risk of new-onset hypertension after SOT in the univariate analyses and were used as covariates in the genetic models. A non-significant association was observed between *CRTC2* *rs8450G>A* and new-onset hypertension, except for patients carrying the *rs8450-GA* genotype which showed a protective effect against new-onset hypertension (n=491, OR=0.62, 95%CI: 0.42 - 0.94, $p = 0.02$) compared to wild type genotype. *CRTC2* *rs8450G>A* SNP was not associated with systolic or diastolic blood pressure levels (data not shown).

166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190

***CRTC2* SNP and new-onset hyperlipidemia in the STCS:**

Among the non-genetic factors, increased recipient's age ($p=0.003$), baseline BMI ($p=0.006$), type of calcineurin inhibitor ($p=0.005$) and treatment with corticosteroids ($p<0.0001$) were associated with increased risk of new-onset hyperlipidemia after SOT in the univariate analyses and were used as covariates in the genetic models.

191 **Supplementary Table 1: Clinical characteristics of the discovery study sample:**

Characteristic	Total	Non-NODAT	NODAT	p-value
Recipient age at transplantation (years), median (range)	48 (22-68)	46 (22-68)	53 (28-68)	0.002
Recipients sex (Males) [%]	60.9	55.9	73.3	0.05
Living donor [%]	11.5	11.7	11.1	0.92
Donor age (years), median (range)	43.5 (10-73)	41 (10-73)	48 (13-69)	0.04
Donor sex (Males) [%]	56.8	63.2	54.8	0.06
Organ, n [%]				
lung	17 (10.9)	10 (9.0)	7 (15.6)	
kidney	102(65.4)	75 (67.6)	27(60.0)	0.46
liver	37(23.7)	26 (23.4)	11 (24.4)	
BMI pre-transplant (kg/m ²), median (range)	23.5 (15.8-37.3)	22.3 (15.8-36.2)	26.4 (18.8-37.3)	0.0003
BMI 5 year follow-up (kg/m ²), median (range)	26.0 (16.7-43.6)	25.5 (16.7-43.6)	27.1 (19.9-42.3)	0.10
Recipient CMV infection (R ⁺) [%]	49.3	52.5	42.2	0.25
Donor CMV infection (D ⁺) [%]	61.5	59.2	66.7	0.39
Recipient and Donor CMV infection (R ⁺ /D ⁺) [%]	27.6	27.0	28.9	0.81
Calcineurin inhibitors, n [%]				
TAC	54 (34.6)	29 (26.1)	25 (55.6)	<0.001
CSA	102 (65.4)	82 (73.9)	20 (44.4)	
First transplantation [%]	80.1	77.5	86.7	0.19
Acute rejection episode during first year post-transplant [%]	47.4	47.2	47.7	0.96

NODAT: New-onset diabetes after transplantation, BMI: body mass index, CMV: cytomegalovirus, R: recipient, D: Donor, TAC: Tacrolimus, CSA: Cyclosporine.

192

193

194

195

196 **Supplementary Table 2: Clinical characteristics of STCS[§]**

Characteristic	First STCS replication sample (n=958)	Second STCS replication sample (n=667)	combined STCS replication sample (n=1625)
Recipient age at transplantation (years), median (range)	55 (18-79)	55 (79)	55 (18-79)
Recipients sex (Males) [%]	66.2	64.6	65.5
Period of follow-up (months), median (range)	12 (0-60)	6 (0-48)	6 (0-60)
Living donor [%]	28.7	24.0	26.7
Donor age (years)	53 (1-86)	55 (1-88)	53 (1-88)
Donor sex (Males) [%]	53.5	53.4	53.4
Organ, n [%]			
Kidney	624 (65.1)	360 (54.0)	984 (60.5)
Liver	158 (16.5)	142 (21.3)	300 (18.5)
Lung	94 (9.8)	97 (14.5)	191 (11.8)
Heart	65 (6.8)	57 (8.6)	122 (7.5)
Islets and pancreas [£]	17 (1.8)	11 (1.6)	28 (1.7)
BMI pre-transplant (kg/m ²), median (range)	24.6 (13.7-41.2)	24.9 (13.4-43.5)	24.8 (13.4-43.5)
BMI 1 year follow-up (kg/m ²), median (range)	25.3 (15.4-44.6)	24.8 (13.7-41.7)	25.2 (13.7-44.6)
Recipient CMV infection (R ⁺) [%]	57.5	59.5	58.2
Donor CMV infection (D ⁺) [%]	53.4	58.4	56.2
Recipient and Donor CMV infection (R ⁺ /D ⁺) [%]	33.0	36.3	34.3
Calcineurin inhibitors, [%]			
TAC	65.5	67.1	66.3
CSA	26.7	26.1	26.4
None	7.8	6.8	7.3
Incidence of NODAT [%]	27.4	18.7	24.4
Incidence of New hypertension [%]	67.6	32.8	53.6
Incidence of New hyperlipidemia [%]	33.1	13.8	26.6

[§]patients included in this table represent the global STCS sample with the clinical data available for the analyses without the exclusion of patients with diabetes, hypertension or hyperlipidemia before transplantation.

[£]These patients had type I diabetes mellitus as an indication for transplantation and were excluded from NODAT analysis. Abbreviations: BMI: body mass index, CMV: cytomegalovirus, R: recipient, D : Donor, TAC: Tacrolimus, CSA: Cyclosporine, NODAT: New-onset diabetes after transplantation, New hypertension: new onset hypertension after transplantation, New hyperlipidemia: new onset hyperlipidemia after transplantation.

197

198

199

200 **Supplementary Table 3: Genotype frequencies in the discovery sample and in the first, the**
 201 **second and the combined STCS samples. Hardy-Weinberg equilibrium and a comparison**
 202 **between the observed minor allele frequency and HapMap minor allele frequency in the**
 203 **Caucasian population.**

SNP	CRTC2 SNPs Frequencies n (%)			Hardy-Weinberg Equilibrium (p-value)	MAF	
					observed	HapMap
Discovery sample (n=156)						
rs8450	GG	GA	AA	0.87	0.30	0.30
	78 (50.0)	64 (41.0)	14 (9.0)			
rs12117078	GG	GC	CC	0.75	0.10	0.10
	126 (80.8)	28 (17.9)	2 (1.3)			
STCS						
rs8450	GG	GA	AA			0.30
First STCS sample (n=949)[§]	408 (43.0)	429 (44.2)	112 (11.8)	0.96	0.34	
second STCS sample (n=667)	281 (42.1)	302 (45.3)	84 (12.6)	0.84	0.35	
Combined STCS sample (n=1616)	689 (42.7)	731 (45.2)	196 (12.1)	0.92	0.35	

204 [§]9 missing genotypes for this sample.

205 MAF: minor allele frequency.

Supplementary Table 4: Association between *CRTC2 rs8450* SNP and NODAT in total population of STCS in different subgroups at risk of NODAT:

CRTC2 rs8450	Additive model				Dominant model				
	GG	GA	AA	GG	GA&AA				
	N	NODAT OR (95% CI)	p-value	NODAT OR (95% CI)	p-value	NODAT OR (95% CI)	p-value		
First STCS replication sample:									
Total	711	ref	0.97 (0.68 - 1.38)	0.85	1.01 (0.59 - 1.73)	0.98	ref	0.97 (0.70 - 1.37)	0.88
Treatment with TAC & deceased donor	281	ref	1.08 (0.63 - 1.85)	0.79	2.45 (1.14 - 5.26)	0.02	ref	1.31 (0.79 - 2.17)	0.30
Second STCS replication sample:									
Total	438	Ref	0.81 (0.47 - 1.40)	0.46	1.80 (0.88 - 3.71)	0.11	ref	0.99 (0.60 - 1.62)	0.96
Combined STCS replication sample:									
Total	1149	ref	0.89 (0.66 - 1.20)	0.44	1.18 (0.76 - 1.81)	0.46	ref	0.95 (0.72 - 1.25)	0.70
Treatment with TAC & deceased donor	395	ref	0.81 (0.51 - 1.30)	0.39	1.86 (0.99 - 3.52)	0.06	ref	1.00 (0.65 - 1.54)	0.99

TAC: tacrolimus, OR: odd ratio, CI: confidence interval

P values were adjusted (when appropriate) for age of recipient at transplantation, sex of the recipient, hepatitis C status, baseline BMI, type of calcineurin inhibitor and type of donor.

Supplementary Table 5: Associations between *CRTC2* rs1572788 T>C (in complete linkage disequilibrium (LD) with rs8450 G>A) and glycemc, lipidemic and obesity traits in several population-based samples:

	Effect of each <i>rs1572788</i> C-allele (complete LD with <i>rs8450</i> A-allele)			
phenotype	n	beta	p-value	Explained variance*
Glucose [£]	46186	0.01	0.004	0.0002
2h glucose tolerance test [£]	15234	-0.02	0.26	
HOMA-β [£]	46186	-0.008	0.03	0.0001
HOMA-IR [£]	46186	0.0002	0.96	
Total cholesterol [§]	97148	0.005	0.16	
HDL-Cholesterol [§]	96908	-0.008	0.013	0.0001
LDL-Cholesterol [§]	92503	0.006	0.06	
Triglycerides [§]	93562	0.009	0.004	0.0001
BMI [§]	113955	0.004	0.37	
Waist-hip ratio [§]	55282	0.002	0.71	
Waist circumference [§]				
Male	38305	0.008	0.32	
Female	47320	0.01	0.19	

explained variance by the polymorphism (only calculated for p<0.05).

£ This clinical variable was analyzed in the MAGIC study.

§ This clinical variable was analyzed in the “Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides”.

§ This clinical variable was analyzed in the GIANT study.

2h: 2 hours, HOMA: homeostatic model assessment, β : beta-cell function, IR: insulin resistance, BMI: Body mass index.

References:

1. Prokopenko I, Langenberg C, Florez JC, Saxena R, Soranzo N, Thorleifsson G, *et al.* Variants in MTNR1B influence fasting glucose levels. *Nature genetics* 2009; **41**(1): 77-81.
2. Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nature genetics* 2010; **42**(2): 105-116.
3. Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, *et al.* Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 2010; **467**(7317): 832-838.
4. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature genetics* 2010; **42**(11): 937-948.
5. Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, Steinthorsdottir V, *et al.* Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nature genetics* 2010; **42**(11): 949-960.
6. Genome Wide Associations Scans for Total Cholesterol, HDL-C, LDL-C and triglycerides.
7. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, *et al.* Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 2010; **466**(7307): 707-713.
8. Barnes MR. Navigating the HapMap. *Briefings in bioinformatics* 2006; **7**(3): 211-224.
9. Wood SN. GAMMs with R. In: Carlin BP, Chatfield C, Tanner M, Zidek J (eds). *Generalized Additive Models - An Introduction with R*. Chapman & Hall/CRC: Broken, 2006, pp 319-324.
10. Lin X, Zhang D. Inference in generalized additive mixed models by using smoothing splines. *J R Statist Soc B* 1999; **61**(2): 381-400.
11. Davison AC, Hinkley DV. *Bootstrap Methods and their Application*. Cambridge University Press: Cambridge, New York, 1997, 1-592pp.
12. Opelz G, Dohler B, Collaborative Transplant S. Influence of immunosuppressive regimens on graft survival and secondary outcomes after kidney transplantation. *Transplantation* 2009; **87**(6): 795-802.