

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Waddlia chondrophila: from biology to pathogenicity.

Authors: de Barsy M, Greub G

Journal: Microbes and infection

Year: 2013 Dec

Volume: 15

Issue: 14-15

Pages: 1033-41

DOI: 10.1016/j.micinf.2013.09.010

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

1 *W. chondrophila*: from biology to pathogenicity

2 Marie de Barsy¹ and Gilbert Greub¹

3

4 ¹ Institute of Microbiology, University Hospital Center and University of Lausanne, Lausanne,

5 Switzerland

6

7

8

9

10

11

12

13 Corresponding author:

14 Prof. Gilbert Greub, MD PhD

15 Center for Research on Intracellular Bacteria (CRIB), Institute of Microbiology

16 University Hospital Center and University of Lausanne,

17 Bugnon 48, 1011 Lausanne, SWITZERLAND

18 Phone : +41 21 314 49 79, Fax : +41 21 341 40 60

19 e-mail : gilbert.greub@chuv.ch

20

21

22

23

24

25

26 **Abstract**

27 *Waddlia chondrophila* is an emerging pathogen causing miscarriages in humans and abortions in
28 ruminants. The full genome of this *Chlamydia*-related bacterium has been recently completed,
29 providing new insights into its biology and evolution. Moreover, new cell biology approaches and
30 the use of novel inhibitors have allowed detailed investigations of its interaction with host cells.

31

32 **Keywords:** intracellular bacteria, *Chlamydia*-related organism, *Waddlia chondrophila*,
33 pathogenesis

34

35 **Abbreviations:** elementary bodies (EBs), reticulate bodies (RBs), aberrant bodies (Abs),
36 endoplasmic reticulum (ER), peptidoglycan (PG), outer membrane protein (OMP), Type III
37 secretion system (T3SS), bacteria containing vacuole (BCV), post infection (p.i.)

38

39

40

41 **1. Introduction**

42 *Waddlia chondrophila* is an intracellular obligate bacterium that belongs to the *Chlamydiales* order.
43 The *Chlamydiales* order includes six different families, including the *Waddliaceae* family. *Waddlia*
44 *chondrophila* was first isolated from an aborted bovine fetus in USA in 1990 [1]. Twelve years
45 later, *W. chondrophila* was also isolated from a second bovine case [2]. In 2005, a novel species
46 which shares 91 % of identity with *Waddlia chondrophila*, was identified in Malaysia from a fruit
47 bat and called *Waddlia malaysiensis* [3]. *Waddlia chondrophila* is considered as an abortigenic
48 bacteria in ruminants [1, 2] and is likely responsible of economical losses. The pathogenic role of
49 *W. chondrophila* in humans is supported by a strong association between *W. chondrophila*
50 seropositivity and human miscarriage [4-6]. Moreover, *W. chondrophila* was also detected in
51 respiratory samples from patients with bronchiolitis or pneumonia [7, 8]. Despite the clinical and
52 veterinary importance of this pathogen and its zoonotic potential, little is known regarding the
53 biology and pathogenicity of this bacterium. In this review, we summarize the current knowledge
54 on *Waddlia chondrophila* biology by focussing on its cell biology, genome, metabolism, and
55 membrane proteins.

56

57 **2. Nascent role of *W. chondrophila* in human miscarriages and respiratory diseases**

58 The potential abortigenic role of *W. chondrophila* in cattle raises the hypothesis that *W.*
59 *chondrophila* could be involved in human miscarriages, as reported by Baud and colleagues [4].
60 This prospective study on 438 women comprised 69 women with sporadic miscarriages, 200 with
61 recurrent miscarriages and 169 control women with uneventful pregnancies [4]. By
62 immunofluorescence, 100 women were positive for *Waddlia* with an anti-*Waddlia* IgG titer ≥ 64 .
63 The seroprevalence for women with sporadic miscarriages (31.9%) and women with recurrent
64 miscarriage (33%) was higher than for control women (7%, $p < 0.05$). The presence of anti-*Waddlia*
65 IgG was confirmed for 97 women by western blot using *W. chondrophila* as the antigen [4]. The
66 specificity of the anti-*Waddlia* antibody was demonstrated by adsorption experiments. In a second

67 study, a case was documented, which not only exhibited a positive serology for anti-*Waddlia*
68 antibodies, but was also confirmed by *Waddlia* specific real-time PCR performed on cervicovaginal
69 swabs and by immunohistochemistry [5]. Immunohistochemistry demonstrated the presence of *W.*
70 *chondrophila* in glandular epithelium cells. This is the first evidence of the presence of this
71 bacterium in the human placenta itself.

72 The transmission mode of *Waddlia* to human remains an important question. Since *C. trachomatis*
73 is a sexually transmitted pathogen whose role in miscarriages is suspected [9], sexual transmission
74 may be possible for *Waddlia*. In the first study, the *C. trachomatis* seropositivity did not differ
75 between the 97 women seropositive for *Waddlia* and the seronegative women [4]. From this
76 absence of correlation between *C. trachomatis* and *Waddlia*, we can conclude that there is neither a
77 significant cross-reactivity nor the same transmission mode. This suggests that *W. chondrophila* is
78 not sexually transmitted. Interestingly, further statistical analysis showed that *Waddlia*
79 seropositivity is associated with animal contact, supporting the zoonotic risk of *W. chondrophila*
80 [4]. This potential is further strengthened when considering that the host range of *W. chondrophila*
81 is known to include cattle [1, 2]. Other transmission modes such as the water network are possible
82 since free-living amoebae present in water are permissive to *W. chondrophila* infection and might
83 represent a huge widespread reservoir for this intracellular bacterium [10]. A recent study on
84 drinking and well water (n=70) confirmed the presence of *W. chondrophila* by qPCR on 10 of the
85 40 well water samples investigated [11]. Infection by ingestion of contaminated meat or milk is also
86 possible [4].

87 The implication of *Chlamydia*-related bacteria such as *Simkania negevensis* and *Parachlamydia*
88 *acanthamoebae* in respiratory disease are well documented [12-14]. The presence of *W.*
89 *chondrophila* in respiratory samples was also investigated. In one study on 389 patients with
90 community-acquired pneumonia, one case was positive by PCR and sequencing of the 16S rRNA
91 gene fragment revealed 98% and 99.7% sequences similarity to the 16S rRNA gene of *W.*
92 *chondrophila* isolate WSU 86-1044 and 2032-99, respectively [7]. As part of a second study, a *W.*

93 *chondrophila* specific real-time PCR was developed and applied to 32 nasopharyngeal samples from
94 children with bronchiolitis not due to respiratory syncytial virus (RSV), the most common etiology
95 of bronchiolitis [8]. Three samples were *Waddlia* positive. Among these three patients, one was also
96 infected by another microbe [8].

97 In conclusion, *W. chondrophila* is likely involved in human miscarriage and possibly involved in
98 respiratory disease. Additional studies are needed to precisely evaluate the incidence and
99 importance of this bacterium in these clinical settings.

100

101 **3. *W. chondrophila*, an emerging veterinary pathogen**

102 *W. chondrophila* was isolated twice from aborted bovine fetuses in USA and in Germany [1, 2].
103 The German fetus was coinfecting with *Neospora caninum* which precluded a definitive diagnosis
104 on the cause of abortion. A serological test was developed and applied to bovine sera, showing an
105 association between anti-*Waddlia* antibodies and bovine abortion [15]. Moreover, 2 bovine fetuses
106 were experimentally infected with *W. chondrophila* and one fetus died within 2 weeks [15]. These
107 results support an abortigenic potential of *W. chondrophila* in cattle.

108 Previous studies reported in this review, showed the potential role of *W. chondrophila* in abortion in
109 ruminants and humans, raising the question of whether this bacterium has the same role in other
110 mammals. In this prospect, the potential role of *W. chondrophila* in porcine abortion was assessed
111 by Koschwanez and colleagues [16]. None of the aborted fetuses (n=286) were positive for the
112 *Waddlia* specific real time PCR suggesting that *Waddlia* is not an abortigenic agent in Swiss sows
113 [16].

114 Recently, it was reported that *W. chondrophila* is able to enter and proliferate in two different fish
115 cell lines suggesting a potential pathogenicity toward fishes that could constitute a potential
116 reservoir of *W. chondrophila* [17]. This hypothesis is supported by the identified role of other
117 *Chlamydia*-related bacteria, such as *Candidatus Piscichlamydia salmonis* [18] and *Candidatus*
118 *Clavochlamydia salmonicola* [19], in epitheliocystis disease in fish.

119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144

4. The biphasic developmental cycle of *W. chondrophila*

W. chondrophila was first cultivated in bovine turbinate cells (BT). A cytopathic effect was observed about 2-3 days post infection (p.i.). Light microscopy revealed bacteria within cytoplasmic inclusions ranging in size from 0.2-0.4 μm [1]. Further studies in BT cells and in P388D1 mouse macrophages demonstrated a biphasic developmental cycle similar to these of other members of the *Chlamydiales* order. This cycle begins with infectious elementary bodies (EBs), which enter in the host cells. Once inside the cells, EBs convert to metabolically active reticulate bodies (RBs), which divide by binary fission (**Fig. 1**). Finally, RBs redifferentiate into EBs, which are released after cell lysis and can initiate a new infectious cycle [20]. EBs of *W. chondrophila* are characterized by a nuclear condensation (**Fig. 1A**). The mechanism regulating nuclear condensation is still poorly understood.

The bacteria of the *Chlamydiales* order also exhibit an alternative developmental stage characterized by an abnormal size, enlarged RB-like structures called aberrant bodies (ABs) [21]. Aberrant bodies are considered as persistent forms. In endometrial cells (Ishikawa), *W. chondrophila* develop into aberrant bodies as early as 72h p.i. and their number and their size increase over time [21]. When the culture medium is renewed every day, aberrant bodies are observed in only 10 % of bacteria containing vacuole compared to 100 % without medium change suggesting that aberrant bodies developed in response to starvation [21]. Interestingly, these persistent forms can revert to proliferating bacterial stages when fresh medium is added after 6 days p.i. [21]. We think that persistent forms, present in endometrial cells of the glandular decidual epithelium, will revert to proliferating bacteria in early days of pregnancy due to local metabolic changes, then leading to inflammation and miscarriage. This explains the occurrence of recurrent miscarriages [4, 6].

145

146 **5. An overview of the *Waddlia* Genome**

147 The complete annotated genome sequence is required for the thorough characterization of a
148 genetically intractable obligate intracellular bacterium such as *W. chondrophila* since it allows the
149 identification of virulence factors by homology and the understanding of its biology.

150 *Waddlia chondrophila* WSU 86-1044 genome consist of a circular chromosome of 2.1Mb and a
151 circular plasmid of 15.6 Kb, with a G+C content of 43.8% and 37.6% respectively [22]. This
152 corresponds to a 2 fold larger genome size compared to the genomes of *Chlamydiaceae* (**Table 1**).
153 *W. chondrophila* possesses two copies of the gene *dnaA* whose positions are not linked to the
154 minimum of the cumulative GC skew corresponding to the origin of replication [23-25]. The
155 genome encompasses 2 rRNA operons, 27 tRNA genes and 1934 protein coding genes [22]. Among
156 these proteins, 65% possess a putative function or a family membership, 13% are conserved
157 hypothetical proteins and 23% show no similarity to known proteins [22]. A core set of
158 *Chlamydiales* genes was determined and comprises, among others, all essential genes coding for
159 proteins involved in DNA replication, transcription and RNA translation (493 genes). A large
160 proportion of these genes encode for family specific proteins and proteins poorly conserved at the
161 amino acid level.

162 The *W. chondrophila* plasmid is present in about 11 copies per cell and encodes 22 proteins that
163 show no homology to other chlamydial plasmid proteins, except an integrase that exhibits 52% of
164 identity to the plasmid integrase pCpA1_003 of *C. psittaci* [22]. The plasmid encodes two
165 transposases and 7 chromosomal regions, ranging from 57 bp to 889 bp, that show 99% to 100% of
166 identity with these two transposases [22]. Two adjacent genes present on the plasmid were also
167 found integrated in the *W. chondrophila* chromosome sharing 88% of identity with their plasmid
168 counterparts. One of these two genes encodes a protein homologous to MazF, an endoribonuclease
169 of the toxin-antitoxin system, MazEF [22, 26]. This system could be involved in the plasmid
170 stability during cell division. Surprisingly, the adjacent gene of *mazF* does not show sequence

171 similarities with *mazE* gene which encodes the labile antitoxin that prevents the lethal effect of the
172 stable toxin, MazF [22]. The biological function of the plasmid is still unknown. *W. chondrophila*
173 and almost all bacteria of the *Chlamydiales* order are considered as untractable organisms since
174 targeted mutagenesis is not possible. Nevertheless, the presence of the *W. chondrophila* plasmid
175 implies that plasmid-based genetic could be possible for this bacterium as recently reported for *C.*
176 *trachomatis* [27], leaving an opened-window to genetic approaches.

177 Recently, the second isolate of *W. chondrophila* (strain 2032/99) identified in Germany was
178 sequenced [28]. The general features of the chromosome are similar to *W. chondrophila* strain WSU
179 86-1044 (Table 1). Interestingly, *W. chondrophila* 2032/99 does not harbor a plasmid. However, 9
180 plasmid proteins have homologs in *W. chondrophila* 2032/99 genome distributed over 5 contigs,
181 suggesting a chromosomal integration in this strain [28]. A common ancestor is shared by plasmids
182 of *Chlamydiaceae* [29]. The presence of one protein encoded on the *W. chondrophila* plasmid
183 homologue to a plasmid-encoded protein of the *Chlamydiaceae*, support the hypothesis of a
184 common evolutionary origin of all chlamydial plasmids.

185

186 **6. Metabolic secrets uncovered by the *Waddlia* genome**

187 *W. chondrophila* genome analysis revealed that this strict intracellular bacteria is able to produce
188 energy independently from its host through oxidative phosphorylation. *W. chondrophila* harbours a
189 complete TCA cycle and glycolysis allowing production of reduced cofactors that are funnelled
190 along the electron transport chain to generate ATP [22]. In addition to a V_1V_0 ATPase complex
191 conserved in the *Chlamydiaceae*, *W. chondrophila* possesses a F_1F_0 ATP synthase complex
192 increasing ATP production. This feature could improve the adaptation of *W. chondrophila* to
193 energy-depleted environment. Moreover, a glyoxylate shunt is present allowing utilization of fatty
194 acids or acetate as carbon source [22].

195 Interestingly, *W. chondrophila* possesses genes to produce at least ten of the twenty classical amino
196 acids compared to *C. trachomatis* that is only able to produce three amino acids (**Table 2**). Unlike

197 *Chlamydia* spp., *W. chondrophila* genome completely lacks the genes involved in the tryptophan
198 biosynthesis. This bacterium seems also unable to synthesize tyrosine and phenylalanine [22].
199 Nevertheless, five transporters dedicated to general or specific aromatic amino acids such as
200 tyrosine and phenylalanine are encoded in the *W. chondrophila* genome. Several oligopeptide and
201 amino acid transporters or permeases are also identified allowing importing these compounds from
202 the environment [22].

203 Concerning lipid metabolism of *W. chondrophila*, additional enzymes for glycerophospholipid,
204 glycerolipid and sphingolipid metabolism are found compared to other members of the
205 *Chlamydiales* order. *W. chondrophila* also possesses a complete operon encoding the mevalonate
206 pathway in the biosynthesis of isoprenoid precursors that is not present in the *P. amoebophila* and
207 *C. trachomatis* genomes [22].

208 Unlike other *Chlamydiales*, *W. chondrophila* possesses all enzymes to convert L-glutamine in UMP
209 and all derivatives of pyrimidine. In contrast, this bacterium does not possess a complete purine
210 biosynthesis pathway but an active purine conversion, specific to *W. chondrophila*, is present. In the
211 *W. chondrophila* genome, no homolog to the *P. amoebophila* NAD⁺/ADP transporter were
212 identified but it seems that *W. chondrophila* is able to synthesize NAD from an intermediary
213 metabolite such as quinolinate or nicotinamide imported through another system. Interestingly, *W.*
214 *chondrophila* genome encodes five nucleotide transporters similar to *ntt_1,2* and *3* of *P.*
215 *amoebophila* potentially involved in the import of all nucleotides including ATP [22]. These genes,
216 likely originated by serial duplications, suggest that the *Chlamydiales* ancestors were already
217 intracellular and imported nucleotides from host cells more than 1 billion years ago [30].

218 Taking together, this genome analysis showed that *W. chondrophila* harbours many enzymes for the
219 synthesis of co-factors, nucleic acids and amino acids, and a complete central metabolism providing
220 energy necessary for biosynthesis. This analysis also suggests that *W. chondrophila* is more
221 independent of the host cell compared to other *Chlamydiales* and represent the best chance to date
222 for cultivating a member of the *Chlamydiales* order on an axenic medium.

223

224 **7. Insights into the evolution of the chlamydial cell wall from *Waddlia* genome**

225 The presence of peptidoglycan (PG) in the chlamydial cell wall has been debated for a long time.
226 Recent studies demonstrated that *Chlamydiaceae* exhibit almost a complete pathway for
227 peptidoglycan biosynthesis. Moreover, some PG synthesis enzymes are functional. Interestingly,
228 PG synthesis genes are expressed primarily during reticulate body development and division
229 suggesting a potential role of PG in cell division. *Chlamydiales*, including *W. chondrophila*, do not
230 possess the *ftsZ* gene. FtsZ is a highly conserved tubulin-like protein involved in cell division of
231 most bacteria. FtsZ localizes at midcell, polymerize to a ring structure called Z-ring and recruits
232 proteins involved in cell division such as FtsI, FtsW and AmiC. The absence of *ftsZ* gene, in *W.*
233 *chondrophila* and in other *Chlamydiales*, supports a FtsZ-independent cell division mechanism
234 where PG could be involved. MreB, an actin homolog which localized at midcell, might represent a
235 functional homolog of FtsZ in *W. chondrophila* (Jacquier *et al.*, submitted).

236 The inability to detect PG could be explained by the fact that all attempts were made on EBs [31].
237 In EBs, a highly disulphide-linked proteinaceous layer serves as a functional equivalent to PG [32].
238 In contrast, RBs contain less cross-linked membrane proteins. Several proteins form this cross-
239 linked proteins complex via disulfide bond, including, OmcA and OmcB which belong to the
240 polymorphic outer membrane protein (OMP) family, and the outer membrane protein beta-barrel
241 porins OmpA and PorB.

242 OmpA was shown to be involved in different mechanisms such as attachment, infection, surface
243 exposure and has antigenic properties. Previous studies, performed on *Chlamydia* spp.,
244 demonstrated that OmpA is an adhesin promoting non-specific interaction with host cells [33].
245 OmpA may also act as porin during chlamydial proliferation [34]. Surprisingly, a novel OMP
246 family comprising 11 putative β -barrel proteins or porins with C-rich signature was identified in *W.*
247 *chondrophila* genome [22]. Characterization of these putative adhesins is in progress.

248 Homologs to *omcA* and *omcB* genes were detected in the *W. chondrophila* genome. Furthermore,
249 the five adjacent genes shares similarities with the N-terminal region of OmcA/B and possess
250 conserved cystein residues, supporting an extended *omc* family in *W. chondrophila*. OmcB protein
251 of *C. trachomatis* was shown to be a surface-exposed protein that functions as an adhesin
252 suggesting the potential role of these Omc family proteins in adhesion [35].

253 A putative autotransporter protein sharing similarities with *P. amoebophila* gene was also
254 identified. This protein likely belongs to the chlamydial polymorphic membrane protein (PMP)
255 family. The pmp proteins are classical autotransporters with a passenger domain surface-localized
256 or secreted, responsible of their function such as adhesion [36]. The *C. trachomatis* pmp family has
257 9 members and *C. pneumoniae* 21 members [37, 38]. This autotransporter could function as an
258 adhesin necessary for *W. chondrophila* infection.

259 Type III secretion system (T3SS), also called injectisome, translocates bacterial proteins into host
260 cell cytoplasm [39]. This machinery is composed of about 25 proteins, localized in the bacterial cell
261 envelope, i.e. the inner and the outer membrane, and is able to span eukaryotic plasma membrane to
262 inject bacterial proteins [40]. Many pathogens, such as *Salmonella* and *Yersinia*, possess a T3SS in
263 order to hijack host cell processes [41, 42]. The T3SS components are highly conserved compared
264 to its effectors which are mostly species-specific. All *Chlamydiales* genomes encode an almost
265 complete T3SS distributed in 4 clusters of genes. This unusual spread of T3SS-encoding genes
266 along the chlamydial chromosome indicates that the T3SS was already present in the common
267 ancestor of *Chlamydia* and *Waddlia* which diverged about 1 billion years ago. Like other members
268 of the *Chlamydiales* order, *W. chondrophila* possesses almost all genes coding the T3SS. The main
269 differences reside in the effector proteins of *W. chondrophila*. The only waddliial effector identified
270 by homology to known *Chlamydiaceae* effectors, is Pkn5, suggesting that filling the gap in this area
271 will probably improve our knowledge of *W. chondrophila* biology.

272 The functionality and the role of the T3SS in *Waddlia* survival and proliferation in human
273 macrophages were demonstrated by using T3SS specific inhibitors, which inhibited bacterial

274 growth in human macrophages. It could be interesting to assess the role of the T3SS in specific
275 events such as adhesion, internalization and recruitment of mitochondria or endoplasmic reticulum
276 (ER).

277

278 **8. Growth of *Waddlia* in different cell lines**

279 *W. chondrophila* was first reported to be able to grow in BT cells and in mouse macrophages. *W.*
280 *chondrophila* can also be cultivated in buffalo green monkey cells, Mc Coy cells, human fibroblasts
281 [2], peripheral blood mononuclear cells (PBMC) [43], Vero cells, A549 pneumocytes and in
282 Ishikawa endometrial cells [21].

283 In PBMC-derived macrophages, the entry of bacteria is taking place during the first 8h after
284 inoculation. In about 24-36 h, the number of bacteria increase by 2-3 logs and the generation time is
285 estimated to 2.8 h. At 48 h p.i., a cytopathic effect is observed with a host cell survival rate
286 decreasing to 50%. This cytopathic effect is associated to the increase in bacterial number at 24 h
287 p.i. [43].

288 In Vero cells and A549 pneumocytes, the growth kinetic of *W. chondrophila* is very similar to that
289 observed in PBMC-derived macrophages, with a latent phase lasting 8 h during which bacteria enter
290 in the cells and differentiate into RBs [21]. This phase is followed by a proliferation phase where
291 the number of bacteria per cell increase by 2.5 log between 8 to 24 h p.i.. The infected cells are then
292 lysed and released bacteria can initiate a new infection cycle. At 72 h p.i., the number of bacteria
293 reaches a plateau because all the cells are infected. In Ishikawa endometrial cells, the growth kinetic
294 of *W. chondrophila* is similar until 24 h p.i. [21]. The bacterial number then slightly decreases
295 between 24 h and 96 h, and no second round of infection was observed. A cytopathic effect of *W.*
296 *chondrophila* is shown in Vero and A549 cells by measuring cell viability. In contrast, in Ishikawa
297 cells, the bacteria do not exert a significant cytopathic effect [21]. This suggests that the bacteria
298 remain in the cell without proliferation after 24 h p.i. and without inducing cell lysis. Interestingly,
299 aberrant bodies appeared in Ishikawa cells at 72 h p.i. (See section 4.).

300 As *Candidatus* Piscichlamydia salmonis, a new family member belonging to the *Chlamydiales*
301 order, is responsible of a common infection in many fish species, called epitheliocystis, the
302 permissivity of two fish cell lines, to *W. chondrophila* was investigated [17]. By quantitative PCR
303 and immunofluorescence, it was shown that *W. chondrophila* is able to proliferate in these two fish
304 cell lines, EPC-175 (Fathead Minnow) and RTG-2 (rainbow trout). This rapid proliferation was
305 associated with a cytopathic effect [17]. Thus, *W. chondrophila* can be used to develop an *in vivo*
306 model of epitheliocystis in fish species.

307

308 **9. Cell biology studies provide insight into the relationship between *Waddlia* and its host**

309 As previously reported in this review, *W. chondrophila* extensively proliferate within human
310 macrophages (PBMC) and induce cell lysis. Different strategies can be adopted by bacterial
311 pathogens to survive within macrophages. The bacterial pathogen can avoid delivery in a
312 degradative compartment such as lysosomes or develop strategies to survive in this compartment. A
313 common tactic is to hijack the endocytic pathway to finally create a proliferation niche that has
314 unique properties. In order to understand how *W. chondrophila* survive in human macrophages,
315 intracellular trafficking was studied [44]. The bacterial intracellular trafficking is followed by the
316 successive acquisition of organelle-specific marker on the bacteria-containing vacuoles (BCV). It
317 was shown that *W. chondrophila* transiently acquired the early endosome marker, EEA1, with a
318 peak at 15 min [44]. Interestingly, *W. chondrophila* never co-localize with LAMP1 and v-ATPase,
319 two late endosome-lysosome markers. Moreover, by using the LysoTracker probe, a fluorescent
320 acidotropic probe, to monitor phagosome acidification, it was shown that vacuoles containing *W.*
321 *chondrophila* never co-localize with this probe suggesting that the vacuoles are not acidic [44].
322 These results demonstrate that *W. chondrophila* BCVs interact with early but not late endosomal
323 compartment and that *W. chondrophila* rapidly evade the endocytic pathway [44].
324 Host cell mitochondria are rapidly and gradually recruited to *W. chondrophila* BCVs, with 50%
325 positive at 2 h p.i. and more than 80% at 8 h p.i. (**Fig. 2A**). Electron microscopy shows an intimate

326 association between mitochondria and *W. chondrophila* BCVs (**Fig. 2B**). Some vacuoles containing
327 *W. chondrophila* are already surrounded by mitochondria as early as 30 min p.i. suggesting that
328 recruitment start just after phagocytosis [44]. A specific association between *W. chondrophila* and
329 host cell mitochondria was already shown in 1990 using BT cells and P388D1 mouse macrophages
330 [1, 45]. A clear association between host cell mitochondria and the bacteria-containing vacuoles
331 was also observed in Vero and Ishikawa cells [21]. One hypothesis could be that mitochondria are
332 required to support bacterial proliferation. After treatment of PBMC-derived macrophages,
333 simultaneously, with Nocodazole and Cytochalasin D, which are respectively, a microtubule-
334 depolymerizing agent and an actin microfilament-depolymerizing agent, mitochondrial recruitment
335 on the BCVs and bacterial growth are inhibited without affecting bacterial entry [44]. However,
336 when used alone, these inhibitors did not impair the mitochondrial recruitment suggesting that two
337 independent mechanisms, i.e. actin-dependent and microtubule-dependent, are involved in this
338 recruitment.

339 To further characterize *W. chondrophila* trafficking, acquisition of ER (Calnexin) and Golgi (GM
340 130) marker were assessed. At 8 h p.i., when bacterial proliferation starts, *W. chondrophila* BCVs
341 are clearly positives for the ER marker but not for the Golgi marker. The *W. chondrophila* BCVs
342 positives for Calnexin or PDI (another ER marker), increase from 4 to 8 h. No close association is
343 observed between BCVs and ER membrane. The *W. chondrophila* inclusions are surrounded by two
344 layers, an inner layer composed of mitochondria and an outer layer composed of ER (**Fig 2A** and
345 **2B**). This results from a sequential process where vacuoles containing *W. chondrophila* first
346 associate with mitochondria and secondly recruit ER.

347 To evaluate the potential role of the COPI-dependent ER-Golgi vesicular transport on the
348 generation of proliferation niche sustaining *W. chondrophila* proliferation, infected cells were
349 treated with BFA which inhibits ARF1, the small GTPase regulating COPI vesicle formation. It was
350 shown that the intracellular growth is inhibited when cells are treated with BFA from 2.5 to 5.5h
351 p.i., supporting the role of the COPI-dependent ER-Golgi vesicular transport during a short period

352 for the biogenesis of the *Waddlia* proliferative organelle [44].
353 In summary, after phagocytosis, the *W. chondrophila* BCVs transiently interact with early
354 endosomes and rapidly evade from the endocytic pathway. Host cell mitochondria are then
355 gradually recruited and form an intimate association with the BCVs. Finally, ER co-localized with
356 vacuoles containing *W. chondrophila*. At 8h p.i., mature *W. chondrophila* proliferative vacuoles are
357 formed with intimate association with mitochondria and ER co-localization (**Fig. 3**). Further
358 experiments are required to shed light on molecular mechanisms involved in the subversion of the
359 endocytic pathway and in the establishment of proliferative vacuoles.
360 Recently, it was reported that *W. chondrophila* possesses a functional catalase (KatA) [46].
361 Catalases are involved in the degradation of reactive oxygen species (ROS) produced in
362 macrophages, by the NADPH oxidase, as microbicidal effectors. KatA is likely a second factor
363 involved in the resistance to macrophage killing in addition to the avoidance of the lysosomal
364 degradation.

365

366 **10. Conclusions**

367 In conclusion, *W. chondrophila* is an emerging pathogen likely involved in human miscarriage and
368 respiratory disease, as well as in bovine abortion. This bacterium is also considered as a potential
369 zoonotic agent. Altogether, the study of the biology of this zoonotic agent is essential to prevent
370 *Waddlia*-associated morbidity.

371 Two aspects of the *W. chondrophila* biology, shared with other *Chlamydiales*, make difficult the
372 study of this bacterium: 1) *W. chondrophila* is an obligate intracellular bacteria; 2) *W. chondrophila*
373 is a genetically untractable organism. Interestingly, *W. chondrophila* is more independent of the
374 host compared to other *Chlamydiales* and is probably a good candidate for growth on cell free
375 medium.

376 *W. chondrophila* possesses a plasmid which suggests that plasmid-based genetic could be possible.

377 In this respect, development of axenic medium (cell free) and transformation techniques would

378 considerably facilitate the experiments and give us the opportunity to better explore the *W.*
379 *chondrophila* biology, a still poorly known landscape. This would significantly improve our
380 knowledge on *W. chondrophila* biology that could be extended to other members of the
381 *Chlamydiales* order.

382

383 **Acknowledgment**

384 The group of Prof. G. Greub is currently supported by the Swiss National Science Foundation
385 grants n° 310030-141050, PRODOC PDF-MP3-127306 and Sinergia CRSII3-141837. We
386 acknowledge C. Kebbi-Beghdadi and A. Croxatto for providing electron microscopy pictures and
387 confocal microscopy pictures.

388

389

390 **References**

391

- 392 [1] Dilbeck PM, Evermann JF, Crawford TB, Ward AC, Leathers CW, Holland CJ, Mebus CA,
 393 Logan LL, Rurangirwa FR, McGuire TC, Isolation of a previously undescribed *Rickettsia* from an
 394 aborted bovine fetus, *J. Clin. Microbiol.* 28 (1990) 814-816.
- 395 [2] Henning K, Schares G, Granzow H, Polster U, Hartmann M, Hotzel H, Sachse K, Peters M,
 396 Rauser M, Neospora caninum and *Waddlia chondrophila* strain 2032/99 in a septic stillborn calf,
 397 *Vet. Microbiol.* 85 (2002) 285-292.
- 398 [3] Chua PK, Corkill JE, Hooi PS, Cheng SC, Winstanley C, Hart CA, Isolation of *Waddlia*
 399 *malaysiensis*, a novel intracellular bacterium, from fruit bat (*Eonycteris spelaea*), *Emerg. Infect.*
 400 *Dis.* 11 (2005) 271-277.
- 401 [4] Baud D, Thomas V, Arafa A, Regan L, Greub G, *Waddlia chondrophila*, a potential agent of
 402 human fetal death, *Emerg. Infect. Dis.* 13 (2007) 1239-1243.
- 403 [5] Baud D, Goy G, Osterheld MC, Borel N, Vial Y, Pospischil A, Greub G, *Waddlia*
 404 *chondrophila*: from bovine abortion to human miscarriage, *Clin. Infect. Dis.* 52 (2011) 1469-1471.
- 405 [6] Baud D, Regan L, Greub G, Emerging role of *Chlamydia* and *Chlamydia*-like organisms in
 406 adverse pregnancy outcomes, *Curr. Opin. Infect. Dis.* 21 (2008) 70-76.
- 407 [7] Haider S, Collingro A, Walochnik J, Wagner M, Horn M, *Chlamydia*-like bacteria in
 408 respiratory samples of community-acquired pneumonia patients, *FEMS Microbiol. Lett.* 281 (2008)
 409 198-202.
- 410 [8] Goy G, Croxatto A, Posfay-Barbe KM, Gervaix A, Greub G, Development of a real-time
 411 PCR for the specific detection of *Waddlia chondrophila* in clinical samples, *Eur. J. Clin. Microbiol.*
 412 *Infect. Dis.* 28 (2009) 1483-1486.
- 413 [9] Baud D, Goy G, Jatton K, Osterheld MC, Blumer S, Borel N, Vial Y, Hohlfeld P, Pospischil
 414 A, Greub G, Role of *Chlamydia trachomatis* in miscarriage, *Emerg. Infect. Dis.* 17 (2011) 1630-
 415 1635.
- 416 [10] Lamothe F, Greub G, Amoebal pathogens as emerging causal agents of pneumonia, *FEMS*
 417 *Microbiol. Rev.* 34 (2010) 260-280.
- 418 [11] Codony F, Fittipaldi M, Lopez E, Morato J, Agusti G, Well water as a possible source of
 419 *Waddlia chondrophila* infections, *Microbes Environ* 27 (2012) 529-532.
- 420 [12] Corsaro D, Greub G, Pathogenic potential of novel *Chlamydiae* and diagnostic approaches
 421 to infections due to these obligate intracellular bacteria, *Clin. Microbiol. Rev.* 19 (2006) 283-297.
- 422 [13] Friedman MG, Dvoskin B, Kahane S, Infections with the *chlamydia*-like microorganism
 423 *Simkania negevensis*, a possible emerging pathogen, *Microbes Infect.* 5 (2003) 1013-1021.
- 424 [14] Greub G, *Parachlamydia acanthamoebae*, an emerging agent of pneumonia, *Clin.*
 425 *Microbiol. Infect.* 15 (2009) 18-28.
- 426 [15] Dilbeck-Robertson P, McAllister MM, Bradway D, Evermann JF, Results of a new
 427 serologic test suggest an association of *Waddlia chondrophila* with bovine abortion, *J. Vet. Diagn.*
 428 *Invest.* 15 (2003) 568-569.
- 429 [16] Koschwanez M, Meli M, Vogtlin A, Greub G, Sidler X, Handke M, Sydler T, Kaiser C,
 430 Pospischil A, Borel N, *Chlamydiaceae* family, *Parachlamydia* spp., and *Waddlia* spp. in porcine
 431 abortion, *J. Vet. Diagn. Invest.* 24 (2012) 833-839.
- 432 [17] Kebbi-Beghdadi C, Batista C, Greub G, Permissivity of fish cell lines to three *Chlamydia*-
 433 related bacteria: *Waddlia chondrophila*, *Estrella lausannensis* and *Parachlamydia acanthamoebae*,
 434 *FEMS Immunol. Med. Microbiol.* 63 (2011) 339-345.
- 435 [18] Draghi A, 2nd, Popov VL, Kahl MM, Stanton JB, Brown CC, Tsongalis GJ, West AB,
 436 Frasca S, Jr., Characterization of "*Candidatus piscichlamydia salmonis*" (order *Chlamydiales*), a
 437 *chlamydia*-like bacterium associated with epitheliocystis in farmed Atlantic salmon (*Salmo salar*), *J.*
 438 *Clin. Microbiol.* 42 (2004) 5286-5297.
- 439 [19] Karlsen M, Nylund A, Watanabe K, Helvik JV, Nylund S, Plarre H, Characterization of
 440 '*Candidatus Clavochlamydia salmonicola*': an intracellular bacterium infecting salmonid fish,

- 441 Environ. Microbiol. 10 (2008) 208-218.
- 442 [20] Moulder JW, Interaction of *Chlamydiae* and host cells in vitro, Microbiol. Rev. 55 (1991)
- 443 143-190.
- 444 [21] Kebbi-Beghdadi C, Cisse O, Greub G, Permissivity of Vero cells, human pneumocytes and
- 445 human endometrial cells to *Waddlia chondrophila*, Microbes Infect. 13 (2011) 566-574.
- 446 [22] Bertelli C, Collyn F, Croxatto A, Ruckert C, Polkinghorne A, Kebbi-Beghdadi C, Goesmann
- 447 A, Vaughan L, Greub G, The *Waddlia* genome: a window into chlamydial biology, PLoS ONE 5
- 448 (2010) e10890.
- 449 [23] Lobry JR, Asymmetric substitution patterns in the two DNA strands of bacteria, Mol. Biol.
- 450 Evol. 13 (1996) 660-665.
- 451 [24] Mackiewicz P, Zakrzewska-Czerwinska J, Zawilak A, Dudek MR, Cebrat S, Where does
- 452 bacterial replication start? Rules for predicting the oriC region, Nucleic Acids Res. 32 (2004) 3781-
- 453 3791.
- 454 [25] Collyn F, Roten CA, Guy L, Solving ambiguities in contig assembly of *Idiomarina*
- 455 *loihiensis* L2TR chromosome by in silico analyses, FEMS Microbiol. Lett. 271 (2007) 187-192.
- 456 [26] Engelberg-Kulka H, Hazan R, Amitai S, mazEF: a chromosomal toxin-antitoxin module that
- 457 triggers programmed cell death in bacteria, J. Cell Sci. 118 (2005) 4327-4332.
- 458 [27] Wang Y, Kahane S, Cutcliffe LT, Skilton RJ, Lambden PR, Clarke IN, Development of a
- 459 transformation system for *Chlamydia trachomatis*: restoration of glycogen biosynthesis by
- 460 acquisition of a plasmid shuttle vector, PLoS Pathog. 7 (2011) e1002258.
- 461 [28] Collingro A, Tischler P, Weinmaier T, Penz T, Heinz E, Brunham RC, Read TD, Bavoil
- 462 PM, Sachse K, Kahane S, Friedman MG, Rattei T, Myers GS, Horn M, Unity in variety--the pan-
- 463 genome of the *Chlamydiae*, Mol. Biol. Evol. 28 (2011) 3253-3270.
- 464 [29] Thomas NS, Lusher M, Storey CC, Clarke IN, Plasmid diversity in *Chlamydia*,
- 465 Microbiology 143 (Pt 6) (1997) 1847-1854.
- 466 [30] Greub G, Raoult D, History of the ADP/ATP-translocase-encoding gene, a parasitism gene
- 467 transferred from a *Chlamydiales* ancestor to plants 1 billion years ago, Appl. Environ. Microbiol. 69
- 468 (2003) 5530-5535.
- 469 [31] McCoy AJ, Maurelli AT, Building the invisible wall: updating the chlamydial peptidoglycan
- 470 anomaly, Trends Microbiol. 14 (2006) 70-77.
- 471 [32] Hatch TP, Disulfide cross-linked envelope proteins: the functional equivalent of
- 472 peptidoglycan in *chlamydiae*?, J. Bacteriol. 178 (1996) 1-5.
- 473 [33] Su H, Watkins NG, Zhang YX, Caldwell HD, *Chlamydia trachomatis*-host cell interactions:
- 474 role of the chlamydial major outer membrane protein as an adhesin, Infect. Immun. 58 (1990) 1017-
- 475 1025.
- 476 [34] Bavoil P, Ohlin A, Schachter J, Role of disulfide bonding in outer membrane structure and
- 477 permeability in *Chlamydia trachomatis*, Infect. Immun. 44 (1984) 479-485.
- 478 [35] Fadel S, Eley A, *Chlamydia trachomatis* OmcB protein is a surface-exposed
- 479 glycosaminoglycan-dependent adhesin, J. Med. Microbiol. 56 (2007) 15-22.
- 480 [36] Molleken K, Schmidt E, Hegemann JH, Members of the Pmp protein family of *Chlamydia*
- 481 *pneumoniae* mediate adhesion to human cells via short repetitive peptide motifs, Mol. Microbiol. 78
- 482 (2010) 1004-1017.
- 483 [37] Kalman S, Mitchell W, Marathe R, Lammel C, Fan J, Hyman RW, Olinger L, Grimwood J,
- 484 Davis RW, Stephens RS, Comparative genomes of *Chlamydia pneumoniae* and *C. trachomatis*,
- 485 Nat. Genet. 21 (1999) 385-389.
- 486 [38] Grimwood J, Stephens RS, Computational analysis of the polymorphic membrane protein
- 487 superfamily of *Chlamydia trachomatis* and *Chlamydia pneumoniae*, Microb. Comp. Genomics 4
- 488 (1999) 187-201.
- 489 [39] Hueck CJ, Type III protein secretion systems in bacterial pathogens of animals and plants,
- 490 Microbiol. Mol. Biol. Rev. 62 (1998) 379-433.
- 491 [40] Plano GV, Day JB, Ferracci F, Type III export: new uses for an old pathway, Mol.
- 492 Microbiol. 40 (2001) 284-293.

- 493 [41] Shea JE, Hensel M, Gleeson C, Holden DW, Identification of a virulence locus encoding a
494 second type III secretion system in *Salmonella typhimurium*, Proc. Natl. Acad. Sci. U. S. A. 93
495 (1996) 2593-2597.
- 496 [42] Haller JC, Carlson S, Pederson KJ, Pierson DE, A chromosomally encoded type III secretion
497 pathway in *Yersinia enterocolitica* is important in virulence, Mol. Microbiol. 36 (2000) 1436-1446.
- 498 [43] Goy G, Croxatto A, Greub G, *Waddlia chondrophila* enters and multiplies within human
499 macrophages, Microbes Infect. 10 (2008) 556-562.
- 500 [44] Croxatto A, Greub G, Early intracellular trafficking of *Waddlia chondrophila* in human
501 macrophages, Microbiology 156 (2010) 340-355.
- 502 [45] Kocan KM, Crawford TB, Dilbeck PM, Evermann JF, McGuire TC, Development of a
503 *Rickettsia* isolated from an aborted bovine fetus, J. Bacteriol. 172 (1990) 5949-5955.
- 504 [46] Rusconi B, Greub G, Discovery of catalases in members of the *chlamydiales* order, J.
505 Bacteriol. 195 (2013) 3543-3551.
- 506 [47] Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger L,
507 Tatusov RL, Zhao Q, Koonin EV, Davis RW, Genome sequence of an obligate intracellular
508 pathogen of humans: *Chlamydia trachomatis*, Science 282 (1998) 754-759.
- 509
510
511

512 **Table 1. Genome comparison between *Chlamydia trachomatis* D/UW-3/CX, *Waddlia***
 513 ***chondrophila* WSU 86-1044 and *Waddlia chondrophila* 2032/99 (adapted from [28])**
 514

	<i>Chlamydia trachomatis</i> D/UW-3/CX [47]	<i>Waddlia chondrophila</i> WSU 86-1044 [22]	<i>Waddlia chondrophila</i> 2032/99* [28]
Genome size	1'042'519	2'116'324	2'141'577
GC content	41%	44%	43%
% coding	89%	92%	93%
Predicted CDSs	895	1934	2'028
Nb of tRNAs	37	37	34
Nb of rRNA operons	2	2	2
Plasmid size	7'493	15'593	-

515 *Unfinished genome
 516
 517
 518

519

520

521

522

523

Table 2. Presence of genes, in *Chlamydia trachomatis* D/UV-3/CX and *Waddlia chondrophila* WSU 86-1044 genome, allowing production of amino acids (adapted from [22]). Genes present are highlighted in green and these absent in red.

Amino acids	<i>Chlamydia trachomatis</i> D/UV-3/CX	<i>Waddlia chondrophila</i> WSU 86-1044
Alanine	Red	Green
Arginine	Red	Red
Asparagine	Red	Green
Aspartate	Green	Green
Cysteine	Red	Green
Glutamate	Red	Green
Glutamine	Red	Green
Glycine	Green	Green
Histidine	Red	Red
Isoleucine	Red	Red
Leucine	Red	Red
Lysine	Red	Red
Methionine	Red	Green
Phenylalanine	Red	Red
Proline	Red	Green
Serine	Red	Green
Threonine	Red	Green
Tryptophan	Green	Red
Tyrosine	Red	Red
Valine	Red	Red

524

525 **Figure legends**

526

527 **Figure 1. Electron microscopy of macrophages infected with *W. chondrophila* at 16h p.i..**

528 A. EBs (arrows) and RBs (arrow heads) present in one inclusion. Scale bar represents 2 μ m.

529 B. Binary fission of RBs. Scale bars represent 1 μ m on the top panel and 0.5 μ m on the bottom panel.

530

531 **Figure 2. Mitochondrial recruitment to *W. chondrophila*-containing vacuoles in Ishikawa cells**

532 **(A). *W. chondrophila* inclusion, in macrophages, surrounded by a bilayer of mitochondria and**

533 **ER (B).**

534 A. Confocal microscopy analysis at 12h p.i.. Mitochondria are stained with mitotracker (in red) and

535 *W. chondrophila* are detected with antibody (in green).

536 B. Electron microscopy analysis at 8h p.i.. Arrows indicate ER. M: mitochondria. Scale bars

537 represent 2 μ m on the right panel and 0.5 μ m on the left panel.

538

539 **Figure 3. Adhesion, internalization and trafficking of *W. chondrophila* during infection of**

540 **macrophages.**

541 1. Adhesion is a critical step for bacterial infection. Several protein families could be involved in
542 this step such as the T3SS, the Omc family, the OmpA family and the pmp family.

543 2. Internalization could be induced by the T3SS effectors translocated directly to the host cell
544 cytoplasm or to the BCV membrane. Once inside the cell, *W. chondrophila* BCVs transiently
545 interact with the early endosome. The catalase KatA degrades ROS produced by the NADPH
546 oxidase in the phagosome lumen.

547 3. After evasion of the endocytic pathway, BCVs quickly recruit mitochondria. This recruitment
548 could be induced by T3SS effectors. Mitochondria recruitment and proliferation are inhibited by
549 nocodazole and cytochalasin B, microtubules and actin filaments inhibitors, respectively.


550 4. The COPI vesicular transport is required to reach the ER, as BFA treatment decrease bacterial
551 proliferation. T3SS effectors could be involved in this step. Finally, *W. chondrophila* proliferate in
552 this particular compartment surrounded by a bi-layer of organelles, an inner layer composed of
553 mitochondria and an outer layer composed of ER.

554 Little is known on the molecular mechanisms involved in adhesion, internalization and trafficking
555 steps during *W. chondrophila* infection. Further investigations on the potential role of the T3SS
556 effectors, the Omc family, the OmpA family and the pmp family in these steps are required to
557 improve our knowledge on *W. chondrophila* pathogenicity.


558

Figure 1
[Click here to download high resolution image](#)


A





B








A



B



-  T3SS
-  Omc family
-  OmpA family
-  Pmp
-  Effectors

1. Adhesion

2. Internalization

