became stronger, but in the course of a few minutes the breathing, which had become very feeble and gasping, with lividity of the face, altogether ceased, and death ensued apparently from a combination of syncope and asphyxia.

Autopsy, twenty-four hours after death.—Brain perfectly healthy. Heart considerably hypertrophied, especially left ventricle, weighing eighteen ounces, somewhat fatty on exterior; no evidence of any valvular disease; right side and both venæ cavæ distended with dark fluid blood, and free from clots. Lungs: Traces of emphysema present, both much congested, especially the left lower lobe. Left pulmonary artery completely blocked by a dark-red, firm, but somewhat friable coagulum of uniform appearance and consistence, which commenced about half an inch on the proximal side of its bifurcation, and was continued on into the lower of its two divisions for nearly an inch; there was no clot in its upper division nor in the right pulmonary artery. Kidneys showed signs of slight commencing granular degeneration. Left leg: The tissues over the inner ankle were considerably thickened, being infiltrated with extravasated blood in process of absorption. The posterior tibial vein (the more internal of the two venæ comites) was filled with a dark-red firm blood-clot, which extended from the ankle to its termination in the popliteal vein. The lower portion of the thrombus was somewhat softer than the upper, and appeared to be partially adherent to the walls of the vein, which, however, showed no signs of inflammation, being merely a little thickened, feeling like a cord when pressed between the fingers. The upper two-thirds was firmer and of a darker colour, lying loose in the vessel, and entirely unconnected with its walls, which appeared to be perfectly healthy. The accompanying vena comes and both saphena veins were quite flaccid and free from thrombus, as also the popliteal, femoral, iliac, and inferior vena cava.

Remarks.—Thrombosis of the veins in the immediate neighbourhood of a fracture, independently of any actual wound of the vein itself, in all probability not unfrequently occurs as the result of the bruising to which the soft parts surrounding the ends of the bones are subjected, the extravasated blood and infiltration of the tissues compressing the veins, and, as a consequence, causing stasis and coagulation of the blood within their walls. In the present instance this condition, which was confined to one of the posterior tibial veins, commenced probably at the ankle (the tissues over the internal malleolus having been severely bruised at the time of the accident), and thence extended along the course of the vessel as far as its termination in the popliteal vein. Here the process appears to have been arrested by the current of blood, and the thrombus probably terminated in a somewhat conical end of considerable size, which, projecting into the popliteal vein, was, after a time, broken off

by the stream of blood, and carried through the heart to the left pulmonary artery, where it was suddenly arrested and found impacted after death. The uniform appearance and consistence of the clot, without indication of any central nucleus, along with the absence of any premonitory symptoms of cardiac or pulmonary obstruction, also point to the fact that the embolus had entirely originated in the popliteal vein, and indicate that when arrested it had suddenly so adapted itself to the shape of the vessel as to at once completely occlude it without giving time for any secondary thrombosis to take place. The absence of the ordinary symptoms of thrombosis—e.g., pain, œdema of the limb, &c.—would, of course, be accounted for by the small size of the obstructed vessel, its deep situation, and by the free maintenance of the venous circulation through the saphenæ, anterior tibial, and accompanying posterior tibial veins.

Complications of fractures such as this do not fortunately often occur, but Dr. H. Tomkins, late house-surgeon here, tells me that a similar accident happened to a patient in the hospital at the commencement of last year. In this case the patient, a woman aged sixty-five years, suffering from a simple fracture of the fibula, died very suddenly on the sixteenth day after the receipt of the injury—without any premonitory symptoms whatever—with well-marked symptoms of asphyxia. On post-mortem examination the heart and other organs were found to be in all respects quite healthy, but an embolus, about two inches in length, was found completely blocking up the right pulmonary artery.

Unfortunately, no examination was made of the condition of the veins of the injured limb, but, in the absence of any other apparent origin for the embolus, it is not improbable that, as in the case I have described above, it became detached from a thrombus which had formed in one of the veins in the immediate neighbourhood of the fractured bone.

HYDATID DISEASE IN AUSTRALIA.

By DAVID THOMAS, M.D., L.R.C.P.

It is well known that Australia presents an extraordinary prevalence of hydatid disease, but, as far as I know, no definite statistics have been published to illustrate the fact. Consequently, some months ago, with the kind assistance of the Hon. W. Morgan, the present Chief Secretary of South Australia, I endeavoured to procure reports from the Governments of Victoria, New South Wales, Queensland, New Zealand, Tasmania, and Western Australia, upon this question. Unfortunately, the mode of registration of the

TABLE OF DEATHS RETURNED AS BEING DUE TO HYDATID DISEASE IN THE PROVINCE OF VICTORIA DURING THE TEN YEARS 1867-77.

Year.	Liver.	Lungs and pleuræ.	Brain and spinal cord.	Heart and Pericardium.	Kidney.	Spleen.	Pancreas.	Omentum.	Ovary.	Womb. (♀)	Abdominal cavity.	Situation not recorded.	More than one organ invaded.	Total annual deaths.
1868	17	5	—	1	—	—	—	—	—	—	2	7	1 (a)	33
1869	6	8	1	—	—	—	1	—	—	—	2	1	3 (b)	22
1870	10	2	1	—	—	—	—	—	—	—	—	4	—	17
1871	4	6	1	—	—	1	—	—	—	1	—	1	1 (c)	15
1872	19	3	1	1	—	—	—	—	—	—	2	2	1 (d)	29
1873	17	3	2	—	—	—	—	—	—	—	—	2	5 (e)	29
1874	21	10	1	—	—	—	1	—	—	—	—	6	2 (f)	41
1875	29	7	—	2	—	1	—	—	1	—	3	2	2 (g)	47
1876	23	1	—	—	2	—	—	—	—	—	1	4	5 (h)	36
1877	20	8	—	1	—	1	—	1	—	—	3	2	2 (i)	38
Totals ...	166	53	7	5	2	3	2	1	1	1 (?)	13	31	22	307 in 10 years.

(a) Lungs and kidneys.

(b) In two instances liver and lungs; in the third case liver and brain.

(c) Lungs and liver.

(d) Lungs and abdominal cavity.

(e) In three cases liver and lungs; in one kidney and abdomen; in another liver, lungs, omentum.

(f) Liver and lungs.

(g) One liver and lungs; the second case lungs and heart.

(h) Three cases liver and lungs; one liver and kidneys.

(i) Liver and lungs.

causes of death in most of these colonies was such as prevented the necessary replies being supplied. However, it appears that in Tasmania no deaths were returned from this disease in the ten years 1867-77. During the greater part of the same period no separate classification of hydatid disease had been adopted in New South Wales; but in 1875 four deaths were attributed to hydatids; in 1876, eleven were so returned. In Victoria, however, the record of deaths from this cause is far more complete, and I append a table based upon the returns from that colony.

One case, in which hydatid of the kidney was present, accompanied by malignant disease of bladder, with stricture, was not returned as a case of hydatid causing death.

During the decade 1867-77, 2.5 per 1000 deaths were due to hydatid disease. In 183 out of the 307 cases the liver was either solely or conjointly with other organs the seat of disease. Holding the second place in frequency come the lungs in 71 cases—i.e., 53 simple and 18 complicated.

Adelaide, South Australia.

A Mirror OF HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum
et dissectionum historias, tum aliorum, tum proprias collectas habere, et
inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

LONDON HOSPITAL.

MR. HUTCHINSON'S CLINIQUE.

FOR the following running notes of some cases lately introduced at Mr. Hutchinson's clinique, we are indebted to Mr. Frederick Treves, F.R.C.S., surgical registrar.

1. *A case of urticaria.*—The patient is a man of twenty-five, who states that he has a "skin disease," and that the eruption appeared only three days ago. It is necessary first to determine the definite form of this skin affection, and then to endeavour to ascertain its cause or its individual peculiarity. The eruption occurs freely on the arms and legs, is less profuse on the trunk, and is evidently a form of urticaria. Has the rash any distinctive characters to mark it? On that point it can only be said that the "wheals" are less noticeable in this case than they are in a typical case of common nettle-rash, and that in typical nettle-rash, on stretching the skin of the affected part, the red elements of the eruption not only entirely disappear, but are replaced by a distinct white counterpart, so that on the stretched skin is seen the eruption faintly but perfectly figured in white instead of in red. In the present patient this feature is scarcely noticeable, but in spite of that there is no obvious reason why this should not be an ordinary case of common nettle-rash. The patient now volunteers the statement that he has the "bad disorder." The statement is vague in the extreme, but before requiring from him a more definite history, its very vagueness may be made a means of suggesting a diagnosis. If the disorder he refers to be gonorrhœa, and the patient be taking copaiba, the rash might be due simply to the well-known action of that drug in occasionally producing urticaria. But suppose the disorder proves to be syphilis, may this rash be syphilitic?

With regard to syphilitic eruptions, let it always be held in mind that there is no one particular form of skin disease to which one can point and say this is always syphilitic. Syphilis has, strictly speaking, no skin disease of its own. Syphilis when it displays itself upon the skin appears there as an imitator—an imitator of the greatest subtlety, capable of representing any form of cutaneous eruption. Syphilis produces an eruption of the skin so like that of variola that over and over again has its true nature been mistaken, and, for a time at least, undiscovered. It imitates the rash of measles, the features of psoriasis, the peculiar action of lupus, and, indeed, the peculiarities of all forms of skin disease. It rarely, however, imitates urticaria, and still more rarely herpes.

On examination, it is found that the patient has gonorrhœa, and that he is taking copaiba. On the frenum pre-

putii, however, is a small scar, and he states that he had a sore in that situation a year or so ago, and that this sore was followed by a "breaking out" on the skin. An examination of his throat reveals nothing positive. A history of syphilis is, however, at least suggested. An examination of the rash reveals nothing that will render the diagnosis certain; treatment, therefore, is postponed, for some days at least, except the negative treatment of discontinuing the administration of copaiba. If in a little while the rash passes off, the case may safely be taken to be one of common nettle-rash, most probably due to copaiba or to some error in diet. If it does not fade in a day or so, the suggestion of syphilis will be rendered more prominent, and the action of specific remedies might be tried.

2. *Syphilitic ulceration of the tongue.*—This patient is a soldier's wife, aged forty-one. She has an ulcer on her tongue, and has been told it is cancer. The tongue being protruded, there is noticed on its left edge a very distinct ulcerating surface, about equal in extent to the surface of a shilling. The ulcer is prominent, raised up, its edges elaborately defined, and its base nearly clean. It presents, however, no exuberant granulations, no warty growths, no clear eversion of its edges. On closer inspection, it is most important to observe that the diseased part is really composed of two ulcers, very close together it is true, but still quite distinct from one another; and on pinching up the part between the finger and thumb, the base of each ulcer is felt to be very thick and very hard, and although so close as almost to be touching, still it is possible to clearly make out that there are two indurated parts, not one. She has had this ulcer for five years. She has a distinct history of syphilis. She has some decayed teeth, presenting sharp and rough projections on the same side of the mouth as the seat of disease. There are no enlarged glands under the jaw. The diagnosis of tertiary syphilitic ulcer proceeding from gum-mata is based, among other points, upon the following, which are selected as affording elements for contrast with epithelial carcinoma: The patient has scarcely reached what might be termed the cancerous age. The ulcer has existed for five years without affecting the glands—a prolonged immunity not to be expected in carcinoma. Above all, there are two ulcers side by side, and evidently having a separate individuality. This is rare in epithelial carcinoma; rare to find, either side by side or less closely connected, two points of origin of malignant growth. Moreover, on feeling the surface of the tongue, the finger encounters here and there a little hard mass about the size of a pea—each one a gumma; and it is hardly necessary to say that one would scarcely find masses of epithelioma dotted about the organ with this pathological generosity. Finally, although this affection is simple enough thus far, it is proper the patient should be warned that it is certainly possible for it to become in the course of time a malignant ulcer, if she neglects its treatment. This warning is given for these reasons:—It is very common, especially about this part, to meet with undoubted carcinomatous ulcers that have emerged from, or have been grafted upon, as it were, long neglected syphilitic sores. Not that there is a structural connexion between the two, or any necessary pathological sequence, but simply because the ulcer has been a cause of prolonged local irritation. This woman is nearing the age when cancer is most frequent. If she neglects treatment, allows the raw surface on her tongue to be constantly irritated by the adjacent teeth, it would be no matter of surprise if the sore should assume sooner or later a malignant aspect, and the glands become involved. If such an event should happen, it would be justifiable to regard this present state as a pre-cancerous period of cancer, so to speak. This pre-cancerous period in the life history of malignant disease is very often indeed to be observed, and is of the greatest moment in influencing early treatment in the removal of sources of local irritation. A middle-aged man came to the hospital a few days ago with a huge mass of ulcerating carcinoma under his jaw. This mass had grown there within the last few months. On his lip was a trivial little pimple, and near it an equally trivial scar. Without especial examination of the lip this little spot would have been overlooked. The patient, however, stated on inquiry that this insignificant little spot had been there for no less a time than ten years; that it was worse at some times than at others; that he would now and then pick it, when it would probably bleed; that it was very sore at times; that he had smoked a great deal, using clay pipes. Had this little spot, really terrible in its insignificance, been examined a few