of the same lung, but at the apex it was distinctly cavernous. The expectoration was nummular and occasionally frothy. The thermometer indicated a temperature of 100° 5' in the evening, and 98° in the morning. In addition to these symptoms, it may be mentioned that there was great weakness, a feeble appetite, and very profuse night-sweats. After allowing her to remain *in statu quo* for five days, she was ordered the hypophosphite of lime in the same doses as in the previous case.

Four days afterwards, on the 22nd of April, she voluntarily stated that she felt much better than she had done for months, and that the night-sweats had disappeared. She was kept on the same treatment six weeks, and the report stated that she was "bright and cheery," and that she had gained a pound in weight since her admission.

On May 21st it was deemed advisable, for the sake of experiment, to stop the hypophosphite and try her with cod-liver oil. On May 24th patient complained of sickness, but no vomiting. She had slight sweating on the previous evening, the first time for fourteen days. On the 26th the sickness was gone, but the night-sweats were profuse. On the 27th it was considered right to order the hypophosphites again in the same doses. The treatment thus embraced hypophosphite of lime and cod-liver oil.

During the next six days the sweating continued, but in a less degree. On June 2nd the patient was much better; no night-sweating. She was weighed on this day, and found to be 8 st. 2 lb., being 6 lb. to the good.

From this date the improvement continued. She went to the Convalescent Home, and on her return from it three weeks afterwards, on August 9th, she stated "she was now perfectly well; cough, spitting, and night-sweats entirely gone; and she had resumed her occupation as an umbrella-maker."

One of the most successful cases was a private patient, A. D.—, aged seventeen. The family history showed that the mother and two sisters had died of consumption, and at no stage of their illness could cod-liver oil be taken without producing nausea and vomiting, however small the dose might be. She had been kept in the house nearly six months, was rapidly losing flesh, sweating greatly at night, and in a highly nervous state through the knowledge of the previous deaths in the family and fear that she was rapidly sinking. She had an irritable, hacking cough, without much expectoration, and, in her own words, stated "she always felt languid, lazy, and tired out, as if it would be no hardship to be in bed all day."

On examination, I found dulness on percussion on the right side, immediately below the clavicle and beside the sternum. This dulness measured transversely and perpendicularly in the former an inch, in the latter an inch and a half. There was no crepitation, but there was bronchial breathing and bronchophony. Without entering into any theoretical inquiry as to causation, there was little doubt that here was consolidation of the lung, limited in character and extent, and all the general indications of phthisis. I use her own words again, because they are plainer and pithier and freer from technicalities than any I could employ. "I sweated a good deal. I wore flannel, and sometimes it was quite damp, and my face and neck so wet that I had to dry them. I did not sleep for some hours after going to bed, and I was very feverish." This was her state when I saw her first, at the end of May. Cod-liver oil and the usual remedies had been tried and abandoned. I ordered her three grains of the hypophosphite of lime twice daily, and also to make careful thermometric observations, which were continued for two months. For the first two nights the temperature was 101° at night and 100° in the morning. For two weeks afterwards the thermometer was found to stand at 100° 5' at night and 99° in the morning. From this time it gradually came down until it was 98° 5' at night and 98° in the morning, and it has continued thus until the present time. With regard to the other symptoms, the sweating stopped the first night and did not recur. The cough became less and less, and in six weeks ceased entirely.

In two months from the time of her commencing the hypophosphite treatment she wrote to me from the country, stating that she often walked six miles in the day. In addition to the hypophosphites she inhaled twice daily a weak solution of ipecacuanha wine.

(To be continued.)

NOTE ON THE USE OF THE FORCEPS.

By THOMAS SAVAGE, M.D., F.R.C.S.,

SENIOR SURGEON TO THE BIRMINGHAM LYING-IN CHARITY.

WHILE I admire and appreciate to the full the work performed by Dr. Johnston during his seven years of mastership of the Rotunda, and published in the *Dublin Medical Journal*, I feel that it would have been better had his remarks in the last report just issued been less strongly expressed in recommending the use of the forceps before the os is fully dilated. Of course, in Dr. Johnston's able hands we know that every care would be used, and that good instead of harm would result; but, in recommending such a practice as a general rule in obstetrics, I cannot help thinking that a door is opened to abuse, and therefore to the production of injurious consequences. Speaking for myself, if I may presume to dictate after so eminent an authority as the Master of the Rotunda, I should feel it to be my duty to be very cautious when setting forth rules of practice for others, especially junior practitioners, and rather to err on the side of timidity than the reverse.

In a large number of cases where forceps are applied before the os is fully dilated a very moderate amount of traction is sufficient to complete delivery; but in a large number also a considerable amount of force will not of itself always liberate the head from the cervix and os, which will require for this end manipulation, and that of a most gentle kind, with the fingers, rather than mere force.

I am very pleased Dr. Johnston lays great stress on the withdrawal of the blades when the head is close to the outlet, as I feel sure a laceration, more or less in extent, is by no means unfrequently produced in the soft parts by either the edges of the blades at the moment of exit, or the too rapid stretching involved; and I have taken for some time the slight trouble to examine carefully the vagina and vulva for a tear after forceps have been used, and I have often found it, notwithstanding great care has been exercised in their use, and with the knowledge of this accident before the practitioner's eyes. Of course I do not apply this liability to the large capacious maternal passages so often met with in pluriparæ as to those which are smaller and more rigid; and I believe the timely removal of the blades has much to do in preventing this accident, while, at the same time, there is less tendency to hæmorrhage if the uterus is not emptied too suddenly, but allowed a short time to contract and expel the child.

Now that the early use of the forceps is advocated by nearly all practitioners, and with so much benefit, all means should be used to make it an instrument as near perfection as possible in its effects, and to this end the formation of any raw surface for the possible absorption of septic matter should be guarded against to the utmost—first, by not tearing the cervix in our perhaps too hasty endeavours to bring down the head; and, secondly, by avoiding any, however trivial, laceration in the soft parts below the cervix.

Birmingham.

A Mirror

OF

HOSPITAL PRACTICE,

BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum, tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

UNIVERSITY COLLEGE HOSPITAL.

REMARKABLE CASE OF PISTOL-SHOT WOUNDS OF HEAD AND CHEST.

FOR the notes of the following case, which presents several points of medico-legal interest, and also for the remarks, we are indebted to Dr. Radcliffe Crocker, resident medical officer.

A young man, aged seventeen, was brought to the hospital on April 4th, at 4 P.M., having shot himself in two places about ten minutes previously. One shot had entered in the cardiac region, the other in the right temple. As