4. That it is necessary to keep the patient on one side, and that the paralysed side should be downwards.
 5. That mucus and other fluids gravitate into, and fill up, the lower lung ; and therefore that if the sides be reversed the mucus will find its way into the opposite lung.
 6. That the fluid, crossing from the large bronchi of one lung to those of the opposite, becomes churned into foam, and causes dangerous obstruction to the respiration.
 7. That the lung is not injured by remaining inactive and filled with mucus for a long period.
 8. That these principles apply to all conditions in which mucus or fluid exists in the lungs, and also to all conditions allied to the apoplectic, whether there be mucus or not.
- Folkestone.

THE ETIOLOGY OF CANCER: STATISTICS AND REMARKS.

By HERBERT L. SNOW, M.D. LOND.,
SURGEON TO THE CANCER HOSPITAL.

IN our present state of helplessness as regards the therapeutic treatment of malignant disease, any information tending to throw light on the origin of that class of maladies may (as a step in some measure towards their prevention) be deemed not unworthy attention—the more especially as I think very undue weight attaches (in the public, and even in the professional, mind) to some causes ; while others are unfairly depreciated. I therefore venture to submit some statistics, taken chiefly from the records of the Cancer Hospital during the past two years ; with the deductions which these seem to me to warrant.

1. *Hereditary tendency.*—Of 146 cases of uterine cancer there was some history of malignant disease having occurred in at least one other member of the family in 12 instances, with 7 very doubtful ditto. Of these 12, 4 patients reported relatives dead of the same disease ; 3 of breast cancer ; 2 of cancer in side (whatever that may mean) ; while in 3 cases the site could not be ascertained. Of 205 cases of cancer of the breast, there was a more or less credible family history in 27 instances ; 6 more are marked doubtful. Of those 27, in 11 cases the mother, sister, or aunt had died of the same complaint (in 4 the mother, in 1 the mother and grandmother) ; 2 were stated to have died of uterine cancer, 2 of hepatic ditto. One relative (father) had died of cancer of the lip ; 1 (grandmother) of cancer of the mouth : these affections being so invariably due to local irritation, I have no hesitation in excluding the cases altogether from the calculation. Again, 1 patient stated that her father had died of cancer of the breast ; considering the rarity of such cases I think her statement must be regarded with doubt. In 13 cases the statement was confidently made that relatives had died of cancer, but the site was unknown. Thus of these latter cases we can only (with some show of reason) regard 15 as occurring in families where the disease might be hereditary. Taking the whole number, we find that in *breast* cancer hereditary tendency occurs in 13·17 per cent. ; but, with the above deductions, in only 7·31 per cent. While in *uterine* ditto the percentage amounts to 8·21. Now, I conceive the only inference we can possibly draw from the above is, that if hereditary tendency to cancer has any predisposing influence whatever in its causation, such influence is of the very slightest, and in practice had much better be ignored altogether, as tending to mislead rather than to assist. Not long since I saw an unmistakably adenoid tumour removed from the breast of a woman, three members of whose family were stated to have died of cancer. I much doubt whether the small percentage recorded above would not be equalled by taking as many hospital patients at random, and inquiring as to the occurrence of cancer in their families.

2. *Injury.*—Of 143 cases of breast cancer, in which special inquiry was made as to preceding injury, 32 patients returned an answer in the affirmative, or 22·37 per cent. ; while 12 gave doubtful replies. Among the uterine cases the idea of injury from external violence is necessarily almost excluded, yet 4 attributed their disease to falls or kicks, while many dated it from a confinement or miscarriage. Among the 205 breast cases quoted above 3 occurred in males, and in these it is noteworthy that 2 attributed their disease, with every appearance of probability, to blows, while the third gave an obscure history of injury. There can be no doubt that

mechanical injury is a real exciting cause of malignant disease ; but (except in epithelioma) the frequency with which the latter so arises is greatly exaggerated, both in medical text-books and in the public estimation. I must here point out one source of fallacy ; women very frequently present themselves, within a week or two after some injury, complaining of severe pain in the breast, and in a state of the greatest alarm ;—and only recover their equanimity after a lapse of many months, coupled with repeated assurances of immunity from cancer. By these cases I am led strongly to consider that in many instances the blow originates the cancer, not *directly* but *indirectly*, by producing mental depression, and by concentrating the attention on the part affected.

3. *Nervous depression (especially mental trouble).*—Of 103 uterine cases in which preceding causes of mental or physical nervous depression (trouble or very hard work) were sought for, 55 gave a history of preceding mental trouble (often combined with hard work), 10 of hard work alone, while 10 gave doubtful replies. Of 38 breast cases 24 patients gave a history of sorrow and anxiety, 4 of hard work, while 1 case was doubtful. Thus of the breast cases (although I must confess the number specially investigated in this connexion is smaller than I could wish), 73·68 per cent. gave an account of nervous depression, either mental or physical, immediately preceding the appearance of the disease, while in the uterine cases the percentage amounts to 63·1. From this I cannot help inferring that sorrow, anxiety, and hard work are very powerful factors in the production of malignant disease. The following case, in its exclusion of every possible cause of cancer save one, I regard as so typical that I cannot forbear relating it here ; it is not included in the above 38. A lady aged forty, wife of a gentleman in comfortable circumstances, strong and previously healthy, leading an easy life, with no family, having a cheerful disposition, without family history of cancer, and never having sustained any injury, became greatly affected at the death of her brother after a short illness, and allowed this to prey on her spirits for some time afterwards. Within a few weeks a small lump appeared in her left breast ; she paid no attention to it till six months afterwards, when, on examination, there was found a well-marked scirrhus tumour, with enlarged axillary glands.

I have above confined my remarks to malignant disease when affecting the breast and uterus, both as affording the largest number of cases for comparison and as practically the most important. I may add that, of 7 cases of rectal cancer, 1 gave a doubtful history of hereditary taint ; 24 of cancer of lips ; tongue, face, or mouth, give 4 instances, and 2 more very doubtful ; 21 ditto of vulva, vagina, and other parts, give 2 instances. Epithelioma, wherever it occurs, is almost invariably (if not always) due to continued mechanical irritation. Nearly all the cases in which the lip, tongue, or mouth were affected could be traced to the irritation caused by a projecting tooth, or by a pipe-stem ; and where these factors were absent the disease often originated at or near the angle of the mouth, where any little fissure would naturally be kept open by the motion of the parts. Two patients with epithelioma of the lip dated their malady from a piece of flesh being torn out during the extraction of a tooth ; one, with epithelioma of the vulva, from a fall on the edge of a piece of board. So-called "ichthyosis" of the tongue, of course, is very apt to end in epithelioma. In 12 out of the 205 breast cases, both breasts became implicated. One case only was supposed to have followed "eczema" of the nipple.

Although I have not yet been able to procure a sufficiently large number of cases for comparison, I submit there is a strong probability that what is here affirmed of mammary and uterine cancer is also true of malignant disease (excepting, of course, epithelioma) wherever that may be situated. And I now beg to recapitulate the principal conclusions to which, I conceive, the foregoing statistics point :—

1. Hereditary tendency, as a predisposing cause of cancer (at all events, of mammary and uterine cancer), is almost valueless, if not entirely so ; and in practical diagnosis should be altogether ignored, as misleading.

2. Mechanical injuries directly produce cancer in a certain percentage of cases ; but this percentage is small.

3. As direct and immediate causes of cancer (especially, in my own experience, of uterine cancer), mental trouble and hard work are very potent agents ; and exert more influence than any other antecedent within our present knowledge.

Bayswater, W.