b. *Folia angusta* ($\frac{1}{4}$ – $\frac{3}{8}$ ad $\frac{1}{4}$ – $\frac{3}{4}$ unc. longa), *marginibus revolutis*.

1. Capitula (4–7) subcorymbosa; pedunculis $\frac{1}{2}$ –1 unc. longis.

51. E. paucidentata (*Steetz*).

2. Capitula solitaria; pedunculis 3–5 unc. longis.

52. E. ciliata (*Benth.*).

Note on the Species of *Nissolia*. By ASA GRAY, M.D., F.M.L.S.

[Read March 15, 1860.]

IN the 'Flora Brasiliensis,' Mr. Benthham has reduced the three species of *Nissolia* proper to one (the *N. fruticosa*, Jacq.), and in a foot-note has briefly characterized a second species, *N. platycarpa*, from Coulter's Mexican collection. The latter is, as I suppose, the plant which, before the fruit was known, I had described under the name of *Chatocalyx Wislizeni**, and afterwards, on receiving the fruit in Wright's second New Mexican collection, had still left in that genus, because the fruit of DeCandolle's genus (at least, of the original species) still remained unknown.

More recently, Dr. Torrey has described and figured another North Mexican species, nearly related to *C. Wislizeni*, but different in several particulars,—among others, in the calyx-teeth, unless these should vary here as they appear to do in *N. fruticosa*. The species then are:—

1. *N. FRUTICOSA*, Jacq.; *Benth. in Fl. Brasil., Legum.*, p. 77, t. 19.

This is No. 194 of Fendler's Venezuelan collection, in fruit.

2. *N. PLATYCARPA*, *Benth. l. c.* Calycis dentibus subulatis tubum subæquantibus; legumine plano longitudinaliter 1-costato, articulis quadratis distinctis, ala samaroidea rectiuscula articulis latiore et 2–3-plo longiore.

3. *N. WISLIZENI*, *A. Gray*. Herbacea, puberula; foliolis rotundatis subreticulatis; calycis dentibus subulatis tubo brevioribus; staminibus diadelphis (9 et 1); legumine plano longitudinaliter 3–5-nervi, articulis quadratis oblongisve distinctis, ala samaroidea parvula sæpius recta articulis parum latiore.—*Chatocalyx Wislizeni*, Gray, Pl. Wright. i. p. 51, and ii. p. 45; Torr. Bot. Mex. Bound. p. 56, t. 18. f. 5–7.

* Although Dr. Gray is quite right in referring his *Chatocalyx Wislizeni* to *Nissolia*, it proves on comparison to be distinct from *N. platycarpa*, both in the venation of the pod and in the terminal wing, which is very much smaller. I have therefore inserted, above, a somewhat fuller diagnosis of my *N. platycarpa*.—G. BENTHAM.

4. *N. SCHOTTII*, *A. Gray*. Fere glabra; foliolis ovatis eximie setaceo-mucronatis; calycis dentibus setaceis tubo longioribus; staminibus inferne monadelphis, superne æqualiter diadelphis; legumine immaturo 2-3-articulato, ala samaroidea anguste oblonga recta brevior.—*Chatocalyx Schottii*, Torr. Bot. Mex. Bound. p. 56, t. 18.

A list of plants observed or collected at Mogador and in its immediate environs during a few days' visit to the place in April 1859; with notes and observations. By the Rev. R. T. LOWE, M.A. Communicated by the SECRETARY.

[Read January 19th, 1860.]

A WEEK'S detention at Mogador, on the coast of Barbary, when returning from Lanzarote to England in the North African Company's Steamer, 'Warrior,' in the spring of the present year, induced naturally some comparison between the general aspect of the sea-coast vegetation in the neighbourhood and that of the Canary Islands and Madeira, in which I had been for some months previously resident; and although losing, by a casual attack of illness, three out of the six days spent on shore, I collected in the immediate environs, partly by my own exertions, and partly by the valuable help of kind acquaintances, a sufficient number, both of the plants and also of the land and sea mollusks, to support the conclusions formed on the spot by observation, and so, perhaps, to warrant an attempt to furnish some slight sketch of the general features of Mogadorian botany and malacology*.

For, beside the interest belonging to a place so little visited, and since the time of Broussonet so wholly unexplored by any naturalist, a peculiar importance is attached to a correct appreciation of the vegetable and zoological conditions of this part of the great African continent. Its intermediate position between the well-searched shores of Algeria in a northerly, and Senegal conchologically, or Sierra Leone botanically, in the opposite quarter of the same continent, no less than its proximity in a nearly easterly direction to Madeira, and in a north-easterly to the two easternmost islands of the Canarian group, Lanzarote and Fuerteventura, the former lying only one degree, and the two latter only two or three degrees south of its own parallel of latitude, mark it as a spot possessing special claims on the attention of naturalists,—bespeaking, I may hope, their favourable acceptance of any con-

* The enumeration of the shells here referred to will be found in the Zoological portion of the present volume.