Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

LOCOMOTOR ATAXY.

BY DAWSON F. D. TURNER, B.A., M.D., &C.

THE following family history may be of interest to those who regard syphilis as one of the important causes of tabes dorsalis. A married couple, now beyond middle life, suffered some years ago severely from syphilis, and the woman had several abortions before giving birth to a living child. Both the man and woman have developed locomotor ataxy in varying degrees; the man suffers principally from ataxia, is unable to balance himself with his eyes shut, is constantly afraid of stumbling and is always looking after his feet, cannot walk in the dark without holding on to something, and has the characteristic gait. The woman has but little ataxia, but suffers severely from lightning pains; she has the Argyll-Robertson symptom, and the loss of the knee-jerk. One of the children has aortic regurgitation, stammers, cannot write steadily, suffers from twitchings of the voluntary muscles and from nystagmus when asked to fix the eyes upon an object, but has no ataxia or loss of the knee jerk. We have here locomotor ataxy occurring in both man and wife, and as the disease is unusual in women, so must it surely be very unusual in both man and wife. It is interesting, too, to note that the manifestations of the disease in the one seem to be complementary as it were to the manifestations of the disease in the other, while the hereditary taint is seen in the child, who has aortic regurgitation and some of the symptoms of Friedrich's disease. The history and the interesting nature of this case seem to me to warrant its being put on record.

Edinburgh.

A CASE OF INTERSTITIAL SALPINGITIS; BOTH TUBES AND BOTH OVARIES REMOVED BY ABDOMINAL SECTION.

BY JAMES OLIVER, M.D., F.R.S. EDIN.

ALICE B—, aged twenty-six, and married seven years, has had one child and two miscarriages. The child was born four years ago, but both miscarriages occurred before this. Menstruation, established at the age of thirteen, had usually lasted four days. For two years after the birth of the child the patient continued to lose blood per vaginam almost every day. During the next eighteen months periodic menstruation was re-established, but the hæmorrhagic discharge continued, not four days as it had done before marriage, but fourteen days, and recurred every fourteenth day. For five months now the hæmorrhage has been almost constant again, just as it was during the two years subsequent to the birth of the child. If the patient is free from hæmorrhage little or no discomfort is experienced; but no matter how frequently the hæmorrhage recurs, pain is complained of during its continuance, and the pain is referred to the right iliac region, the right hip, and down the front of the right thigh as far as the knee. For five months there has been pain whilst voiding urine, and this pain is referred to the hypogastrium. *Physical signs, vaginal examination:* The cervix uteri is directed towards the right, whilst the fundus is deviated towards the left side of the pelvis. In the right fornix and in close apposition with the body of the uterus is felt a small, hard, and regular swelling, which is tender to the touch and appears to be firmly attached to the pelvic floor. The left fornix appears to be free except for the fundus uteri, which can be pressed down on this side. Whilst the patient was under immediate observation, the dates of hæmorrhage were as follows:—Dec. 27th; Jan. 2nd, 4th, 5th, 6th, 7th, 8th, 9th, 13th, 14th, 28th, 29th, 30th, 31st; Feb. 1st, 2nd, 3rd, 4th, 14th, 18th, 19th, 20th, 24th, 25th. Of a total of sixty-three days, there was profuse discharge of blood

during twenty-four, the patient all the while maintaining the recumbent posture. Had this position not been maintained the hæmorrhage would most probably have been more free and more constant, as gravitation would undoubtedly have aggravated the disturbance. On March 6th the abdominal cavity was opened. The right ovary and tube were intimately incorporated with each other, and firmly attached to the pelvic floor; the adhesions, however, were broken down and the tube and ovary were removed together. Although no evidence of disease had been detected per vaginam on the left side, the tube and ovary were here found to be disorganised and matted together, and these, too, were removed. Macroscopically both Fallopian tubes were in a state of fibroid induration, the circumference of the right measuring nearly one inch and a half throughout its extent, whilst the circumference of the left was slightly less than an inch. The lumen of both tubes was patent except at the fimbriated extremity, where it was obliterated, and towards the uterine end where it was occluded by firm bands of constriction. Both ovaries were slightly enlarged and cystic. Sections of both tubes revealed great increase in the connective tissue, but no other evidences of change. No trace of gland tissue was detected in the sections made from the right Fallopian tube, but here and there traces of tubular glands were detected in the sections made from the left tube. In this case there was no history of specific infection as far as one could judge, still it is quite possible that gonorrhœal inflammation may have been the cause of all the inflammatory associations noted in the pelvis. On the other hand, the pelvic peritonitis may have been puerperal and the interstitial change detected in both tubes may have been gradually produced in consequence of an interference with the return of blood from the parts. The patient made a good recovery; and in spite of the fact that both tubes and both ovaries were removed, she has menstruated once since the operation. The discharge was moderate in amount and lasted four days, beginning on April 11th and ceasing on the 14th.

Gordon-square, W.C.

BOWEL OBSTRUCTION CAUSED BY A PIN; REMOVAL OF LATTER; RECOVERY.

BY A. IRWIN BOLTON, A.B., M.B.

ON Oct. 6th, 1890, I was called in great haste to see a Spanish Jew girl, aged twelve years, who was suffering great agony, and twisting about in all directions in her bed. Her mother stated that on the morning of the 4th her daughter said she had swallowed an ordinary pin by accident, but that she made no complaint whatever until the evening of the 5th. She then felt a great desire to stool, but every attempt caused her intense pain and proved abortive. On the afternoon of the 6th I found the patient in great pain. Abdomen very much swollen and slightly tympanitic. Every home remedy had been previously tried. Suspecting that the pin had fixed itself crosswise in the bowel I introduced my finger up the anus, and discovered it as I suspected. The point had penetrated the posterior wall of the bowel, and the head lay against the anterior wall, but on a higher level than the point. Introducing a long forceps up the rectum, I pressed the head of the pin up the bowel, and thereby relieved the point from the posterior wall. Then, catching the point with the same forceps, I extracted it. No further trouble followed. The case shows with what speed a pin can pass all the tortuosities and valves &c. of the bowels without being arrested in its course.

Kustendjie, Roumania.

OBSTETRICAL SOCIETY OF LONDON.—At the meeting of the Society held on Oct. 2nd, the following vote of condolence on the death of Dr. Matthews Duncan was unanimously passed: "The President, Council, and Fellows of the Obstetrical Society of London take the earliest opportunity of offering Mrs. Matthews Duncan and family the expression of their deep sympathy in the great loss they and the profession have sustained in the recent lamented death of their highly distinguished Honorary Fellow, Dr. Matthews Duncan."