INTRODUCTORY.

*Study of Cross-Movements—Special Literature of the District—
Author's Position in 1899—Author's Advance since 1899.*

ELEVEN years have passed since, in 1891, I began my observations in the Dolomites, and in proportion as my geological experience has extended, I have been able to take a fuller grasp of the difficulties of that region. Each new paper that I have published marks a stage in advance of its predecessor; and all the facts obtained by me in the Dolomites lead towards the same general result.

The paper¹ published in 1893 included geological maps of three typical areas in the Dolomites. The map of Dürrenstein mountain (Map C, p. 32) and the sections (l.c. pp. 34 and 35) show that overthrusts and downslips had there taken place with E.N.E.-W.S.W. strike and low inclination towards S.S.E.; and that subsequently the inclined shear-planes and the overthrust and underlaid rock masses had been dragged and displaced by faults directed N.N.E.-S.S.W., N.N.W.-S.S.E., and nearly north-south; the downthrows were on the west of these diagonal and transverse faults, the thrust having acted from E.N.E. (l.c. fig. 9). The map of Falzarego Valley and Cortina (Map B, p. 28) shows a leading fault, curving from W.S.W.-E.N.E. direction to N.E. and N.N.E., and a group of N.N.W., N.W., W.N.W., and west-east faults radial in direction to this curve and intersecting it. In the tectonic chapter I pointed out that the strike of the strata in Falzarego Valley, like the curved fault, veered from an almost east-west direction and dip north to an almost north-south direction and dip west, and that the "bending-round" of the strata had been accomplished by means of these radial faults dislocating an older fault (l.c. p. 98). In the same paper a

¹Miss Ogilvie, "Contributions to the Geology of the Wengen and Cassian Strata in Southern Tyrol," Quart. Journ. Geol. Soc., 1893.

section in Abtey Valley (fig. 5, p. 28) shows that a series of strata has been overcast towards the south and cut by a reverse fault inclined northward, so that whereas the overthrusts at Dürrenstein had been directed towards N.N.W., this was an example of an overthrust towards the south.

In the autumn of 1893, I laid before Professor Rothpletz my geological map of Sella Massive and Buchenstein Valley—the same which was published six years afterwards by the Geological Society of London (see footnote, p. 3). Professor Rothpletz disadvised the publication of this map in 1893 because the observations did not agree with any previously made in the Dolomites, and also because I had not determined whether the N.N.E.-S.S.W. fault on the east of Sella Massive continued across Gröden Pass. The novelty of my observations consisted chiefly in the evidence they afforded that within a comparatively small area a number of folds, overfolds, and inclined planes of overthrust and downslip had been developed along curved directions of strike, or crossing one another, and inclined towards quite different points of the compass. According to my map, the Sasso Pitschi thrust-plane on the south of Sella Massive inclined north, overthrust having taken place towards the south, but on the east side of Sella overthrusts had occurred towards the east, and on the north of Sella overthrust had been towards the N.N.W. These and certain observations of radiating fault-bundles, and of oblique faults cutting older thrust-planes, had more in common with Professor Lossen's work on the Harz mountains¹ than with any Alpine structures then known. My map and sections failed to convince Professor Rothpletz, and as I could give no sufficient explanation of the observed phenomena I agreed it was best to postpone their publication. I limited myself at that time to a brief statement of the chief tectonic and palæontological features, with some diagrams.²

During the next few years, observations of cross-movements began to be obtained in Switzerland. Dr Burckhardt's

¹ Professor Lossen, "Geol. u. Petrogr. Beiträge zur Kenntniss des Harzes," *Jahrb. K. Pr. Geol. Landesanstalt in der Akad.* 1882.

² Miss Ogilvie, "Coral in the Dolomites," *Geological Magazine*, Jan. and Feb. 1894.

work¹ on the Klön and Linth district made known the complex character of the folds that commenced to form in that district during late Eocene or Oligocene time. Dr Burckhardt showed that a system of folds overcast towards the north had developed in the southern part of the district but that in the area of the northern facies two obliquely intersecting fold-systems had been produced. Thus in Switzerland phenomena were described resembling those I had found in the Dolomites.

In 1897, Professor Rothpletz published a preliminary account² of his work on the structure of the Glarus Alps.

When I read these Swiss papers, I took courage to send my maps of Sella and Buchenstein to the Geological Society of London, spending three or four days at Gröden Pass to complete my investigation of the N.N.E.-S.S.W. fault and to confirm my previous observations on the distribution of the igneous rocks in the Pass; then going to Switzerland to see for myself how far my observations in the Dolomites corresponded with Swiss Alpine structures.

In 1898, a preliminary account of my paper on cross-folding in the Dolomites was given in the Proceedings of the Geological Society and in 1899 the full paper³ appeared. It was entitled "Torsion-Structure" in deference to the resemblance between my observations and those to which Professor Lossen had applied the term "torsion."

Immediately after the publication of this paper I contributed to *Nature* (Sept. 7, 1899) a short article indicating the most striking features of general resemblance between the geological structures I had demonstrated in the Dolomites and those in other Alpine regions. In 1899, I also read two special papers⁴ on the causes and effects of crust-torsion, at the British Association Meeting in Dover, and at the Inter-

¹ Burckhardt, "Monographie der Kreideketten zur Klönthal, Sihl, u. Linth," Beitr. zur Geol. Karte der Schweiz, 1896, Neue Folge, V. Lieferung.

² Rothpletz, "Ueber den Geol. Bau des Glärnisch," Zeitschr. d. D. Geol. Gesell. 1897.

³ Mrs Ogilvie Gordon, "The Torsion-Structure of the Dolomites," Quart. Journ. Geol. Soc., 1899.

⁴ Mrs Ogilvie Gordon, "On Sigmoidal Curves," Brit. Assoc. Reports, 1899. "The Crust-Basins of Southern Europe," Intern. Geogr. Congr. Report, Berlin, 1899.

national Geographical Congress in Berlin, showing different aspects of the subject of crust-deformation in relation to local intersecting directions of "strike" either simultaneously or successively induced, and illustrating these from several special areas in Southern Europe.

I purposely selected as the text upon which I demonstrated cross-movements in the Alps, not any area in the Dolomites, but the better known area of Glarus that had been made famous in geology by the researches of Professor Heim. I showed that Glarus was comprised in a local area of subsidence liable to subsidiary movements during or intermittently with the sequence of Alpine movements. I drew a sketch-map of the area that I defined as the Glarus Basin and showed the intersection of different folds and fault-systems, viz., the *leading Alpine strike* at this part directed N.E.-S.W., and the N.W.-S.E. and other *local strikes* following the smaller strike-curves of this Glarus Crust-Basin.

Passing from my illustrative sketch-map of Glarus, I discussed the subject of intersecting faults on general physical principles; showing how the leading strike-curve in any plicated region had reference to some basin of subsidence in the earth's crust, how in one and the same region strike-curves might form with reference to subsiding basins of quite different magnitudes, and that the subsidiary curves relating to the smaller local areas of subsidence were bound to intersect, and be intersected by, the major curves that related to some much wider area of subsidence. I defined the chief local crust-basins which had been determined in the Alpine and Mediterranean region, and demonstrated from my own results in the Dolomites how in all cases local strikes and strike-curves bore relation to local deformational movements initiated in virtue of the local subsidences, relative compression, and different resistances of rock-masses, and that such movements had occurred intermittently with the grander regional movements affecting Southern Europe. While a certain simultaneity of the local and regional movements must have been unavoidable, I emphasised the numerous evidences in the Dolomites that *intensity in the local adjustment movements*, the local inthrows, the local intrusions,

was a characteristic which signalised the decadence period of some important regional movement.

Referring to my proofs that in the Dolomites a prolonged sequence of Tertiary movements had occurred, I led forward numerous evidences that cross-deformation with similar effects of crust-torsion had occurred throughout the Alps, and might be traced (1) to the inevitable *superposition of local and regional strains* in Alpine areas during regional Tertiary movements, seeing that the rock-developments in contiguous Alpine areas exhibited remarkable differences in their physical character; (2) to the superinduced *transversely or obliquely-acting* strains either local or regional in character initiated during a regional movement; (3) to the *periodic variations* in the intensity and horizontal direction of the thrusts causing the grander deformational movements themselves, or those causing subsidiary local movements at special areas, or at certain limited horizons of the crust.

I illustrated from my own maps the leading phenomena due to cross-movements in the Dolomites—such as the curved strikes and unequal thicknesses of the more yielding stratigraphical horizons; radiating fault-groups in the less yielding horizons; faults found to be inclined at varying angles to the horizontal as they are followed from place to place; fault-lines dying out in one direction and the line of throw continuing in a different direction; overfolded masses subjected to subsequent cross-deformation and segmentation, and the segments thrown back at some oblique angle to, or opposite to, the former direction of thrust; intersecting faults initiated at different ages, but along both, or all of which at a still later age differential shearing took place and gave rise to the loop-shaped, sickle or sigmoidal fault-curves, which might be regarded as the most notable tectonic feature in the Alps. In more general terms, I traced the marked effects of crust-torsion to the local deviations produced in virtue of local conditions when any powerful horizontal thrust from a particular direction affects an area that has already been folded and faulted along some other direction or directions of strike,—an area, that has within itself definite lines, localities, and horizons of special crust-weakness

developed in virtue of crust-deformation, and quite additional to the original lines, localities, and horizons of crust-weakness that arise from the natural differences of rock-deposits, and the contiguity of areas with very different types and thicknesses of rock-formations.

In 1898 and 1900, Professor Rothpletz published his important works on the Glarus¹ and Rhaetikon² Alps. Professor Rothpletz showed that folds had formed in curved directions of strike, and had been overcast towards different points of the compass. A main Glarus Thrust-mass had travelled from the east or south-east. On the other hand in the Schild district in the west of Glarus, thrust-masses subsequently travelled from the N.W. direction. East of the Rhine Valley, Professor Rothpletz described a Rhaetikon Thrust-mass which had advanced from the east upon the eastward continuation of the Glarus Thrust-mass. As I reviewed these works in *Nature* (Jan. 24, 1901) it is not necessary to discuss them farther here.

At the close of 1900, Dr Lorenz published a highly interesting geological monograph of an area,³ close to the Rhine Valley, between Glarus and the Rhaetikon. Dr Lorenz determined that cross-folds parallel with two strikes had been developed in this area, a N.W.-S.E. and a N.E.-S.W. strike. He associated the N.W.-S.E. strike with the curve round the north of a *local Glarus crust-basin*; the N.E.-S.W. strike with the Swiss-Alpine strike. Dr Lorenz inclined to the opinion that the determination of the local crust-basin was older than the regional Alpine folding in N.E.-S.W. direction of strike.

A more recent work⁴ by Dr Lorenz treats the stratigraphy of an adjoining district in the Rhaetikon.

¹ Rothpletz, "Das Geotektonische Problem der Glarner Alpen," Jena 1898.

² Rothpletz, "Geol. Alpenforschungen I. Das Grenzgebiet zw. den Ost u. West Alpen u. die Rhätische Ueberschiebung," München, 1900.

³ Lorenz, "Monographie des Fläscher Gebirges," Beitr. zur. Geol. Karte des Schweiz 1900, Neue Folge X. Lieferung.

⁴ Lorenz, "Geol. Studien im Grenzgebiete zw. Helvetischer u. Ostalpiner Facies," II. Theil. "Südlicher Rhaetikon" (Ber. d. naturf. Gesells. zu Freiburg Bd. xii.) 1901.

Special Literature of the District.

The contributions to the geology and the mineralogy of this district have been very numerous. In the course of my researches I have had occasion to refer among the older standard works to these of Baron von Richthofen,¹ Professor Doelter,² Geh. Dr. von Mojsisovics;³ among the newer works to those of Professor Broegger,⁴ Professor Salomon,⁵ Dr Weber,⁶ Herr O. v. Huber,⁷ Professor Romberg.⁸ In July 1902 I published a short account of the sequence of the igneous intrusions in Monzoni and Upper Fassa, based upon my own researches in that district.⁹

Author's Position in 1899.

The sequence of the Triassic rocks in the Dolomites was given in detail in several of my papers; reference may be made to a special paper "On the Upper Cassian Zone" ¹⁰ wherein I pointed out that the horizons of Upper Cassian strata above the *Koninckina Leonhardti* and *Cardita crenata* zone ("Stuores" or Lower-Cassian zone) were those in which the faunas were most local in character and in which the occasional beds of coral reef limestone were most developed. This zone is succeeded in the Sella-Cassian area by a varying thickness of Schlern dolomite and Raibl strata.

¹ V. Richthofen, "Geogr. Karte der Umgegend von Predazzo," 1859.

² Doelter, "Der Geolog. Bau, die Gesteine und Mineralfundstätten des Monzongebirges in Tirol," Jahrb. der k. k. Reichsanst., 1875.

³ V. Mojsisovics, "Die Dolomit-Riffe von Süd-Tyrol u. Venetien," 1879.

⁴ Broegger, "Die Eruptivgesteine des Kristiania-gebietes." II. "Die Eruptionsfolge der Triadischen Eruptivgesteine bei Predazzo in Süd-Tyrol," Kristiania, 1895.

⁵ Salomon, "Geol. u. paläont. Studien über die Marmolata," Paläontographica, 1895, pp. 1-210. "Ueber Alter; Lagerungsform u. Entstehungsart der periadriatischen granitischkörnigen Massen," Tscherm. Min. u. Petr. Mitth. vol. xviii. pp. 109-283.

⁶ Weber, "Die Contactverhältnisse vom Monzonithal nach Allochets." (Inaugural Dissertation) Würzburg, 1899.

⁷ O. v. Huber, "Beitrag zu einer geolog. Karte des Fleimser Eruptivgebietes," Jahrb. der k. k. geol. Reichsanst., 1901.

⁸ Romberg, "Vorarbeiten zur geol.-petrogr. Untersuchung des gebietes von Predazzo (Südtirol) Sitzungsber. der k. pr. Akad. der Wiss., Berlin, 1901.

⁹ Ogilvie Gordon, "Monzoni and Upper Fassa," Geological Magazine, 1902.

¹⁰ Ogilvie Gordon, "Ueber die Obere Cassianer Zone an der Falzarego-Strasse" (Südtirol) Verhandl. der k. k. Geol. Reichsanst., Vienna, 1900. Cf. Geological Magazine, 1900.

The paper published by me in 1899 in the Quarterly Journal of the Geological Society of London was specially devoted to the tectonic problems, and to my mind proved the following propositions :—

(1) The only effusive and contemporaneous igneous rocks originally interbedded with the sediments are the contemporaneous tuffs and lavas in the Wengen-Cassian series.

(2) Large sheets and many small occurrences of augite porphyrite and the amygdaloidal “melaphyre” types are of subsequent age and are intrusive in the sediments as sills and dykes of Tertiary age. My map demonstration of the occurrences of these igneous rocks at the Gröden Pass in intersecting Tertiary faults, and at Col di Lana and in Buchenstein Valley also in Tertiary faults, as a system of dykes and sills, brought forward *the first proof of Tertiary porphyrites and melaphyres* in this region.

(3) The so-called “Buchenstein agglomerates” are not Triassic volcanic products, but shear-breccias or “fault-rock,” sometimes formed from the Wengen mixed tuffs, lavas, and shales, sometimes from Mendola dolomite or Buchenstein calcareous horizons, and sometimes impregnated with ramifications from intrusive igneous material in the fault-zone.

(4) The sedimentary rocks of the Sella-Cassian country were folded by distinct regional crust-movements, an earlier East-Alpine series directed along an almost east-west strike and a later Asta-Judicarian series with a highly complex system of strikes in E.N.E.-W.S.W., N.E.-S.W., and N.N.E.-S.S.W. directions. The sedimentary rocks were during Asta-Judicarian epochs of movement subjected to complicated torsion phenomena, originating definite assemblages of torsion-lines, and faults and overfaults of curious types and effects. Interference phenomena were associated with local deformational movements around masses of more resisting rock such as the Permian Quartz Porphyry, or areas of local subsidence.

(5) The highest horizons of sedimentary rock preserved in the Sella-Cassian districts examined by me are the Jurassic deposits, and they were folded by both movements. In adjoining parts of the Dolomites and in the neighbouring

INTRODUCTORY.

9

Adige Basin, the Asta-Judicarian faults, folds and thrusts have affected all the geological horizons, from Permian to Eocene inclusive, and cannot be older in initiation and development than the late-Eocene, Oligocene-Miocene, and later ages.

Author's Advance since 1899.

During the two summers, 1900-1901, I spent much time in mapping the region lying to the south of the Sella country, as far as the southern slopes of Monzoni, in order to ascertain if the structure of this southern region corresponded with that which I had already worked out to the north. I find that from end to end of this southern district, the geological, petrological, and structural phenomena are the natural expression of those in the Sella-Cassian country, and admit of the same interpretations. There is no evidence that the massive thicknesses of calcareous rock originated as *isolated* reef-growths, either coral or algal—they are marine formations and contain locally coral or algal remains, but also rich molluscan and other faunas; the calcareous rocks are ordinary erosion remnants of a wide sheet of deposit.

The sedimentary succession in the Fassa-Monzoni country is simply the extension of that in the Sella country, with the same lithological and palæontological individualities except in the case of the Wengen-Cassian horizons and the calcareo-dolomitic rocks above. The "Schlern Dolomite" is represented here by a bedded calcareous or calcareo-dolomitic series at whose base there is only a thinly-developed succession of Cassian shaly and impure limestone and of Wengen tuffs, lavas, tufaceous grits, marls, and plant shales. My discovery of Wengen-Cassian strata in this locality for the first time and of a definite fossiliferous horizon in the passage-bands between Upper Werfen strata and the horizon of Mendola Dolomite greatly aided the geological mapping (*cf.* pp. 20, 29).

Owing to the transitional state of the mineralogical nomenclature in the case of the monzonites and other igneous rocks of Monzoni, and to the fact that the best mineralogists in Berlin and Vienna were to revise the mineralogy of Monzoni and Predazzo, I made no attempt to distinguish the fine

gradations of igneous varieties, but concentrated my attention on the relation of the leading types to the deformational lines of structure, and I discovered (*a*) that a very important fault-zone runs through Monzoni on the north slope from Allochet nearly to Malinverno, then bends to S.S.E., this line of fault being parallel with the limiting faults on the north and west of Monzoni; (*b*) that several cross-faults segment the Monzoni massive and are more particularly the seat of the youngest injections.¹

By close attention to the cleavages, faults, and foldings in the different igneous rocks in Monzoni, and by comparison of these structural lines or forms with the structural lines and forms exhibited in the sedimentary strata, and in the augite porphyrite, augite and plagioclase porphyrite, and melaphyres of neighbouring areas, I devised a new method for that country of determining the ages of the intrusive masses and dykes, a method based upon their relation to the succession of deformational movements, of which I had for many years made a special study. The results obtained by this field-method can only be of a general character, and it is scarcely possible to compare them with the results obtained up to the present time along mineralogical lines. Moreover it would take several years to make a complete map of Monzoni; and not until both the cartography and mineralogy are exhaustively investigated in Monzoni and Fassa, can the results be entirely satisfactory. But, while I am thoroughly conscious of the inadequacy of my own individual work, it has at least the merit of demonstrating securely upon stratigraphical and upon structural grounds that the intrusions in Monzoni and Fassa were associated with that long series of Tertiary crust-movements, coming now from one direction, now from another, which gave rise to so many apparent anomalies of structure in the Dolomites.

In delineating the sequence of igneous rocks in Monzoni, I pointed out (*Geol. Mag.*, 1902) the great tectonic importance of distinguishing amongst the varieties of monzonite in Monzoni these biotite and augite varieties, which have always

¹ I made known these discoveries in June 1902 at the Royal Society of London, and they were published in the Abstract (*Geol. Mag.*, July 1902).

been regarded as the "normal type" of monzonite in Monzoni, from certain younger types, chiefly fine-grained, richly felspathic types and quartz-bearing varieties. I showed that whereas the "normal monzonite" had been at the leading fault-zone very much altered, sheared, and slickensided, and intercalated with copious injections of gabbros, diorites, and pyroxenites, the *younger* highly felspathic or quartz-bearing monzonites had been intruded very much later into fault and contact limits as veins or dykes in the gabbro, diorite and pyroxenite series as well as in the *older* monzonites. Their mode of distribution showed that the younger monzonite varieties belonged together with granite, quartz-syenite, syenite, and syenite porphyry intrusions to that advanced epoch in the history of the Judicarian cross-movement when the N.W.-S.E. and almost north-south faults were being developed in Monzoni, and *after* the copious sills of porphyrite had been intruded in the neighbouring area of Fassa.

With regard to the form assumed by the several intrusive masses in the Dolomites, I have throughout my whole series of papers avoided the use of any such term as *laccolite*, *batholite*, etc., as the cases examined by me do not correspond to any of the recognised descriptions of such masses. The forms assumed by the intrusive masses in the Dolomites are so varied that it is impossible to select any one form as a set standard. The igneous rocks have entered fault-zones and horizons of weakness in the sedimentary strata of the earth's crust, and are essentially *dyke-and-sill* systems. Where, as at Col Guschel, Pocol, Col Lares, Varos, and other places the covering strata above a copious sill of porphyrite are partially preserved, these are not gently arched, but are crumpled and tilted, brecciated or schistose, show contact alteration and are penetrated with finer sill-and-dyke ramifications from the larger sill. According to my observations, no large intrusive mass occurs except in the line of some pre-existing fault or faults, and I have found that from a comparatively narrow fault-zone in which the rock-magma has ascended, it has sent sill-ramifications of various sizes here and there into the successive horizons of the crust, the sills ascending both in bedding and cleavage-planes. The sills have spread most

copiously at critical zones of weakness in the crust, more especially in the mixed Upper Werfen Series which had previously been very much split up by inclined fault and cleavage planes, and in the lower and upper limiting-horizons of the thick calcareous deposits, the calcareous rocks becoming locally engulfed or closely threaded with smaller sills and dykes. Sometimes the horizontal extension of the magma has taken place only on one side of a leading fault; *e.g.* at Monzoni, Col del Lares, Chertz Hill, Col di Lana, etc., where the extension of the sills has taken place on the south side of leading east-west faults. This is common in Fassa and Enneberg, because of the predominance of bedding-planes and inclined faults inclined towards north or north-west direction, hence, as seen in my geological sections, many of the sills lie wedge-shaped in the strata. Sometimes horizontal extension of the magma has taken place on both sides of a leading fault, not by any means equally, still in such a way as to give a certain resemblance to a saucer-shaped or cup-shaped intrusive mass, passing downward into one or more narrow fault-dykes; Mt. Donna is a fairly good example. Other local forms, such as the domal sill are called attention to in the course of the paper.

Above all, my previous paper in 1899 and this paper establish the fact that the earlier intrusions ran along the N.N.E.-S.S.W., N.E.-S.W., and E.N.E.-W.S.W. lines of disturbance, as well as the E.-W. or W.N.W.-E.S.E. fault-lines,—in short, that they ascended intersecting faults. And where the magma spread on one side only of two intersecting faults, in an angle as it were, the intrusive mass was afterwards bounded by more or less curved or hook-shaped limits, and these limits were after the intrusion more than ever defined as places of differential strains. My new field-observations bring out more and more clearly the determining influence exercised by pre-existing planes of stratification, cleavage, or faulting, upon the particular local distribution of the igneous magma. It may be said generally that whether connected with longitudinal or cross-faults, the larger sills are limited in their east and west extension by some well-marked flexure of the sedimentary strata associated with the cross-

plication of the rocks; and farther, that such flexures are proximal to or continuous with leading N.N.E.-S.S.W., N.E.-S.W., or E.N.E.-W.S.W. lines of disturbance, many of which were subject to differential movements subsequently to the earlier intrusions.

This all-important structural fact, viz., the intercalation of igneous material in the several Judicarian lines of disturbance and in the series of thrust-masses, as well as in the east-west faults, is really the key to the geological structure of the Dolomites. In the absence of any remnants of Eocene strata in the neighbourhood of Fassa, it alone gives direct evidence of the Tertiary age of the intrusions; for these faults belong to a system that cannot have begun to develop until the late-Eocene or Oligocene Alpine movements, and therefore the relation of the igneous rocks to them shows that the intrusions were at anyrate not earlier than that geological epoch. I brought forward in 1898-99 this evidence for the area of Enneberg and Sella, and now in 1902 am able to establish it for the far grander and, geologically, more widely known occurrences of igneous rock in Fassa and Monzoni.

Briefly stating the more general tectonic features regarding the intrusive rocks, all the massive sheets of igneous rock in the Fassa-Monzoni district are intrusive and were intercalated during the Asta-Judicarian epochs of movement, *i.e.* they are of Tertiary age. The chief intrusions took place at localities where the leading east-west or W.N.W.-E.S.E. faults were intersected by later faults and fault-groups. In the case of copious sill-flows, the igneous magma sought out more especially as its floor these critical horizons or bands of weakness in the crust which on account of the strongly contrasted physical characters of subjacent layers had been the seat of extreme differential strains during the previous crust-disturbances, and had been cut up by innumerable shear-planes. On these planes the torsion of individual rock-segments was slowly effected, and from time to time fresh fractures formed across them, to be followed by later intercalations from a still more specialised magma.

In the Fassa and Monzoni district there are the same evidences as in the Sella country of cross-folding and cross-

thrusting. But now I furnish a mass of new evidence to show how greatly extended in time these movements were, how extremely complex in their deformational effects, and how essentially the history of intermittent intercalations of igneous material was knit up with a long history of local subsidences taking place within the Periadriatic region of the Alps and producing effects which inevitably interfered with the movements of Alpine distribution.

The Fassa and Monzoni district is essentially composed of a series of thrust-masses which represents a dislocated series of overcast folds with steep southern flexure. The fundamental strike of the strata in these thrust-masses is W.N.W., E.S.E. (N. 80° W.), nevertheless there are several leading faults which run due east-west, and appear to be of more ancient date. The stratified rocks of the thrust-masses have been gently folded parallel with the W.N.W.-E.S.E. strike, and dislocated by W.N.W.-E.S.E. strike-faults against which the compression has then been intensified. The strike-faults form a "step" series in Fassa, with downthrow on the north, so that subsequently to the intense epoch of thrust towards the south (or S.S.W.) there took place a certain reversal or relaxation of the north-south strains, and bands of rock slowly subsided in areas *over which previously crust-creep had advanced*.

This folding, step-faulting, and inthrowing of *the thrust-masses parallel with their strike* exercised an important retaining influence upon all subsequent deformational events. It was followed by intensified action of the crust-stresses *transverse or locally oblique to those previously-acting*. On the opposite sides of leading N.N.E.-S.S.W. faults cross-bands of the thrust-masses were vertically displaced, local basins of inthrow were defined, limited by the intersecting faults, and warped fold-systems formed following a curved strike round the south-east, and with steep knee-bent flexures facing the south-east or east. In addition to the leading N.N.E.-S.S.W. faults, fan-like groups of N.N.E., N.E., and E.N.E. faults were initiated, the area of convergence being always located where an older W.N.W.-E.S.E. or east-west fault was intersected by a later transverse fault. The strike of the strata in close

proximity to any of the N.N.E., N.E., or E.N.E. faults shows a series of deflections from an east-west or almost east-west strike to the particular direction of the fault, and the strata are at these faults and at the W.N.W.-E.S.E. strike-faults closely folded and puckered, *but in a distorted way*, the folds like the strike undergoing rapid deflections, varying in form, and owing to the previous displacements affecting quite different horizons of the crust in adjacent localities. The phenomena of deformation attending the development of these fan-like groups of faults, and warped folds and over-folds, were those which I termed characteristic Torsion-phenomena.

During the more advanced stages of the transverse movement, there was, in the immediate proximity of areas of local subsidence of rock-masses, a marked tendency to overthrow of the steep flexures and distorted folds, and the development of reverse and normal fault-planes, followed by the inflow of rock magma into various horizons of the sedimentary mantle of strata. In Rodella and Fassa the chief overthrusting at this period took place from N.N.W., N.W., or W.N.W. direction towards S.S.E., S.E., or E.S.E.; and normal fault-planes with downthrow towards N.N.W. or N.W. were then initiated, some of these having as low an inclination as 15° or 20° , so that the strains were essentially gliding and shearing strains at critical horizons of weakness in the crust. The resisting homogeneous groups of calcareous or calcareo-dolomitic strata were penetrated by systems of fine divisional planes inclined N.W. or S.E., many of which afterwards acted as planes of differential movement.

Over the old thrust-masses now much dislocated and distorted, and embodying invasions of igneous rock, there passed a folding-movement whose general strike was directed N. 40° - 50° E., but was subject to local deflections. It was again followed by overthrusts, downslops, inthrows, and igneous intercalations, having relation to the newer and older directions of crust-fracture and to new local readjustments. As local differential strains transverse or oblique to the N.E.-S.W. strike increased in influence and the axes of elongation or least resistance altered, very curious effects were

produced of back-folding and thrusting towards the directions opposite, or slightly oblique to, the former overfolds and overthrusts. The portions of the older thrust-masses that had in earlier stages swung round towards the N.E.-S.W. direction of strike and been thrust S.S.E. upon planes inclined N.N.W., were naturally specially susceptible of torsional movements at any fresh horizontal compressive stress in the crust. In the course of the N.E.-S.W. folding and subsequent deformation, they were crumpled, faulted and cleaved along N.E.-S.W. strike, and sheared slices locally glided or were carried back on planes inclined towards E.S.E., S.S.E., or S.S.W. In many cases these were gradually nipped laterally from all connection with the rest of the thrust-mass, and left as curious, isolated remnants resting upon low shear-planes, inclined oppositely or obliquely to one another. The shear-planes at a first glance might be very easily mistaken for mere bedding or cleavage planes. The reverse and normal fault-planes inclined S.S.E., S.E., or E.S.E. in the leading examples in this area, *e.g.* the faults round the north and west of Bufaure, Forca Alpe, Varos, Monzoni, are planes of differential movement between rocks of markedly different petrological character and elasticity.

The differential strains transverse to the N.E.-S.W. direction of strike also induced the development of local groups of torsion-faults roughly transverse to the N.N.E., N.E., and E.N.E. fault-bundles and therefore following W.N.W.-E.S.E., N.W.-S.E., and N.N.W.-S.S.E. directions. These have as a rule a very short course, merely from one of the older leading faults to the next on the south. The N.N.W.-S.S.E. lines of faults have sometimes a longer course. There likewise developed a vertical or high-dipping system of cleavage-planes or steep joints in N.N.W.-S.S.E. direction, sometimes almost north-south, well evinced in the igneous and calcareous rocks. A gentle roll of the strata parallel with some of these later fault-lines or parallel with older strike-directions has been subsequently developed in several localities, but this feature is merely an example of the fact that at any or all stages of the crust-deformation local plications were induced in the proximity of

many of the pre-existing fault-lines. The Fassa-Monzoni district gives conclusive evidence of the influence of faults upon forms of deformation. After the fault-lines had developed in diverging bundles and in obliquely and rectangularly intersecting directions, there was little opportunity for the formation of linear and symmetrical folds; the recurrences of horizontal compression only effected partial and unsymmetrical plications following curved strikes determined by local conditions.

A marked feature throughout the whole sequence of deformation was the tendency for certain groups, or limited shear-slices, of the interbedded Werfen marls and marly or granular limestones to be nipped up as short folds, and strongly squeezed and attenuated between down-slipping masses of the calcareous, or in later stages the calcareous and igneous rocks, which were primarily above them in position in the crust. Such groups are preserved as irregular crush layers, lenticles or wedges wholly environed by the rocks into whose midst they were squeezed.

The streaming of the igneous magmas seems to have practically followed the same laws of movement as the rock-strata; the igneous material ascended in the crust at different periods in correlation with local dislocation and subsidence of rock-masses; the igneous inflows, while certain of them retained connection with the deeper-seated reservoirs, were enwrapped as lenticles or wedges of incompletely-consolidated magma amidst the Triassic succession; there undergoing still farther stages of segregation, alteration, consolidation, and deformation; and being repeatedly intercalated with fresh protrusions or threads ramifying in the fissures and cleavages. The crumpling, cleavage, faulting and thrusting followed the same directions in the igneous masses and in the sedimentary strata; but the igneous rocks tended at all surfaces to move differentially in relation to the sedimentary strata, and there were consequently formed those curious igneous and sedimentary breccias which occur here and there throughout the district. At the same time the large intrusive masses acquired schistose, gneissose, or brecciose structure at zones of contact and pressure within themselves.

The fossils which I collected in the field were very numerous, but, except the Wengen-Cassian fossils of Sella Pass, were almost all of them Werfen fossils. These have been identified for me by Dr Broili, Assistant-Custos in Munich Palæontological Museum; and I take this opportunity of thanking him for the kind assistance thus rendered to me. I desire also to thank Mr Gibb, Lecturer in Geology at Aberdeen University, for his careful microscopic examination of a number of thin sections prepared from my collection of igneous rock specimens. Throughout this paper Mr Gibb's descriptions are placed in inverted commas.