IX. *On the Variations in the Internal Anatomy of the Gamasinæ, especially in that of the Genital Organs, and on their Mode of Coition.* By A. D. MICHAEL, F.L.S., F.Z.S., F.R.M.S.

(Plates XXXII.-XXXV.)

Read 3rd March, 1892.

CONTENTS.

	Page
Introductory Observations	281
Bibliography	282
Objects of this Paper	285
Modes of Investigation	286
The Species used	287
The Male Genital System	289
The Mode of Coition	294
The Female Genital System	297
The Alimentary Canal.....	309
Excretory Organs	311
Musculature.....	311
Summary	311
Descriptions of New Species	312
Explanation of the Plates	317

INTRODUCTORY OBSERVATIONS.

THIS paper records the results of a series of investigations extending over the whole of the years 1890 and 1891 and the early part of the year 1892. They have been conducted partly in London, through the kindness of friends, from whom I have received frequent supplies of fresh material from which I could obtain living specimens of the various species I was studying; partly at the Land's End, Cornwall, during an autumnal stay there; partly near Ragaz, in Switzerland, during the summer of 1890; and partly near Innsbruck, in Tyrol, during the summer of 1891. In the two last-named localities I had the advantage of conducting the work jointly with my cousin, Mr. M. J. Michael, who has cut the very numerous serial sections which have formed one of the methods of investigation; he also drew fig. 68. In Tyrol I also had the assistance of Mr. E. Bostock, of Stone, in collecting specimens. I beg to tender my thanks to both those gentlemen.

This paper is not intended as a general treatise upon the anatomy of the Gamasinæ; it is confined, as strictly as may be, to those parts which I believe to be new or to vary from existing records—previously-known organs, which do not vary, being only so far referred to as seems to me desirable in order that the new matter may be clearly under-

stood. The same remarks apply to the Plates; for although many organs are necessarily drawn, particularly in the sections, as to which I have little or nothing new to say, yet it is only in order to show their relation, or position relatively, to those which are new or which do vary from known forms. It will, I hope, be found that there is not any drawing which does not contain something not hitherto known or figured.

The Gamasinæ are the typical subfamily of the Gamasidæ, the other subfamilies being the Uropodinæ, the Dermanyssinæ, and the Pteroptinæ; the two latter subfamilies are wholly parasitic, and their anatomy, so far as it is at present known, is, as might be expected, not so highly developed as that of the other two subfamilies. The Uropodinæ differ considerably in internal organization from the greater part of the Gamasinæ, and their internal anatomy has been dealt with by myself and others, as mentioned below.

Mégnin has claimed that the Gamasinæ are the most highly organized of all the Acarina; and although they are devoid of eyes, which are possessed by several other families, the great development of the so-called brain—*i. e.* the circumœsophageal ganglion, for such it is, all commissures being lost in one mass—the elaboration of the trophi and muscular system, the existence of a heart or dorsal vessel, &c. probably entitle them to that position. They are mostly very active creatures, having fully chitinized dorsal shields and legs, and a ventral surface either entirely or partially covered by chitinized plates, the arrangement and form of which is often characteristic of species or of genera.

It used to be supposed that the Gamasinæ fed upon vegetable matter in process of decay. In the year 1880, however, when investigating the life-histories of some of the species for the purposes of a paper in the 'Journal' of this Society, I found, as there stated, that they would not live in my cells on vegetable matter, but that they thrived admirably on a diet of live cheese-mites (Tyroglyphidæ), and since then I have had frequent opportunities of observing the predatory nature of many of the species. I believe also that Col. Blathwayt, of Bath, who had also been unsuccessful in rearing Gamasinæ upon a vegetable diet, has been perfectly successful over a considerable period since he has adopted the mode of feeding suggested in that paper; he also uses small insects*. Some of the species apparently feed upon small dead insects and other creatures.

A general idea exists that the Gamasinæ are parasitic. This, however, is not by any means universally the case; the greater number of species are free-living, and not parasitic in any stage; some are parasitic in immature stages, but not in the adult. The adult male is very rarely parasitic, and where parasitism exists, even in the young, it is often very doubtful whether they require anything from their hosts beyond conveyance.

The presence of some amount of moisture in the atmosphere and surroundings is essential to the existence of most, if not all, of the species.

BIBLIOGRAPHY.

The anatomy of the Gamasinæ is not by any means a new subject. The memoirs regarding it are few, but some of them are very good.

* "On some common Species of Gamasidæ," *Journal of Microsc. and Nat. Sci.* n. s. vol. ii. (1889) p. 102.

First in order of date must be mentioned the paper by M. Mégnin, of Versailles, in which he expresses the opinion before referred to*. This contains a careful description of the external anatomy and trophi of some of the species of *Gamasus* and *Uropoda*, particularly the latter. The paper also deals, to some extent, with the internal anatomy, but in a far more general and slighter manner, and this portion of the description is far from being entirely correct, particularly that portion which refers to the genital organs—*e. g.*, he attributes to the male the possession of a penis, which does not exist—and the description of the female reproductive organs is still slighter than that of the male; while the information given as to the time and mode of coition was, as will be seen in this paper, entirely incorrect; still M. Mégnin's paper was the beginning of our knowledge of the anatomy of the group, and a large part of the external anatomy especially still stands as quite correct and as the best existing description.

Almost at the same time as M. Mégnin's memoir, Dr. P. Kramer, then of Schleusingen, now of Halle, published a paper on the Gamasidæ †. This paper contains, *inter alia*, considerable information as to the internal organs; chiefly, however, such as might be obtained by a careful study of more or less transparent specimens without actual dissections or sections. The drawings of the internal anatomy are few, and are chiefly confined to outlines of general form, some of which scarcely correspond with anything yet known to me; but the variety of internal organization in the Gamasidæ, and, indeed, in the Acarina generally, is so considerable in different species that I should be sorry to say that they may not be correct. It is in this paper that the existence of a heart in the Gamasidæ was first made known.

In a subsequent paper (1882)‡, which is chiefly devoted to the Uropodinæ, Dr. Kramer gave considerable information as to the various chitinized plates of the external skeleton of the Gamasidæ, but did not add to the then existing knowledge of the internal anatomy.

In a still later memoir § upon *Halarachne Halichæri*, a remarkable parasite belonging to the Gamasinæ, found originally by Prof. Allman in the nares of the Seal ||, the same author carried the knowledge of the internal organization much further; he described and figured the curious tendinous platform in the middle of the body (episternite), which is suspended from the dorsum by muscular bands, and which forms a kind of inner skeleton, whence arise the powerful muscles that move the legs, &c. He also treats correctly of many other matters, which, although described only for the particular species, prove to be of much wider range.

In the following year (1886) Herr Willibald Winkler, with the assistance of Prof.

* "Mémoire sur l'organisation et la distribution zoologique des Acariens de la famille des Gamasidés," Robin's Journ. de l'Anat. et de la Physiol., May 1876, pp. 298-366.

† "Zur Naturgeschichte einiger Gattungen aus der Familie der Gamasiden," Archiv für Naturg. 1876, Bd. i. pp. 46-105.

‡ "Ueber Gamasiden," Archiv für Naturg. 1882, Bd. i. pp. 375-434.

§ "Ueber *Halarachne Halichæri*, Allm.," Zeitsch. für Naturwiss. Bd. lviii. (1885).

|| "Description of a new Genus and Species of Tracheary Arachnidans," Ann. & Mag. Nat. Hist. 1847, vol. xx. pp. 47-52.

Claus, not knowing of Dr. Kramer's paper, independently discovered the existence of the heart in Gamasidæ, and published the result of these investigations in a short but excellent paper *, in which the heart and its accessory organs were described far more exactly than Kramer had done, and which was illustrated by a very good plate, where the surrounding organs, &c., were also shown.

It was possibly his studies for this purpose that drew Winkler's attention to the group; at any rate, in 1888 the first paper was followed by a much more important treatise †. This admirable and beautifully illustrated paper is incomparably superior to anything else which we have upon the internal anatomy of the Gamasinæ; indeed, it may practically be said to comprise all that is, up to the present, recorded on the subject, and it is most clearly explained and carefully figured. The principal types which he takes are what he calls *Gamasus crassipes*, Linn., and *Uropoda scutata*, Koch (?); he, however, less fully refers to and figures *G. fucorum*, De Geer, a nymph of *Pæcilochirus carabi*, Canestrini, and *Hyoaspis nemorensis*, Koch. He also states that he has examined specimens of the genera *Holotaspis*, Kol., *Sejus*, Koch, and *Epicrius*, Can. Still I think the principal stress of the paper must be considered to be laid upon the species called *Gamasus crassipes*. As to this, I have been enabled to follow him, and have frequently dissected this creature, and as a result I wish to express my admiration for the excellence and accuracy of Winkler's work and the beauty of his drawings. I have not myself dissected the other species which he names except *Gamasus fucorum*, and in three genera he does not name the species, so that I cannot say anything about the matter; but I have not any reason to doubt his being as correct with regard to them as with regard to his *Gamasus crassipes*.

It will be observed that I say above, "what he calls *Gamasus crassipes*." I use this expression because it seems to me that the species is not the *Gamasus crassipes* of Linnæus. This is really quite immaterial, and Winkler does not claim to be a specialist in the identification of species of Gamasidæ. I only mention it to avoid confusion in case anybody should at some future time dissect the true species and not find it quite agree with Winkler's description.

When I say that it is not the *Gamasus crassipes* of Linnæus, it must be explained that Linnæus's own description would not enable anyone to identify the species or even the family; it would be equally applicable to a very large number of creatures widely different from each other. Schrank's description in 1781, might enable the student to identify the family, but nothing more. In 1804 Hermann published a description with figures ‡, which are good. This must really be looked on as the first definition of the species; it certainly is not the *Gamasus crassipes* of Winkler.

With regard to the Uropodinæ, I cannot say that I can confirm Winkler's views quite as exactly as I do respecting *G. crassipes*; the larger part of what he says is undoubtedly correct of the species to which he refers, and which again I think is not the *Uropoda*

* "Das Herz der Acariden," Arbeit. d. Zool. Inst. zu Wien, T. vii. (1886) Heft 1, pp. 111-118.

† "Anatomie der Gamasiden," *ibid.* T. vii. Heft 3, pp. 317-354.

‡ 'Mémoire aptérologique,' Strasbourg, 1804.

scutata of Koch, but the *Uropoda ovalis* of the same author—an error which did not arise with Winkler, but is copied by him with a (?) from one of the systematists.

Besides the literature above referred to, there are two memoirs of my own on the internal anatomy of *Uropoda* *. In these I explained in what particulars I was not able to agree with Winkler's description of the anatomy of his species, or perhaps I should rather say what additions I should make to it, and I also showed how widely the internal organization, particularly of the female reproductive organs, varied in different species of the genus.

The Italian works upon the Gamasidæ are chiefly systematic or biological; they contain very little information relative to the internal anatomy.

OBJECTS OF THIS PAPER.

In the paper spoken of in the last section, Winkler, besides treating of the anatomy of his *Gamasus crassipes* (I shall drop the "so called" in future for the sake of brevity) and one or two other species, proceeds to generalize and to take it for granted that other species are similar. I have usually found it unsafe to generalize in matters connected with the Acarina; this probably is partly due to our imperfect knowledge of the group, and partly to the extreme tendency to vary in an unexpected manner which is found on investigation, not only between different families, but even between different species of the same genus when the internal organs are examined. One of the principal objects of this paper is to show that, although, of course, there is a general resemblance between the viscera of the various species of Gamasidæ, which often becomes so marked as to constitute a degree of similarity almost amounting to identity, and although some organs vary but little, yet many most important organs and even systems of the body vary so greatly in different members of the group, and even of the same genus, that whole sets of organs are found in one which are absent in the other, and that the parts which are strictly homologous are often so developed as to be remarkably different from one another; this is especially true of the reproductive organs. It has also been my endeavour to delineate and give definite information with regard to such variations and new organs, which have come under my notice, as appear to me to be of sufficient importance and interest to make them worthy of remark. Again, both on account of its great intrinsic interest and, incidentally to the anatomy, in order, as far as possible, to make certain of the exact functions and mode of employment of some of the organs, a series of observations have been made as to the mode of coition in the species of Gamasids which I have been dealing with. This has not been by any means easy; it has required great care and patience; it is not every species of *Gamasus* which will copulate in confinement under circumstances which render observation possible; and even when such a species is obtained and the requisite surrounding conditions successfully maintained, observation is difficult, and the student must be content to bear with

* "Observations on the special Internal Anatomy of *Uropoda Kramerii*," Journ. R. Microsc. Soc. 1889, pp. 1-15.

"On the Variations of the Female Reproductive Organs, especially the Vestibule, in different Species of *Uropoda*," *ibid.* 1890, pp. 142-152.

many disappointments. The result, however, has been the gaining of considerable information on the point, and I think the process observed is sufficiently curious to make it by no means the least important part of this paper; it is, I believe, totally unrecorded except so far as some portions of it are treated of in my own notice, referred to below, of *Gamasus terribilis*, when I discovered that species, which was the first upon which I made observations that have induced me to follow out the subject.

Finally, there are unrecorded facts tending to prove that in some cases the internal organs are important in the classification of the Gamasidæ by showing relations between subfamilies or genera not previously understood.

Of course it is in the highest degree improbable that in the limited number of species to which my attention has been turned I have found more than a small proportion of the variations existing in this large family of Acarina; doubtless numerous others remain to reward future research.

MODES OF INVESTIGATION.

The methods which I have employed in the present inquiry have been three, viz. 1. Dissection; 2. Serial sections; 3. Observation of the living creature. Of these methods actual dissection has been chiefly employed; I have endeavoured, as far as possible, to confine myself to species where I could obtain an ample supply of specimens; and although I could not altogether succeed without neglecting many interesting cases, yet it has been the rule; and where I have had a sufficient supply I have not confined myself to a few dissections, but have repeated them over and over again, until I had either thoroughly satisfied myself upon the points in question, or else found it impossible to carry the inquiry further by this means. I have not ever relied upon a single dissection. All the figures of whole organs in the Plates illustrating this paper have been drawn from such dissections.

I have found that with these delicate and minute creatures I have not been able to dissect specimens which have been treated with hardening agents such as alcohol or picric acid, or any other of the well-known reagents, as successfully as fresh specimens; I have therefore usually dissected them immediately after death. I at first employed boiling water to kill with, but have subsequently abandoned it in favour of the fumes of chloroform, when the specimen was intended for dissection. Staining has been done either after or during dissection. All dissections have been done by the aid of Stephenson binocular microscopes.

Serial sections, sagittal, horizontal, and transverse, of each species have been employed, both to check the dissections and to trace out points not ascertainable by dissection; here again I have endeavoured, in each instance, to have several series of sections of each species in each direction. The sections have been chiefly cut with a Cambridge rocking microtome. The creatures used for the purpose were either killed in the same manner as those for dissection, or else by boiling after Prof. Lowne's method to solidify the blood-plasma around the organs; some were then treated with dilute picrosulphuric acid, and subsequently passed through alcohol, of gradually increased strength

up to absolute, before imbedding. Latterly, in numerous cases, the picro-sulphuric acid has been omitted, and the creature passed direct from boiling water to dilute alcohol, I think with advantage.

Both dissections and sections have been stained either with picro-carmin, borax carmin, or hæmatoxylin.

The observations on living specimens have been made by the aid of the cells I have usually employed in rearing and observing Acarina, viz. glass rings cemented to ordinary 3×1 in. microscopical glass slips and covered by another similar slip which is kept on by two elastic bands or a clip; the bottom of the cell thus formed is covered with moist blotting-paper, and the whole apparatus is placed on the horizontal stage of a Stephenson microscope for observation.

THE SPECIES USED.

A number of species have been examined of which no record appears in this paper, because I did not find them vary sufficiently from Winkler's type to render it necessary to mention them.

In selecting the species of those that departed far enough from that type I have given the preference to sorts which I could obtain in considerable numbers in one place and in a particular class of material, so that I could get it sent me by country friends when I was in London, or collect them myself when I was in the country, without the great loss of time involved in hunting for solitary wandering species of these active creatures. For this reason *Laelaps stabularis*, Koch, was one of those chosen, because it can be obtained in large numbers in the chaff and fodder in farmhouse stables; my principal supplies of this Acarid came from Cornwall. For the same reason a species which I believe is *Holotaspis montivagus*, Berlese*, was also used because the females of it can be obtained in large numbers in cucumber and other hot-beds and in similar places. Most of my specimens of this species were found and investigated at Ragaz in Switzerland. I, however, did not succeed in finding a single male of this Gamasid, although I was anxious to do so. Another species found with it in the same place, but far less abundantly, was *Holotaspis marginatus*, Herm., of which I obtained one male, but one only. In the other species investigated the males were nearly in as great numbers as the females.

For the same reasons Gamasids which lived in the nests of other animals were very convenient, and indeed were chiefly utilized, not from the variations in organization being in any way connected with parasitism, for the larger number were not parasitic, but only used the nests as a convenient place to live in, or as one that attracted small

* My species closely resembles Berlese's drawing and description; but it is impossible to be certain, because we neither of us found the male, and that is the distinctive sex in Gamasinæ. Although he names his species, he suggests that it was a nymph; mine certainly was not immature, as the female constantly contained a fully-developed egg with the embryo formed and almost ready to emerge (Pl. XXXV. fig. 71). I have drawn the epistome of my species (Pl. XXXII. fig. 18), as it perhaps varies a little from Berlese's plate.

insects upon which they fed, or else were commensalists*. The nest of the mole was most serviceable, because it contained several suitable species in large numbers, and I could almost rely on finding them in any good fresh nest; and because I was able to make arrangements to have fresh moles'-nests sent me at regular intervals all through the winter, and, indeed, whenever I was in London. From this source I obtained the followingspecies, viz.:—1. *Gamasus terribilis*, a large, well-marked, and interesting species, which I first discovered in moles'-nests some few years since†, and which is specially favourable for dissection and section-cutting. 2. *Hæmogamasus hirsutus*, Berlese, at least I believe my creature to be that species, which was found upon the mole, and with which mine agrees thoroughly in size, general appearance, and in almost all other respects; but as the mandible of the male and the epistome vary a little from Berlese's figure, I have given drawings of those organs in my creature (Pl. XXXII. figs. 9, 10, 11), so that if by any possibility there should be two almost similar species, mine may be known; however, the mandible as figured by me differs from Berlese's figure principally in possessing some additional parts, which are very hyaline and might possibly be overlooked; the epistome is a very variable organ. 3. Another species bearing considerable resemblance to *Hæmogamasus hirsutus*, but having well-marked differences both in general appearance and in its organs, the mandible of the male being entirely different; this species is unrecorded; I propose calling it *Hæmogamasus horridus*, and have described it below; it is also drawn (Pl. XXXII. figs. 1 to 5 inclusive). I have taken this as the best type of the most important variations in the female genital organs described below. 4. I have also found and investigated two other species coming from the same nests, both somewhat allied, and both unrecorded; these I propose calling *Laelaps oribatoides* and *Hæmogamasus nidi*; the descriptions will be found below, and are illustrated, as far as necessary, by figs. 6, 7, 12, 13, 14, 15. Another class of nests which provided me with numerous specimens for examination were those of ants; the researches on the myrmecophilous species took place in Tyrol: the species were *Laelaps cuneifer*, Mich., *Laelaps lævis*, Mich., *Laelaps vacua*, Mich., and *Laelaps acuta*, Mich.

Some other species will be found noticed in this paper for special variations or points of interest; but those named above have been the principal species which I have used in tracing out types of organization which seem to me to be common, although to a varying extent, to larger or smaller groups of Gamasinæ.

Before leaving the subject of species I think it would be well to say a few words on the principal points by which species are distinguished in this subfamily, in order that the sufficiency of the details of species which I have given may be understood.

Beyond the general appearance, size, style of motion, and the general distribution of the hairs or spines, these characters are—1. The relative length of the legs with regard to the body, the thickness and shape of the respective legs, and their armature; the two latter points especially in the male. 2. The form of the mandibles of the male where

* "On the Association of Gamasids with Ants," Proc. Zool. Soc. 1891, pp. 638–653.

† "Observations upon a Species of *Gamasus* supposed to be unrecorded," Journ. Quekett Microsc. Club, ser. ii. vol. ii. (1886) pp. 260–268.

the sexes show differences in this respect. 3. The form of the epigynum (or external genital plate) of the female. 4. The number, form, and distribution of the other chitinous plates of the ventral surface in both sexes. 5. The shape of the epistome (the "Randfigur" of Kramer).

Of these numbers 1 and 2 are, I believe, perfectly reliable; number 2 is extraordinarily different in different species, and is often most strange and characteristic in form. Numbers 3 and 4, according to late classifications, are more generic than specific differentiations. Number 5, the epistome, is a more doubtful question, and one not fully settled. Dr. Kramer (*suprà*, p. 283, note †) and Prof. Canestrini * were, I think, of opinion that it was a reliable character. Mégnin said that the epistome was not useful in distinguishing species because it was variable, and was different both in the two sexes and in different individuals of the same sex; Kramer replied that it did not vary so widely as to impair its value as a guide to species. My own opinion is that although it is usually similar in the two sexes, yet it sometimes is markedly different—e. g. *Gamasus terribilis*, *G. crassus*; and also that, although the general characters are usually alike in specimens of the same sex of the identical species, the details and exact form of the organ vary considerably in different individuals, e. g. *Hæmogamasus hirsutus* (see Pl. XXXII. figs. 10, 11); and that, consequently, the epistome is useful in identifying species, but only to a limited extent. It should, I think, always be figured; but too much reliance should not be placed on it; it is only an aid, and where it is known to vary as a secondary sexual character the sex from which it is drawn should be stated.

THE MALE GENITAL SYSTEM.

The first departure from Winkler's type which I will mention is the male genital system of *Gamasus terribilis*. I do not for a moment suppose that the form found in that species is confined to it alone, but I select it as an excellent example, and one which I am exceedingly familiar with.

According to Winkler's description and figure (his fig. 18) (the latter stated to be "schematisch") of the sexual organs of *Gamasus crassipes*, they consist of—1, a single, azygous, globular testis with a trifling swelling on the hinder side, which he considers to be the true spermatoblastic region (Keimlager); 2, two paired vasa deferentia, the length of which is about five diameters of the testis, while their own diameter is not above one seventh that of the testis—thus they are long and thin; they arise quite sharply from the testis, preserve an almost even diameter throughout, and terminate in 3, an azygous ductus ejaculatorius (Ausführungsgang) of about the same diameter as the vasa deferentia and about one fifth the length; 4, a large azygous accessory gland lying between and underneath the vasa deferentia and entering the ductus ejaculatorius close to where it springs from the vasa deferentia. This description I believe to be absolutely correct; I have carefully dissected Winkler's species and agree with him in all respects.

* "Osservazioni intorno al genere *Gamasus*," Atti d. R. Ist. Ven. di Sci. &c. (1881). "Nuove specie del genere *Gamasus*," *ibid.* 1881. 'I Gamasi italiani,' Padua, 1882.

There is not any intromittent organ known in the Gamasinæ; the ductus ejaculatorius communicates directly with the exterior by a roundish opening (Pl. XXXIII. figs. 29, 31, 33, *ga*), which, in most genera, is placed in the median line immediately in front of the sternal plate, and thus in the soft cuticle which is found between that plate and the hypostome, *i. e.* the ventral side of the oral tube (or rostrum). Immediately in front of the genital opening is a curious narrow flap of chitin truncated and terminated by two slightly feathered setæ (figs. 2 & 36, *bp*); it is hinged by its proximal edge to the cuticle, is hollow, and has a fine nerve distributed to it. It is found in both sexes, although the female genital aperture is distant from it. Kramer calls this organ the breast-palpus; Winkler considers it to be the homologue of a labium; I feel doubtful about this homology, and therefore speak of it in this paper by the neutral name of "the bipilate appendage." It is usually found in the Gamasinæ, and does not cover the genital aperture.

The genital organs of the male of *Gamasus terribilis* are delineated by fig. 29; a glance at that drawing will show how widely they differ from those of Winkler's species: that difference is not in any degree a question of time of year or surrounding circumstances; I have examined the organs in large numbers of specimens at all seasons, but invariably with the same result. The unpaired, globular, central testis is entirely absent; in its stead we have two very large doubly-pyriform organs, *i. e.* each testis is more or less of the shape of two pears with their thick ends placed together; but the hinder pear stands somewhat perpendicularly, and leans and curves inward and somewhat forward, while the anterior pear lies nearly horizontally, but curves inward. It must not be supposed that these two pear-shaped pieces are in any way separate organs; they are thoroughly continuous; only there is an external, not very deep, rounded constriction at the point where the somewhat perpendicular part ceases and the horizontal commences. The hinder and upper ends of the two bilaterally symmetrical testes are joined together by a very thin connection near the dorsal surface, just in the situation (as regards the genital organs) where the single testis is in Winkler's species; although in that species, with its long vasa deferentia, the testis is much nearer the anal end of the creature. It will be seen from the drawing that the long, thin, well-marked vasa deferentia of *G. crassipes* are not to be found in *G. terribilis*; indeed, vasa deferentia can scarcely be said to exist at all in the latter species: a minute portion of the organ on each side, close to the ductus ejaculatorius, may be called a vas deferens; but there is not any real demarcation between that and the testis. Of course it might be said that the whole horizontal portion of the organ, on each side, was a vas deferens; but the form of that part, its entire continuity with the hinder part of the testis, the solidity of its contents, and their steadily progressive development from the hinder to the anterior end of what I call the testis seem against that view.

The ductus ejaculatorius and the great accessory gland are practically similar to those of *G. crassipes*, except that the former is perhaps a little longer in proportion, and not quite so uniform in diameter as in *G. crassipes*; and that the latter is without the curious narrow recurved portion at the posterior end figured by Winkler.

Doubtless, in spite of the extreme dissimilarity between the above described organs in

the two species, both are developments of the ring system upon which the genitalia of both sexes in the Acarina, and, indeed, in other groups of the Arachnida, are so frequently based; but this is not saying much, for the varieties which nature gives us in this ring are so numerous and so diverse that they would scarcely seem greater if based upon a separate plan.

Winkler mentions that the distal portion of the ductus ejaculatorius is coated with chitin. In *G. terribilis* this chitinous coating rather assumes the form of a short chitinous trough, not entirely surrounding the mouth of the ductus, but in which the latter lies; while a partial chitinous ring, like one of those in a pseudo-trachea of the blow-fly's tongue, is found within the mouth of the tube; the reason for this difference may possibly be the size of the bodies that have to pass through the ductus ejaculatorius of *G. terribilis*.

It may be as well to say a few words as to the maturation of the sperm-elements in this testis, particularly as it affects later portions of this paper.

In the attenuated end of each testis, from the part where it joins its fellow of the opposite side for a short distance onward, we find, inside the tunica propria, &c., which covers the whole organ, a closely-packed quantity of cells, which at the narrowest part are extremely small and gradually increase in size; but in each cell, wherever it is amenable to the microscope, we see the nucleus quite clear and distinct; although in the smaller cells it is scarcely possible to see a nucleolus. As we examine the testis, proceeding toward the ductus ejaculatorius, the cells gradually and steadily increase in size, until we arrive nearly at the point where what I have spoken of as the more perpendicular portion joins the horizontal part. Near this point the cells, which we may call "sperm-mother-cells," have attained a remarkable size, some of them measure as much as $170\mu \times 125\mu$; each cell still retains a large clear nucleus, which in a cell of the above measurement would be about 58μ and would possess a well-marked nucleolus of about 25μ . Before the mother-cell had attained this size a number of minute, nearly globular bodies, clear but highly refractive, and not at first measuring more than 1μ to 2μ , will have formed in contact with the inner side of the peripheral protoplasm; these I consider to be the true spermatozoa in an early stage. When the cell has attained its full size the interior of the peripheral protoplasm is nearly coated with them; the cell, however, still remains fairly transparent, or at least translucent. The nucleolus has already shown signs of breaking up.

On passing from the perpendicular to the horizontal part of the testis a great change takes place in the sperm-mother-cell, its nucleus breaks up entirely, or sinks into the centre of the cell; the spermatozoa break off from the peripheral protoplasm and collect in a mass in the middle of the cell; this mass is closely agglomerated and no longer transparent, and a clear space is left between it and the peripheral protoplasm; the whole cell now presents the appearance of a partially-filled spermatocyst. The spermatozoa continue to increase slightly in size, until in the ripe cells nearest to the vasa deferentia they may measure 3μ to 4μ ; the sperm-mother-cell itself has, however, shrunk a little, and in a testis where the largest mother-cells were of the above measurement those fully ripe would not exceed about $135\mu \times 89\mu$. These measurements are taken all through from

one freshly killed creature, after examining it with a number of others, so as to assure myself that it was a good average specimen. I may state here, though possibly it belongs rather to a later branch of my subject, that these spermatozoa after being discharged from the mother-cell, and as found in the female, where in this species they occupy the recess in the vagina described by Winkler, have increased in size from what they were in the mother-cell and have become slightly elliptical bodies of about 6μ to 7μ long diameter. There are a very large number of spermatozoa in a mother-cell in this species. The large sperm-mother-cells often adhere in groups of 3 or 4 in the testis, which thus seems divided into separate masses.

In *Laelaps cuneifer*, a very much smaller species, we have a nearer approach to what occurs in Winkler's species: there is a single, unpaired, central testis; but instead of being globular, it is pine-shaped, and lies almost transversely in the body (Pl. XXXIII. fig. 30); from near one of the rounded angles at the base of the pine proceed two longish vasa deferentia, which are of nearly even diameter throughout, and start quite suddenly from the testis (fig. 31). The great accessory gland in this species, although placed as in *G. crassipes* and *G. terribilis*, is different in form (fig. 31, *gla*). It widens out towards its distal extremity and is slightly flattened; but its hind margin is deeply indented, so that the posterior portion of the organ becomes bifurcate, each side being a large rounded lobe. In this species the testis is, as in *G. terribilis*, constituted of large sperm-mother-cells, gradually increasing in size (fig. 32); they are much smaller actually than those of *G. terribilis*, but are large relatively to the size of the species. They are not so tightly packed as in *G. terribilis*, and are considerably different in the average class of form, as may be seen from the figures; the individual spermatozoa are larger in proportion and less numerous, the nucleus seems to disappear at an earlier stage, and the mother-cell is less smooth and transparent.

In *Laelaps lævis* the process which we have seen commencing in the last species by the bifurcation of the hinder portion of the accessory gland has gone vastly further; here we find (fig. 33) not one central gland, but two totally separate elliptical glands, discharging by extremely short ducts, which coalesce only where they enter into the ductus ejaculatorius. The same arrangement is found in *Hæmogamasus hirsutus* and in *H. horridus* (see Pl. XXXV. fig. 72, which is a transverse section cutting the two accessory glands (*gla*) just at the point where the large columnar secreting-cells, which form the hinder parts of the glands, are fading into the looser reticulate tissue found in their anterior portions, and which appears more adapted to the collection and storage of secretion). A longitudinal section through the testis of *Laelaps lævis* is drawn at fig. 34 (Pl. XXXIII.). It will be seen that the formation of the ripe sperm-mother-cells is somewhat different from that in the species above described; the spermatozoa are less distinctly formed, and lie in scattered amœboid masses in a looser and more reticulated substance. This drawing was very carefully made from the actual section, but in consequence of the very small number of specimens which I was able to obtain I could not investigate this species as fully as I should have liked.

What struck me as an interesting case of spermatogenesis, which I believe is novel, in the Acarina at all events, although it may possibly bear some slight analogy to

the mode of development in the Crayfish, is that of *Gamasus crassus*, Kramer; at least I believe that this is the species, but as there is a little difference in the epistome from his figure, I have drawn that organ and the mandible of the male (Pl. XXXII. figs. 20, 19)*.

The process of development in this species is delineated by fig. 35; the different stages are numbered successively; it commences with a single very minute cell (1), which gradually enlarges until the nucleus can be clearly seen (2); this then divides into two (3), then into four (4), eight, &c., in the ordinary manner, the cell continuing to enlarge all the time, so that each contained sphere, after numerous divisions, is as large as one of the original two spheres. In this manner a small nuclear-aggregate (Winkler's "Morula von Kernen") is formed inside and detached from the peripheral protoplasm (5). So far, of course, there is not anything unusual; but now commences what strikes me as singular: the whole cell becomes slightly flattened and more discoidal; four, or more rarely three, slight rounded projections appear on the margin at equal distances (6); these projections continue to increase in length (7), and this goes on until the whole thing presents curiously the form of an *Ophiocoma*, except as regards number of rays (8). The contained spherules arrange themselves chiefly in the arms, and either break up or are no longer distinctly visible; an indentation appears in the edge of the disk between each two rays, which deepens until the four rays are only joined by their points (9). In this manner each ray has become a long tubular spermatocyst with a slightly clavate head; this head, however, becomes indented on one side, and the inner end bends outward so as to form a hook (10). This stage is interesting because it is the permanent form of the spermatocysts, or spermatozoa, in some other species, e. g. *Sejus togatus* (see Pl. XXXIII. fig. 45, which represents a group taken from the testis of the female in that species; those from the spermatheca are similar). Finally the hooked end straightens out; both ends of the cell become pointed, and the whole cell might then, as far as appearance is concerned, well be a single spermatozoon were it not for its size (11); it is in this form that it is found in the vasa deferentia; I am not sure that it does not split longitudinally into two spermatozoa. Certainly the single semen, be it spermatozoon or spermatocyst, is often very much finer and thinner even than 11; it becomes quite filamentous, particularly those found in the female and capsule (see explanation of the word "capsule" in the part on the mode of copulation). I think these filaments are the mature spermatozoa.

Before closing this section it is perhaps well to refer to what Winkler says of the spermatogenesis in *G. crassipes*, in order that by describing the whole process for the sake of clearness I may not be supposed to claim the discovery of anything which he has already observed.

The spermatophores, as he finally calls them, are smaller in his species than in many others, and I believe his description to be quite correct. He says that the primary cells divide and form daughter-cells; the nucleus of the daughter-cell divides frequently and becomes what he terms a morula within an envelope or cyst (Umhüllung); the divided nuclei arrange themselves in a strip; the cell becomes clavate, then lengthens out, but

* Prof. Berlese says that this species is identical with *G. coleoptratorum*; but this is an error on the Italian Professor's part; the two species are very different.

retaining the club-shape; it contains minute highly refractive particles, which are to be regarded as the true spermatozoa, the whole cell being, he considers, a spermatophore, although he usually calls it a spermatozoon; he also thinks that the cell breaks up and is transformed within the female.

THE MODE OF COITION.

I place this here, apparently somewhat out of order, because I think it will enable some part of what I have to say relative to the female organs to be more readily understood.

The coition has been carefully watched by me in four species, viz. *Gamasus terribilis*, *Laelaps cuneifer*, *Hæmogamasus hirsutus*, and *Gamasus crassus*, not only once but in several instances, more especially in the cases of the firstly and thirdly named species.

In 1886, in the paper upon the then newly discovered *Gamasus terribilis* (p. 288, note †), I for the first time described this process, which I think will be found exceptional in nature and of considerable interest. I had, however, then only observed it in a single species and during the short period that elapsed between the discovery and the publication of the paper. I have since investigated the matter more thoroughly and have compared other species, with the result that I am able to describe it much more fully and in some respects more accurately; and that I find that this very singular process is not confined to *Gamasus terribilis*, but is common to at least a section of the family. Of course I cannot say whether it is universal, but it has been the method in all species which I have been able to observe sufficiently well since I first noticed it; but these species are much too few to generalize upon.

I will describe the process in *Hæmogamasus hirsutus*, a species in which I have watched it several times, and have been fortunate in obtaining clear observations and in killing and examining the creatures at different stages.

The male, which is smaller than the female, approaches her usually from the side; he occasionally jumps upon the dorsal plate of the female, but the other is the commoner mode. In either case the male clasps its legs round two of the legs of the female, both on one side; the legs seized have usually been the 3rd and 4th, but sometimes the 2nd and 3rd. The male, having secured a firm hold of the legs of the female, slowly turns over so that his dorsal surface is downward, then slips right under the female, keeping further back than the female, so that nearly half the body of the male projects behind the female; the male passes its fourth pair of legs upward behind the posterior edge of the abdomen of the female and lays its tarsi on the dorsal surface of her abdomen, thus obtaining a firm hold. What looks like a hyaline sac or bubble then appears in the mouth of the genital aperture of the male; this sac is somewhat rapidly distended, but still takes some little time, varying in different cases, before it attains its full size. If the creatures be now suddenly killed (which must be done in an instantaneous manner either by chloroform ejected from a pipette or else by boiling water), this sac, which I propose calling the spermatic capsule (*capsula spermatis*) and shall refer to in future by that name, is found in the mouth of the genital opening of the male, the end of the capsule only being attached to the opening, and the other part of

the capsule standing free in air in a longitudinal or diagonal, nearly horizontal, line between the ventral surfaces of the male and female. The capsule itself is an elongated flask almost parallel-sided and with a rounded distal end; it narrows rather suddenly to a short neck near the genital opening. This capsule, while still small compared to what it will become, but already large in proportion to the creature, is drawn *in situ* (Pl. XXXIII. fig. 36) from a specimen of the male killed at this stage of coition; I have not found sperm-mother-cells in it at this period. If the Gamasids be not killed so soon the capsule continues to distend until it is as long as the body of the male, or even longer; it still preserves nearly its original shape, but is rather broader in proportion to its length than it is at first. If the creatures be killed at this precise moment, which is not very easy to catch, the flask will be found still with its small end in the genital opening of the male; but it will be astonishingly large, and will consist of a tolerably thick but clear and structureless exterior wall surrounding a large number of spermatocysts precisely like the ripe sperm-mother-cells found in the proximal part of the testis, and undoubtedly the same bodies. The capsule is filled up by a clear but thickish fluid, and contains a multitude of minute granules; the small end is closed by gummy and granular matter. A capsule taken in this stage is delineated (fig. 37).

If the creatures be not killed so soon, but are allowed to complete the coitus, the male now exerts both its mandibles and passes them one on each side of the spermatocapsule; the capsule, which apparently has about it viscid material which more or less sets on exposure to the air, adheres to the inner sides of both mandibles, usually to the small brush or row of hairs on each mandible which is inserted close behind the articulation of the movable arm of the chela; the mandibles are now moved away, carrying the spermatocapsule, the small end of which is near the mandibles, but is left free; this small end is applied to the female. The capsule is evidently highly elastic, so much so that it is very difficult to preserve it full; its elasticity usually drives the slight stopping out of the small end and the whole contents of the capsule are discharged upon the slide; when this process takes place in nature of course the coitus is complete. After the contents have been discharged the capsule remains attached to the mandibles; it retains its shape, but has shrunk to about the size shown in fig. 38. If the creatures be killed at any time after the capsule has become attached to the mandibles of the male and before the separation of the two creatures, the capsule will probably discharge if it has not already done so; the empty capsule, still retaining its form but not its size, will be found attached to the two mandibles. Fig. 38 shows the rostrum of a male *H. hirsutus* killed at this stage with the capsule still in the mandibles. Fig. 39 shows the rostrum of a male of *Laelaps cuneifer* killed under similar circumstances. If the creature be not killed at all, the male may be watched cleaning his mandibles, and getting rid of the capsule and other matter which adheres to them.

It will be seen that the form of the capsule in *L. cuneifer* is different from that of *H. hirsutus*, being a much shorter and rounder flask; each form is quite constant in the species to which it is referred; otherwise the process is similar. *Gamasus terribilis* has a capsule also constant in form, but almost the same as that of *L. cuneifer*, for which

reason I have not figured it. In *G. terribilis* the genital plate (epigynum) of the female is lifted by the mandibles of the male and the capsule applied to the vagina; in this species the large spermatocysts are not ever found inside the female, only minute spermatozoa in the recess of the vagina which exists in this species as in *G. crassipes*; the spermatocysts break up inside the capsule, which discharges only minute ovoid spermatozoa imbedded in a thick viscid fluid. The outer walls of the spermatocysts are left in the capsule or about it, and do not enter the female. In four cases of *Gamasus crassus* killed during the coitus, the mandibles of the male were actually inside the vagina of the female. This is a species very like *Gamasus terribilis* in general appearance, but the female, although constructed on the *G. crassipes* type, has not merely the domed recess in the vagina to act as a spermatheca; it has a well-marked, almost globular spermatheca, joined to the vagina by a short chitinized tube with a narrow opening (see fig. 42, *spt*).

The mode of formation of the spermathecal capsule in this species is really astonishing. The lower or movable arm of the chela of the male mandible has a large foramen or hole in it, shaped rather like a lemon-pip (fig. 19); the spermathecal capsule passes right through this hole and folds over. Each end is a sac, the anterior pyriform, the posterior globular; neither sac would pass the hole, but they are joined by a long tubular portion which does pass. It seems to me, therefore, that the capsule must be blown like a bubble right through the hole; the globular end of the capsule emerges last from the genital aperture. Fig. 40 is drawn from a specimen killed *in coitu* and the mandibles then freed from the vagina of the female; the anterior sac of the capsule broken. Fig. 41 shows the posterior end of the capsule emerging from the male genital aperture. This species is one with filamentous spermatozoa. The mode in which the sperm is applied to the female in other species will be discussed in dealing with the female genital organs.

I am of opinion that the more liquid contents of the capsule, other than the spermatocysts, are the produce of the great accessory glands of the male (*gla*, figs. 29, 31, 72); but whether the wall of the capsule itself is the product of a separate gland or of a special part of the main gland and is blown out like a soap-bubble by the rush of liquid and other contents at the moment of formation, or perhaps it is a more apt simile to say as a glass-blower blows a flask, or whether it is simply the hardening of the exterior part of the general mass on exposure to the air, I have not been able to determine; I incline to the former hypothesis in consequence of the regular form which the capsule assumes and its elasticity. If this be so, the formation of the narrow tubular portion of the capsule in *G. crassus* would be easily understood; after the anterior end of the flask has been blown through the foramen in the movable arm of the chela, the mandible might, and probably would be, quickly extruded; the effect of this would be to draw out a long tube of even diameter, as will be readily understood by anyone who has seen glass-blowers draw barometer-tubes by fixing one end of the molten flask and then moving rapidly away.

THE FEMALE GENITAL SYSTEM.

It is in this group of structures that we find the widest variations in the Gamasinæ; we here not only have great differences in the same part in distinct species, but we also meet with a whole set of organs in a large section of the family which are absent from those hitherto described, and these new organs present considerable variety amongst themselves. I will now proceed to describe them, but first for comparison I will shortly take from Winkler's memoir the principal features of the organs as hitherto known and described by him.

The female genital system in *Gamasus crassipes* consists of the following parts, viz. :—

1. An almost globular ovary * fairly corresponding to the single globular testis of the same species; in this ovary the germinal portion is the central, and the more developed eggs are found round the periphery in sacs, the walls of which are so fine as not always to be readily seen. These sacs are temporary outpushings of the outer investing tunic of the ovary, and may be called oöcysts or ovicells.
2. A short unpaired nearly straight oviduct.
3. A larger continuation of the same, with more glandular walls, which Winkler calls the uterus, and where he says the egg attains much of its development.
4. The vagina, which is a largish chamber, open below, but with a domed recess above just over the opening, which recess usually or frequently contains a ball of spermatozoa.
5. Two small vaginal glands which open into the vagina just at the base of this domed recess. And finally
6. The epigynum or external genital plate, which in this species is a large, almost triangular, plate, with its straight edge backward, and is hinged by that edge to the ventral surface; it occupies most of the space between the two hind pairs of legs, and entirely closes the genital aperture. Within this plate is a somewhat elaborate structure of chitinous pieces, the precise function of which is not known.

I will contrast with this the organs which I find in *Hæmogamasus horridus*. I select this species as a type, and as that to be first described, because in it the organs which I believe to be entirely undescribed are found in about their simplest form.

It will be most convenient to commence in the reverse order from that used in regard to *G. crassipes*. The epigynum (genital plate) in this species (fig. 2, *epg*), and indeed in all species of the same genus which I have hitherto examined, is no longer a separate plate working on a ginglymous hinge and closing the genital opening; it is a plate, truncated in front, firmly attached by its whole surface to the ventral cuticle; precisely like the other chitinous plates of the underside of the creature, or probably it would be more correct to say that both it and the others are actually portions of the ventral cuticle, only differing from the other portions by the deposition of chitin in the cells. The genital opening is only a long, somewhat curved, slit placed in the flexible cuticle between the epigynum and the sternal plate; it is, however, somewhat protected by a very delicate and narrow flexible border which runs along the anterior edge of the epigynum.

* I use the word ovary in this paper in the sense in which it is used by all writers on the Acarina, viz., as the whole organ in which the eggs originate, and that solid part in which they are developed in oöcysts (as above defined) before they pass into the free and tubular oviduct.

This class of border, although I do not remember its having been described, is frequently found on the movable genital plates of the Gamasinæ; and is attached to the lower (ventral) part of the edge where the plate is movable, so that it lies under and close against the ventral surface; but it is usually so fine and transparent that it cannot be seen until the plate is lifted.

In *H. horridus* there is not a trace of the chitinous structure of *G. crassipes* inside the epigynum, nor is there any sign of the domed recess which practically functions as a spermatheca in *G. crassipes*: the vagina (Pl. XXXV. fig. 68, *va*) is an extremely simple chamber, with soft and thin walls, leading without vaginal glands or other complication to the oviduct; I have not even found spermatozoa or spermatocysts in it. There is not here any marked difference between uterus and oviduct; the whole appears one organ without distinction of structure or constriction between, unless it be considered that in this and allied species the whole represents Winkler's uterus, and that his oviduct is here sunk in the ovary, which would be a theory requiring some evidence. I have therefore used the expression "oviduct" for the whole (figs. 49, 68, *od*). The oviduct leads to the ovary (figs. 48, 49, 68, *ov*), which is placed near the ventral surface and much as in *G. crassipes*, and may be said to fairly agree with the same organ in that species, except in being rather smaller, more flattened, and less regular in form; it fades into the oviduct without its being easy to say exactly where one commences and the other leaves off, although there is a considerable constriction between the two, through which a tracheal trunk passes. The oviduct runs backward and forms a kind of pocket (Pl. XXXIV. fig. 49).

Now commence the undescribed organs. In the centre of the ovary on its upper surface is a rounded elevation which does not show actual eggs developing; from the sides of this elevation spring two great paired flattened arms or branches; the whole structure, including the root and two arms, is somewhat lyre-shaped, and I propose calling it "the lyrate organ" (*organum lyriforme*)*, and shall so speak of it in future; it is irregular in form, never quite agreeing in two individuals or on the two sides of the same individual, but still preserving the same general form in each species; so that, after becoming well acquainted with them, it would, I think, be possible to tell what species a dissection was from by the form of the lyrate organ alone. This organ is one of the largest and, as regards the arms, one of the most solid in the body; it stains deeply and is wholly composed of largish clearly-marked cells averaging about 30 μ diameter, without any intercellular space or tissue; each cell has a well-defined nucleus averaging about 7 μ diameter, but I have not been able to see nucleoli with certainty. The size and appearance of the cells and nuclei is extremely regular all over the surface of the organ, except just near the root, where the cellulation is lost: if the organ be cut into sections the interior cells are found to be similar to the exterior; the cells are not larger near the root than at the distal ends of the arms, nor are the nuclei more distinct or larger in one place than the other. A portion of one arm of the lyrate organ of a closely-allied species, *Hæmogamasus hirsutus*, is drawn on a larger scale, fig. 47, in order to give an idea of the cellulation, &c. The central rounded elevation from which the arms arise does not exhibit the same cellulation and is not solid—it is hollow, with

* The full name should probably be the lyriform portion of the ovary, but I use "lyrate organ" for brevity.

thick fleshy, but probably slightly distensible, walls. At the point where the arms join this portion the cells appear to separate from each other and become loose. This organ exists in the nymph, but is then very small. When the female is adult, but lately emerged, the lyrate organ, although much larger than in the nymph, is still not so large as it will become; it is when the impregnated eggs are attaining (one by one) the immense size and highest state of maturity which they arrive at within the body of the mother, that the lyrate organ is at its greatest size and development, but the cells of which it is composed are not larger than, nor different from, those in the smaller gland.

With regard to the function of the lyrate organ, I think that it must clearly be regarded as a portion of the ovary. Speaking first of the arms only, they remind me in many respects of the terminal chambers of the ovarian tubes in such Coleoptera as *Platysoma frontale*, *Hydrobius fuscipes*, *Byrrhus pilula*, &c. It may of course be questioned whether they are the true germ-bearing portion, or whether they are composed of vitelligenous cells only; it is not perhaps easy at the present time to answer this question with certainty, as biologists do not appear to be entirely agreed as to the office of what are probably the most nearly allied structures in other creatures. Certain considerations would at first seem to favour an idea of their being yolk-glands: these are *inter alia*, firstly, their large size in creatures which do not appear to lay many eggs, but it is not uncommon in nature for the number of germs in the ovaries to be enormously in excess of the ova eventually matured; secondly, the very large size and high state of development which the egg attains within the body of the mother. In this and allied species only one egg is really matured at a time, and the size and state of maturity which it arrives at before deposition may be judged of from Pl. XXXV. fig. 71, which is a sagittal section carefully drawn to scale from an actual preparation of *Holotaspis montivagus*; in the species I am describing, *Hæmogamasus horridus*, I think the mature egg is even larger in proportion, but the embryo is not so highly matured within the maternal body. There is generally a second, much smaller and flatter, but still large, egg showing a moderately forward state of yolk-division, but not any signs of the embryo forming. The other eggs are usually much less advanced, although in various younger stages.

Notwithstanding the above considerations I do not see any sufficient evidence of the vitelligenous nature of the arms of the lyrate organ, and I think they must, at present at all events, be looked upon as the germiniferous part of the ovary; and this view would bring the various organs of the female genital system most into harmony, and explain in a reasonable and probable manner their relations to each other. As far as I know there are not any structures in any other family of the Acarina at all resembling these lyrate organs, so we cannot be guided by any analogy with other members of the order.

The central domed elevation (*cms*), from the sides of which the arms spring, is, as before stated, unlike them both in histology and otherwise: it is a hollow, shallow recess with fleshy walls, and communicates with the arms, the portion of the ovary in which the oöcysts are developed and the eggs matured, with another organ to be mentioned further on, and probably with the oviduct. It is occasionally empty, but is much more frequently more or less filled with matter which closely resembles that found (as before mentioned) in the spermatid capsule of the male *Gamasus terribilis* at the moment of

coition, and which is discharged from thence into the vagina of the female. Had I been forced to rely upon this resemblance alone I should not have felt confident of the function of the organ, because this minute cellular matter and glutinous fluid are not so distinctive but what other products of the body might be mistaken for them; but luckily a closely allied species, *Hæmogamasus hirsutus*, enabled me to decide the question; it has the domed elevation larger than in *H. horridus*, and usually full of unmistakable filamentous spermatozoa, which often crowd the chamber as full as it will hold. These spermatozoa measure about $30\ \mu$ in length by about $1\ \mu$ in breadth.

It certainly seems strange that in two species so nearly related as *Hæmogamasus horridus* and *H. hirsutus* the spermatozoa should in the first case be minute ovoid bodies, and in the other long and filamentous; it is impossible not to suspect that the latter is the ultimate form into which the former at some time and under some circumstances develops; and this seems all the more probable because in *H. hirsutus* the spermatozoa retain the ovoid form until nearly the last, and it is only when they are about to enter this chamber that they assume the filamentous form. The filament may be formed within the ovoid, but it cannot be distinguished. Although one cannot help being impressed by this obvious idea, yet the facts are that the two species live together in the moles' nests under precisely similar circumstances; that I have examined them at practically all seasons of the year, and have made extremely numerous dissections and sections in all directions of both species, and yet in all cases where the chamber has not been empty it has contained the minute ovoid bodies in *H. horridus* and the filamentous in *H. hirsutus*. It cannot be suggested that they are the same species in different stages; the extremely dissimilar mandibles of the adult males (Pl. XXXII. figs. 3, 9) and the epistomes (figs. 5, 10, 11), to say nothing of the internal anatomy, would put this out of the question; and the females in both have mature eggs.

I propose calling this domed chamber the camera spermatis. It will be seen later on that spermatheca or receptaculum seminis, in the ordinary sense, would scarcely be applicable.

Between the arms of the lyrate organ, but somewhat above and totally detached from them, is another large and wholly undescribed organ, which is found only in the female; there is not a trace of it in the male. It is difficult to give the measure of this because it is dilatable and more or less elastic, and subject to endosmosis and exosmosis; and it is difficult to be sure what is exactly its normal size, if it have any; but in every dissection or section of a mature female *H. horridus* which I have made it has been a large organ, often one of the largest in the body. It is a closed sac or vesicle with hyaline, transparent, but not excessively thin, walls. The form is that of a cone with curved sides and rounded apex, or a bell with a closed mouth (Pl. XXXIV. figs. 48, 49, *sa*). It extends almost to the dorsal surface (Pl. XXXV. fig. 68, *sa*), is more or less horizontal, has the base of the cone (or mouth of the bell) directed toward the rostrum, and the apex toward the posterior end of the Acarid. It is generally slightly coated outside, either wholly or partially, with loose granular matter, but this is not in any way a portion of the organ; it is easily cleaned off, and is to be regarded as embedding matter. I propose calling this organ the sacculus foemineus. Its apex is not permanently closed, it is simply constricted, and

may for the moment be either wholly closed or slightly or considerably open; when open it forms the entrance to a much smaller pyriform sac (figs. 48, 49, *cs*), joined to the first by a short wide neck; or, rather, perhaps the second is to be regarded as a continuation of the first, but the neck between is usually slightly twisted. I propose calling this second sac the "cornu sacculi," as after examining its forms in various species I consider it as in effect a prolongation of the sacculus; the histology is identical. The broader end of the cornu is toward the sacculus, but it turns almost perpendicularly downward (fig. 49), so that its apex is the lowest part; this again is not absolutely closed, only constricted, but the constriction is very close. The apex is attached to the camera spermatis, which it enters low down, practically at the base or from below. The wall of the cornu spreading out after the constriction, and being in fact continuous with the tunica propria of the chamber, thus there is an entrance from the cornu through the camera spermatis to the ovary, although it often requires long and careful examination of many specimens to make sure of the fact; as not only is the constriction close, but the point of junction is small and the parts so fine and delicate that they break with a touch, and nothing but the finest hair can be used in manipulating them.

At the opposite end of the sacculus from the cornu, namely, at the base of the cone, which is the anterior end, there are two branches which I propose calling the "rami sacculi." These rami are bilaterally paired structures, really diverticula of the sacculus, forming wide tubes springing from the lateral edges of the base (figs. 48, 49, 67, *rs*); their length is nearly equal to the width of the sacculus, but they vary in proportion in different specimens, and even on the two sides of the same creature; their histology is identical with that of the sacculus, and their lumen continuous with it without any constriction; they are of about even diameter throughout, and run forward somewhat sideways and a little downward. The distal ends are almost closed and are rounded. From the distal end of each ramus springs a much smaller tube longer than the ramus (figs. 48, 49, *ta*); these I propose calling the tubuli annulati, or ringed tubes. They spring from and open into the rami, without any valve or other closing apparatus that I have yet been able to discover between the two; I mention this because it is not the case in many other species. The histology by no means agrees with that of the rami: they appear to have two tunics, which seem almost, if not quite, detached from each other; the outer, which is the thinner, is much the looser and more flexible, and is distinctly ringed or wrinkled transversely, the ringing being rather irregular; this tunic does not pass within the ramus, but stops at or immediately before the point of junction. The inner tunic seems of a closer and more substantial texture, but neither show cellulation. It must not be supposed from the fact of these tubuli being ringed that they are tracheæ; they never by any chance contain air, and they do often apparently contain fluid; moreover, the ringing is not at all like that of tracheæ; and finally the tracheæ of Gamasids, with which they are abundantly supplied, resemble those of most other Acarinæ in not presenting an appearance of ringing or spiral threads, although they might probably be made to break up into spirals by properly applied pressure as in the Oribatidæ. The ringed tube on each side commences, as before stated, at the termination of the corresponding ramus, and it ends, with a more or less trumpet-shaped mouth, in the thin

connective cuticle which binds the coxa of the third leg to the more chitinous part of its acetabulum.

Thus it will be seen that this series of organs or parts—the cornu sacculi, the sacculus foemineus, the rami sacculi, and the tubuli annulati, all of which are strictly confined to the female—form an unbroken connection, varying enormously in size and capacity, but absolutely continuous, between the sperm-chamber in the ovary and the acetabulum of the third leg on each side.

The position of these organs as regards the other viscera may be well gathered from fig. 46, which was carefully drawn immediately after death from a female of *Laelaps lævis*, after the dorsal plate, &c., had been removed.

It will, I think, be convenient to describe some at least of the somewhat numerous and wide variations of these organs which I have found in different species before treating of their contents and supposed function.

In describing all these parts, and, indeed, the internal anatomy generally, I cannot too strongly insist upon the fact that two specimens are never really alike, and that the same specimen varies with age and condition; it is therefore only what may be called a type-form in each case that can be described; but this type is never so widely departed from but what it may be easily recognized in any dissection of the species, and may probably therefore be considered as constant as could be anticipated.

Hæmogamasus nidi (Pl. XXXII. figs. 6, 7) is a smallish new species allied to *H. horridus*, and in which the female genital system is so similar that it is not necessary to give a separate description of it.

Laelaps stabularis is a species in which the new organs above described correspond more closely to those of *H. horridus* than in any other I have found, except *H. nidi* (see Pl. XXXIV. fig. 51). The differences, as will be seen by the figure, are that the lyrate organ of *L. stabularis* is much more enlarged at its distal ends than that of *H. horridus*, and forms broad, somewhat recurved plates; that the cornu is wider in the lumen and much less plainly marked off from the sacculus; and that the rami are much longer, and the ringed tubes consequently very much shorter, than in *H. horridus*.

Hæmogamasus hirsutus is a species closely allied to *H. horridus*, belonging to the same genus, and found with it under similar conditions; and yet the organs we are discussing differ far more widely from the same parts in *H. horridus* than those in *L. stabularis* do, although the latter is a very different creature, belonging to a separate genus. The lyrate organ of *H. hirsutus* (fig. 52) does not vary in any remarkable manner from that of its ally, but when we examine the sacculus (fig. 53, *sa*) the contrast is very striking. Instead of the large, simple, bell-shaped sac we have a small, more or less rhomboid, organ placed transversely across the body; the whole central part of the anterior side of this is occupied by a shortish but very wide cæcal prolongation running straight forward between the rami. From the posterior corners of the sacculus proceed two paired, almost globular, diverticula, immensely larger than the sacculus itself, and joined to it by almost imperceptible necks, with sharp lines of demarcation where they issue from the central portion of the organ. These diverticula extend almost to the dorsal surface and are quite clear and hyaline; the position of one of them is shown in Pl. XXXV. fig. 70, *sd*.

Between the diverticula the cornu arises; it is smaller than in *H. horridus*, but of somewhat similar shape, except that it is generally joined to the sacculus by a small, almost globular chamber with a narrow neck. The rami are short but wide; and the ringed tubes much of the same nature as those before described, but they do not terminate exactly where they reach the rami with a simple opening into the latter, as in *H. horridus*; they enter the rami and continue within for a minute distance, then each ringed tube terminates in a very small, almost globular enlargement within the ramus (fig. 54); this sphere is so clear and delicate that it can only be seen under favourable circumstances and with a highish power, but it certainly exists. I have not been able to determine to my own satisfaction whether this and similar, or more or less similar, terminations of ringed tubes in other species are really permanent closings, or only in the nature of valves; the latter is the more probable, although I have not hitherto detected any openings, but they are so small and fine that proper examination is almost impossible, and I have not been able to trace them in the sections.

Laelaps oribatoides (fig. 56) is another allied species much of the *H. hirsutus* type, but with considerable differences. The two globular diverticula are present, but in this species are smaller than the sacculus, while the central anterior diverticulum of the last species is entirely absent: the cornu, however, in *H. oribatoides* assumes much greater importance; it is an organ containing nearly as much space as the sacculus and winds over and round the lyrate organ down to the underside of the camera spermatis, where it enters. The lyrate organ in this species is singular in form, the distal ends being much enlarged and forming irregular hatchet-shaped pieces, which are much reflexed.

Laelaps vacua is a very small species; in it we find a widely different development of the sacculus organs (fig. 57); that viscus itself is a small globular sac without rami; the ringed tubes enter the sacculus itself and project within it, with a slight bulb, which I do not think is closed; the cornu is a large, more or less discoidal or ovoid body, not regular in form, but considerably larger than the sacculus, to which it is joined by a long fine tube, larger than the ringed tubes, but not much so; the difference in the figure is a trifle too great. This tube springs sharply from the sacculus and from the cornu, with a well-marked line of demarcation; it is shown stretched out in the figure, but *in situ* it winds round the other organs, and is slightly embedded in fatty and loose cellular matter. The cornu lies with a portion of its substance almost embedded in the camera spermatis and ovary. I could not, however, trace the entrance in this species, although it probably exists; I had not a sufficient number of specimens for investigation, and the creature is very small and delicate (extreme length about .5 mm.).

Fig. 58 represents the sacculus and cornu of a small species very like *L. vacua* externally, and with these organs similar in principle.

Laelaps acuta is a smallish and very swift species, which again presents novelties in the set of organs we are considering; it has an elliptical sacculus (fig. 59, *sa*), without any rami: the ringed tubes enter the sacculus itself as in *L. vacua*; they enter it on the underside. The cornu in this species, instead of being a separate sac attached by a long tube as in the last, is a mere short conical projection in the middle of the anterior surface of the sacculus, thus being almost at right angles to the ringed tubes; the apex

of the cornu is not closed, but a long fine tube runs from it to the camera spermatis; this tube also is shown extended in the figure, but *in situ* is wound round the sacculus and is difficult to see.

Holotaspis marginatus is a rather large and powerful species; in it the lyrate organ has a rather different appearance from what it has in the above-described species; the arms are comparatively very thin, but have sharply clavate ends (Pl. XXXIV. fig. 60, *ol*). The cellulation of these arms is apparently similar to that of other species, except that I am not able to detect the nuclei which are so very conspicuous in the others. On the other hand, the camera spermatis (*cms*) is extremely large, and its surface seems to be composed of cells similar to those of the arms, which is unusual; it appears as if it had taken upon itself part of the function of the arms in consequence of the small size of the latter. The sacculus is of the usual class of tissue, but is a long roughly-elliptical sac, with its long diameter transverse to the body. The cornu is not at all of the usual type; it is short, wide, and rather fleshy, and passes straight to the underside of the camera spermatis. In this species there appears a new complication of the organs which is far from easy to understand; the ringed tubes (*ta*), instead of terminating at or shortly after their entrance into the sacculus, continue in an almost straight line nearly to the middle of that organ, and there the two are joined, within the sacculus, by a small oval sac which communicates with each tubulus. There is, springing from the middle of this inner sac, a short, straight, and extremely fine tube running to the cornu.

Fig. 61 represents the sacculus of a small species found in Tyrol, and bearing considerable external resemblance to *Holotaspis montivagus*, but which I am unable to identify, because I dissected the one or two specimens which I found. I mention it because the structure of the organs I am describing is in some respects intermediate between the species last described and the next. Here the sacculus is rather small and bag-shaped; it stands free without cornu or rami or any apparent communication with the camera spermatis or the ovary; its only attachment seeming to be the ringed tubes which enter it at the corners, and terminate by bulbous endings within the sacculus.

Laelaps cuneifer (fig. 62) is constructed much on the same model; the sacculus is free, without cornu or rami; the ringed tubes, however, enter close together on the underside of the sacculus, and terminate where they enter it without continuing inside.

Holotaspis montivagus is the species where neither Prof. Berlese, who originally described it, nor I have ever found the male. I have considerable doubt whether the mode of classification, which includes this and *H. marginatus* in the same genus, be not an artificial one; but this is not the place to discuss that question. In this species there is a new and puzzling complication; it seems to partake of some of the characters of *Holotaspis marginatus* and some of those of *Laelaps cuneifer*. The lyrate organ (fig. 63, *ol*) is large and well-developed; it shows the ordinary cellulation with strongly-marked nuclei; these extend on to a portion of the exterior of the camera spermatis, but not all over it. The sacculus (*sa*) is rather small and globular; it is entirely free, not attached except by the ringed tubes; these enter the sacculus as in *H. marginatus*, and continue until near the centre; but there, instead of both terminating in a single oval sac as in *H. marginatus*, each ringed tube ends in a little hollow ball with rather

chitinized walls, the two not quite touching. The strangest part of the construction, however, is that between these two little globes, and attached to both, and I think (but of this I am not certain) communicating with them, is a third little hollow, slightly chitinized, vesicle, which is not quite a sphere, and from the outer end of which proceeds a fine serpentine tube, with a bulbous end, which tube is wholly within the sacculus and does not emerge at all. The sacculus, with the spheres and spheroid looking together like a trefoil, and the third tube are seen *in situ* in fig. 63, and the trefoil, &c., dissected out at fig. 64; both are carefully drawn from actual dissections; a sagittal section through the sacculus, showing one globe and the central spheroid and tube, will be found at fig. 71, *sa* (Pl. XXXV.).

The last species in which I shall describe these organs is a new one found in Tyrol. I propose calling it *Laelaps ligoniformis* (fig. 65). In this curious creature the lyrate organ is fully developed, but there is not any sacculus, cornu, or rami, only the ringed tubes; these run direct to the camera spermatis, and the two tubes join just at the point where they enter the camera. In this species there exists something not found in any of the others, viz., at or about the point where each ringed tube has its mouth in the acetabulum of the leg there is a small elliptical receptacle (fig. 66), about 40 μ in long diameter, communicating with the exterior by a short stalk about 17 μ long. I had not enough specimens to investigate this species properly.

All the sacculus organs in all species are apt, when empty, to shrivel up and be almost impossible to find; but they may then often be found by placing the creature, after the dorsal plate has been dissected off, in water, when they fill by endosmosis and expand.

It is to be remembered that the sacculus organs and ringed tubes are apparently correlated with the lyrate organ. I have not found the one without the other in any species; *L. ligoniformis* is the nearest approach, but even there the ringed tubes are present, while none of the species of the *G. crassipes* type, which I am acquainted with, show any trace of either organ.

We now come to the question of the contents and function of the sacculus fœmineus and its associated parts or diverticula. Of course the first obvious suggestion of function, from the ending of the ringed tubes in the acetabula of the third legs, would be that the whole structures were coxal glands; but, apart from the contents of the sacculus, the equally obvious reply is that these organs exist only in the female, and that if they were coxal glands they would exist in both sexes, as is the case with the supercoxal glands of the Oribatidæ, and the coxal glands of such other Acarina as are known to possess those organs, including, according to Winkler, the Gamasinæ, where they are very small and totally unlike the organs I have been describing. Winkler figures them in the male.

The presence of the organs in the female only seems to confine our suggestions as to their function to the generative system; and all the other evidence which I have obtained confirms this view. Omitting for the moment the very specialized species *Uotaspis marginatus* and *H. montivagus* and their one or two allies, all the other species investigated which possess the sacculus and its appendages usually have contents therein. Taking again *Hæmogamasus horridus* and *H. hirsutus* as convenient

types, it will be seen by Pl. XXXIV. figs. 48, 49, 53, &c. that the sacculus and cornu usually contain a number of loose bodies, varying in quantity and mixed up with a thickish fluid and granular or minutely filamentous matter. If the testes of the males of these respective species be examined, and the most mature spermatocysts be compared with the bodies contained in the sacculus of the female and its cornu and diverticula, it is impossible to avoid noticing the extreme similarity of the two: those in the sacculus &c. are usually a little smaller than those in the testis, and show signs of having been subjected to rather more pressure; often they have burst and are mere empty shells, but the perfect ones retain the shape and appearance of the male spermatocysts. If a male *H. hirsutus* be killed at that stage of the coitus when the spermatocapsule has been filled but not discharged, and if the capsule be allowed to discharge upon a glass slip and its contents be compared with those of the sacculus of a female freshly killed, immediately after the completion of the coitus, the identity of the two cannot be doubted (Pl. XXXII. figs. 24, 25).

In *Laelaps vacua* the commonest form of the ripe spermatocysts in the testis of the male is rather special; it is what I believe is known as the "curved dumb-bell" shape in sponge-spicules (Pl. XXXIII. figs. 27, 28). This is also the commonest form amongst the bodies found in the sacculus fœmineus, and the two agree in other respects besides form. It is indeed a fact that in all species examined the spermatocyst in the male testis has an appearance more or less peculiar to the species; and that, in spite of slight variation, the bodies in the sacculus usually fairly correspond to the ripe spermatocysts; also that if the latter can be obtained from the spermatocapsule of the male and from the sacculus of the female immediately after coition that resemblance amounts to identity as far as I have been able to investigate the matter. There are, however, some species, such as *Laelaps cuneifer*, *Holotaspis montivagus*, *Holotaspis marginatus*, &c., where the sacculus never contains spermatocysts, but only fine granular matter if anything, which may or may not be spermatozoa; but in these species there generally is not any connection between the cornu and the camera spermatis.

It may, I think, therefore safely be said that the contents of the sacculus are the products of the male genital organs.

In cases where the sacculus has contents but they are small in amount, which most usually occurs where the creature is oldish and the ova very fully developed, or the ovary nearly or partly exhausted, the whole contents will generally be found collected in the cornu. If the organs be fuller, then the sacculus itself and, in *H. hirsutus*, the great globular diverticula are amply supplied, occasionally almost full. If the last-named species be killed immediately after coitus, then the rami will be found to be full also; but the contents, in the cases I have seen, very soon slip out of the rami into the sacculus. I have not been able to obtain *H. horridus* in coitû.

The question arises, how do the spermatocysts get into the sacculus? There appear to be only two possible methods—the one that they are ejected from the spermatocapsule into the vagina, pass thence all along the oviduct, through the camera spermatis and perhaps part of the ovary, thence into the cornu, and then into the sacculus and its diverticula, and even into the rami, whence they finally return, as required, by the same road to the camera spermatis. This seems an extremely elaborate and not very probable

mode, but it has in its favour the fact that the vagina is a large opening, and that such species as *Gamasus crassus* and *G. crassipes*, which have not any sacculus or correlated organs, but which have a spermatheca in the vagina itself, are, as shown at page 296, fecundated by that opening.

The other possible method is that the products of the male organs are ejected from the narrow mouth of the capsule into the more or less trumpet-shaped mouth of the ringed tubes in the cuticle joining the coxa of the third leg to the more chitinized part of its acetabulum, and pass directly up the ringed tubes into the sacculus; this is a sufficiently simple method, and is rendered more probable by the similarity of the situation of the opening to that by which *Astacus* and many other Crustacea and Myriapoda are fecundated, although in these cases it is also the aperture for the deposition of ova. The only objection which I see to this is the fact that the spermatocysts found in the sacculus undoubtedly appear considerably too large to have passed through the ringed tubes; this is a serious difficulty, but it is possible, and even probable, that the ringed tubes may be capable of distension, and that the spermatocysts, which are soft bodies, are capable of compression, and can be forced through very small apertures, just as the eggs are forced through even hard, not distensible, openings which seem quite incapable of allowing them to pass. The weight of the objection is also considerably modified by the circumstance that it practically has to be faced in either explanation, for the large spermatocysts are found both in the cornu and the sacculus; and yet in *Laelaps acuta* the tube leading from the camera spermatis to the cornu and in *L. vacua* the tube leading from the cornu to the sacculus are scarcely larger in diameter than the ringed tubes, and yet if the spermatocysts be introduced into the vagina they must pass through both these tubes before they can get into the sacculus, but in the sacculus they are found abundantly.

I had hoped to settle this question by actual observation, but I have found that it is not possible to see sufficiently clearly to be quite certain whether the small end of the spermatheca is applied to the opening of the vagina or to the mouth of the ringed tube; the whole capsule is certainly not introduced into the vagina as it is in *G. crassus*, but the epigynum, which covers or borders the opening of the vagina, lies just between the third legs, and therefore just between the opening of the ringed tubes, and, being large, almost reaches them. In the position which the male and female occupy during the coitus, it is almost impossible to be absolutely certain whether the narrow end of the capsule, which lies between the mandibles of the male, is applied a little more or a little less to the side; but in the instances of *H. hirsutus* which I have watched it has appeared to me, as far as I could judge, that the mandibles were too far at the side for the vagina, and seemed much more like being at the mouth of the ringed tubes.

I have not even found spermatocysts or spermatozoa in the ringed tubes, but neither have I ever found them in the vagina or oviduct of any species possessing the sacculus or ringed tubes, although I find them abundantly in those situations in species which have not the ringed tubes.

It must be clearly borne in mind that in such species as *G. crassus*, *G. terribilis*, &c.,

where I know that the capsule is discharged into the vagina, there is a distinct spermatheca, usually filled with sperm, leading out of or forming part of the vagina, which does not exist in any of the species that possess the sacculus or ringed tubes. It is also worthy of remark that in the one species investigated which has not possessed either the sacculus or this vaginal spermatheca or recess (*Laelaps ligoniformis*) there is what I should judge to be a spermatheca (Pl. XXXIV. fig. 66) just at the opening of each ringed tube, and this does not exist in any of the other species.

It is also to be remembered that in most spiders, e. g. *Tetragnatha*, *Theridium 4-punctatum*, &c., the spermathecæ do not open into the vulva, but have separate external apertures*. In the spiders, or most of them, the spermatozoa are contained in spermatophores or spermatocysts.

Taking all these matters into consideration, I incline to think, although I should not like to state it as a fixed opinion, that in what I may call the sacculus species the sperm-elements enter the body of the female by the ringed tubes, and not by the vagina.

From the above-described facts, and particularly having regard to the great change which takes place in the sperm-elements of *Hæmogamasus hirsutus* between the time when they are found in the capsula spermatis and that when they enter the camera spermatis, I think that the following is probably the function and action of the female organs above described, viz. :—That the germ-elements are produced in the arms of the lyrate organ, and gradually work their way to the base of that organ, where they break off and pass into or by the base of the camera spermatis; that the sperm or spermatocysts enter the sacculus probably, but not certainly, by way of the ringed tubes and rami; that the spermatozoa are at that time not fully developed, and that they only attain their final development within the body of the female; that the sacculus with its cornu, and diverticula when present, is the organ in which this development takes place, and whose office it is to mature the spermatozoa; that when the spermatozoa are really ripe they pass out of the cornu into the camera spermatis; that there they meet the germs and fertilize them; that the ova then lodge in what is called the ovary and form oöcysts by the outpushing of its walls; that in these oöcysts they attain a considerable amount of development, although probably not as high a state of maturity as they will attain in the oviduct, whence they pass through the vagina to the exterior.

It may be suggested with considerable force that this explanation fails in the case of such species as *Laelaps vacua* and *L. cuneifer*, where there is not any apparent connection between the cornu and the camera spermatis or ovary; I cannot help thinking, however, that, although I have not found it, this connection must exist at some time of year or under some circumstances; I cannot otherwise understand the presence of abundant spermatocysts in the sacculus.

Finally it remains to account for the function of the organs in such a species as *Holotaspis montivagus*, with its three little hollow balls forming a trefoil within the

* Bertkau, " Ueber den Generationsapparat der Araneiden," Archiv für Naturgesch. 1875, p. 285.

sacculus, and its small third tube entirely coiled away within the sacculus; but this, I confess, I am not able to do at present; it must remain for future investigation. I have little doubt that the third tube is the homologue of the tube which joins the cornu to the camera spermatis in *Laelaps acuta*, and it is possible that, under some circumstances, it may emerge from the sacculus and join the camera as in that species. The three chambers of the trefoil are not usually empty; they more generally contain very fine granular matter, which might possibly be spermatozoa, but certainly not spermatocysts. It may be possible that in this species, where the male has never yet been found, the sacculus has become modified into an organ with other functions not yet understood; or it may even be possible that an agamous generation exists, and that the sperm-elements may be preserved in this singular receptacle from the mother within whose body the embryo becomes so highly developed; but these are only mere suggestions of possibilities, and I do not in any way put them forward as facts or even probabilities.

Another totally different but very interesting variation of the female genital organs of the Gamasinæ from Winkler's type (*G. crassipes*) is found in *Sejus togatus*, Koch. These parts are delineated in Pl. XXXIII. fig. 43. It will be seen how unlike they are to anything yet spoken of; they are a thorough return to the ring form. There is a single azygous central ovary (*ov*), having the appearance of a bunch of grapes—all the numerous larger eggs being in oöcysts on the exterior, and the germinal part more central. From the ovary proceed two oviducts (*od*), not one, as in all the other species we have been considering; these two oviducts join in an azygous vagina (*va*), out of which leads an immensely large, flask-shaped spermatheca (*spt*), which lies between the oviducts, has a narrow mouth and stalk, and powerful muscles attached evidently for the purpose of ejecting the contents. It is generally full of spermatozoa of the curious form shown by fig. 45. These spermatozoa are also found in the oviducts and ovary of the female.

The principal interest of these organs lies in the fact that, although very unlike what are usually found in the Gamasinæ, they closely resemble those of the other subfamily, the Uropodinæ; although the general appearance of the creature and the anatomy of the alimentary canal, and most other parts, are of the type of the Gamasinæ, not of the Uropodinæ. This relationship of the female is rendered more interesting from the fact that the male also shows certain affinities to the Uropodinæ.

THE ALIMENTARY CANAL.

I have not many variations to record in this system; one or two, however, that I have observed may be of some interest.

The ventriculus in most species of Gamasidæ is a smallish viscus, having a moderately large and wide cæcal prolongation forward in the central line, and often a corresponding one backward. Two large and long paired sac-like cæca spring from its lateral anterior

part and run forward; and two pairs of similar but larger cæca spring from the lateral posterior part and run backward: these are arranged one above the other; the upper pair are the longer: they usually agree in both sexes.

In *Hæmogamasus horridus* the female agrees with the above description, but the male is entirely without the lower pair of hind cæca (Pl. XXXV. fig. 72).

In *Laelaps cuneifer* the anterior median cæcum is bifid at its anterior extremity, each half curling slightly outward in a rather singular hood-like manner; the anterior lateral cæca also are recurved at their distal ends in a form which I have not seen elsewhere; the hind central cæcum is very wide. I have omitted the drawing which I had made of this canal in order to relieve the already overcrowded Plates.

An interesting feature is found in the pharynx of *Hæmogamasus horridus*. It is well known that in predatory Acarina, and indeed in most that live by suction, the pharynx is the sucking-organ, and that it is provided with a special set of muscles beautifully adapted to dilate and contract it. This, which has been well described by Henking*, MacLeod†, and others, is not confined to the Acarina, but is found also in many other Arachnida. Winkler also, in the paper so often referred to above, describes it well in *Gamasus crassipes*, and shows there that the lumen of the pharynx is an upright oblong with two branches at its upper end; he also mentions that its walls are somewhat chitinized; he gives excellent delineations of the musculature.

In dissecting *Hæmogamasus horridus* I was surprised to find that the development in the same direction had gone much further, and that there was an actual chitinized skeleton to the pharynx, forming a closed chitinous organ of a very definite shape and highly elastic; this is delineated by fig. 23 (Pl. XXXII.), from which it will be seen that the whole structure is lanceolate and comes almost to a point posteriorly, where the œsophagus arises from it; and is truncated anteriorly, where it joins the mouth. The organ is tricarinate, *i. e.*, if three carinæ of birds' sterna were hollow and open at their proximal edges, and were placed with these edges together so as to form one common lumen, and to diverge from one another at equal angles, a very fair model of this pharyngeal skeleton would be obtained; but it is all one piece, without any break or space between the carinæ, which are firmly ankylosed to each other. Thus the lumen in a transverse section of the organ looks like a triradiate sponge-spicule (fig. 73, *ph*). It will be seen from the same figure that one carina goes perpendicularly downward, that the two others branch from it like the arms of the letter Y, and that the large distensor muscles are attached to the side-surfaces of all the carinæ; while the much smaller constrictors pass from one edge to the other, and are doubtless aided by the elasticity of the chitin. In spite of the small size it is quite possible to dissect this organ out, and fig. 23 was drawn from such a preparation, which is still in my possession.

It may be mentioned that in *Hæmogamasus horridus* and *H. hirsutus* the two great

* "Beiträge zur Anatomie, Entwicklungsgeschichte und Biologie von *Trombidium fuliginosum*, Herm.," Zeitschr. f. wiss. Zool. Bd. xxxvii. (1883) pp. 569, 570.

† "La structure de l'intestin antérieur des Arachnides," Bull. Acad. R. d. Belg. 1884.

Malpighian vessels which are usually free are attached to the hind cæca of the ventriculus almost the whole length of the cæca.

EXCRETORY ORGANS.

In addition to the great Malpighian vessels there exist in some Gamasids, although apparently not in all, a pair of organs which I believe have not yet been recorded in the family; they lie in the side of the hinder part of the abdomen close under the skin, and are stout sacs discharging to the exterior by a pore in the cuticle. In *Hæmogamasus horridus* there are two such sacs on each side; they are flattened and the two are close together (Pl. XXXII. fig. 22). They have been practically similar in the other cases where I have found them. These organs are well known in other Acarina, *e.g.* in the Tyroglyphidæ, where Claparède noticed them as long ago as 1868 *, in the Oribatidæ, where I described them and called them the expulsory vesicles †, &c. They contain a yellowish oily fluid, and have been generally admitted to be excretory organs; they appear to be sacs closed on the interior side, their only opening being external.

MUSCULATION.

I do not find that the extremely pretty levatores ani muscles, and the neighbouring muscles which serve to compress the hind part of the abdomen dorso-ventrally, have been figured or described. Fig. 74 (Pl. XXXV.) is from a transverse section of *H. hirsutus* passing through the anus, and the musculature will be so readily understood from that drawing that I do not think I need describe it. The drawing is made from a preparation of a specimen killed with boiling water, and the anus is consequently rather unnaturally protruded, but I selected it as showing the construction better than other specimens where this had not occurred.

SUMMARY.

For the convenience of those who wish to see at a glance whether this paper contains anything likely to interest them I beg to summarize it as follows.

I suggest that it contains :—

1. A series of observations on the coition of the Gamasinæ, showing a mode previously quite unknown except from a partial description of a single case formerly recorded by me; and only less curious than that of the spiders, but entirely different therefrom.
2. Some considerable differences in the male sexual organs from any previously

* "Studien an Acariden," Zeitschr. f. wiss. Zool. xviii. Bd. pp. 445, 546, Taf. xxxv.

† 'British Oribatidæ,' Ray Soc. p. 179.

described in the family, and some singular cases of the formation of spermatozoa and spermatocysts not previously recorded.

3. A description of a set of organs in the female genital system not previously known, and which apparently serve chiefly for the maturation of the sperm-elements within the body of the female, with the varieties of these organs in numerous species, also some other not unimportant variations in the female reproductive organs from any hitherto recorded in the family.

4. A description of the female reproductive system, not before known, of a member of the genus *Sejus* showing its similarity to those of the Uropodinæ.

5. A description of a pharyngeal skeleton and some other points of interest relative to the alimentary canal.

6. A few other novelties of less importance in the internal anatomy of the Gamasinæ.

DESCRIPTIONS OF NEW SPECIES.

HÆMOGAMASUS HORRIDUS, sp. n. (Pl. XXXII. figs. 1-5.)

	♀	♂
Length without rostrum, about.....	1·4 mm.	1·16 mm.
Breadth, about	·77 „	·65 „
Length of legs, 1st and 4th pairs, without caruncle or ungues, about	1·3 „	1·1 „
„ „ 2nd „ 3rd „ „ „ „ „	·90 „	·70 „

Colour darkish umber-brown.

Texture. Almost the whole dorsal surface is covered with a chitinous plate, which, however, is rather thin; it is smooth but not polished; there are not any markings on the chitin. The plate covers the fore part of the body but does not extend to the posterior end; a border of softer white cuticle is shown round the posterior and the greater part of the lateral margins, especially in the female.

Shape longish; nearly parallel-sided, but somewhat wider behind than in front; posterior end rounded, anterior slightly shouldered and narrowed in. Dorsal surface rather flat.

Mandibles. Those of the male are exceedingly curious; they are elaborate and difficult to describe; they will be best understood from fig. 3. The whole mandible is short and thick for the family; the fixed arm of the chela thick, not diminishing in thickness until close to the distal end, when it suddenly curves downwards like a parrot's beak; but the projection of the beak below is very slight. There are two minute teeth close to the beak; the rest of the arm is rather undulated than toothed. The movable arm of the chela is the more singular; it is a broad undulated piece without teeth until nearly opposite the end of the fixed arm, then there is a break or indentation on the under edge; from this point arises a great hook, which first curves forward, then turns strongly backward, almost parallel to its outward

course, and continues in that direction until near its origin, when it turns outward and upward and ends in a sharp point. At the place where the hook arises there are two discoidal or ovoid projections from the upper edge of the movable arm, the proximal smaller than, and somewhat overlying, the distal; these are less highly chitinized than the rest of the chela. The portion of the movable chela which works within the penultimate joint of the mandible forming the articulation is very large, and has a considerable-sized foramen in its upper part; from its lower part there projects a long narrow sac of cuticle, not much chitinized, and which stands free from the rest of the mandible and nearly reaches the indentation before the hook. The mandible of the female (fig. 4) has the distal end of the fixed arm of the chela much more gradually curved and thinner than that of the male; the two teeth behind it are much stronger, and there are two curious transparent blades which descend from its outer edge and appear like guides to the movable arm. Two very much smaller blades of similar material are found on the lower edge of the last-named arm, which otherwise has not anything remarkable about it; it is entirely without the great hook of the male.

Epistome (fig. 5) hyaline, campanulate (with the broad straight end attached to the body), lateral edges curved downward; the whole free edge is bordered by a dense fringe of long fine hairs directed forward.

Legs near together, long; without apophyses in either sex; all legs terminated by long caruncles and claws, and furnished with numerous strong spines and hairs.

Dorsal surface densely clothed with short spine-like hairs, set in transverse wavy rows; the hairs longer than the distance between the rows, and alternate, so that the end of each hair overlaps the row behind and lies between two hairs of that row. These hairs all point backward except those at the edge, which stand slightly outward and form a fringe round the body. The hairs increase a little in length from the anterior to the posterior part of the body; the longest are those on the soft cuticle. Length of the hairs from about .03 millim. at the anterior to about .07 millim. at the posterior end of the body; distance apart laterally about .02 millim. There are two long straight hairs projecting over the rostrum.

Ventral surface haired. Anal plate small, inversely pyriform; there are three long hairs behind it. Genital plate of female sac-shaped, widest posteriorly, attached by its whole surface, not hinged; its anterior edge bordered by a fine membrane. Genital aperture of female a curved transverse slit. There are two small curved metapodic plates, one on each side of, but some distance from, the genital plate. Stigmal plates narrow and curved. Sternal plate with distinct scale-like markings.

Habitat. Found commonly, but not abundantly, in the nests of the Common Mole (*Talpa europæa*) in England. The young may probably be parasitic on the Mole, as I do not often find them in the nest.

HÆMOGAMASUS NIDI, sp. n. (Pl. XXXII. figs. 6 and 7.)

	♀	♂
Length without rostrum, about	·94 mm.	·72 mm.
Breadth, about	·51 „	·40 „
Length of legs, 1st and 4th pairs, without caruncle or ungues, about	·78 „	·60 „
„ „ 2nd „ 3rd „ „ „ „ „ „	·53 „	·45 „

This species is so like *H. horridus* in general appearance that I have not thought it useful to draw a figure of it; there are, however, well-marked anatomical differences; it is also much smaller and lighter in colour. Its movement is different from that of *H. horridus*; *H. nidi* moves in little jerks or springs, *H. horridus* does not.

Colour yellow-brown, not very dark.

Texture. The whole, or almost the whole, dorsal surface is covered with a chitinous plate; the chitin, however, is very thin, and allows the Malpighian vessels and often the ventriculus to be plainly seen through the dorsum; it is smooth, but not polished; there are not any markings on the chitin. Sometimes a narrow border of soft cuticle is seen round the hinder part of the body, especially in the female, but not nearly so wide or conspicuous as in *H. horridus*.

Shape. Long, almost parallel-sided; posterior end rounded, anterior slightly shouldered and narrowed. Dorsal surface rather flat.

Mandibles of the male (fig. 6) with the fixed arm of the chela gradually diminishing in thickness and curving downward, forming a slightly recurved pointed claw-like ending; there is one strong tooth far back. The movable arm is a very powerful strongly curved piece which overlaps the fixed arm, when closed, some distance behind the point of the latter. From the outer side of the movable arm springs a large doubly-curved accessory piece, which extends a little beyond the end of the fixed arm, and curves upward; it is truncated at its distal end; but at the inner edge of the truncated surface arises a sharp spine, directed perpendicularly upward close to the end of the fixed arm; there is also a second accessory piece almost hidden by that last described.

Epistome (fig. 7) rather irregular, sub-triangular, with rounded apex and concave sides; the greater part of the lateral edges and the distal edge are fringed with processes which can hardly be called hairs, but are rather long ragged projecting pieces of the material of the epistome itself; they are not set close together nor bisymmetrically on the two lateral edges.

Legs rather near together, long and thin, without apophyses in either sex. All legs terminated by long-shaped caruncles and claws. Legs furnished with numerous curved hairs and spines.

Dorsal surface densely clothed with short spine-like hairs set in transverse wavy rows, the hairs slightly longer than the distance between the rows, and alternate; they all point backward except those at the lateral edge, which stand slightly outward and form a fringe. The hairs increase a little in length from the anterior to the posterior end of the body; the longest are those on the soft cuticle. Length of

hairs from about .02 millim. at the anterior to about .04 millim. at the posterior end of the body; distance apart laterally about .01 millim.

Ventral surface haired, all the plates of very thin chitin; anal plate small, inversely pyriform. Genital plate of female larger than in *H. horridus*, widened and rounded posteriorly, attached by its whole surface. Genital aperture of female a transverse slit. There are two little lateral metapodic plates as in *H. horridus*; stigmal plates curved and very thin.

Habitat. Found abundantly in the nests of the Common Mole (*Talpa europæa*) in England.

LÆLAPS ORIBATOIDES, sp. n. (Pl. XXXII. figs. 12-15.)

	♀	♂
Length without rostrum, about	.77 mm.	.47 mm.
Breadth, about	.47 "	.38 "
Length of legs, 1st pair, without caruncle or ungues, about	.60 "	.48 "
" " 2nd " " " " " "	.46 "	.36 "
" " 3rd " " " " " "	.42 "	.33 "
" " 4th " " " " " "	.57 "	.50 "

The female of this species may generally be recognized among its neighbours of the same locality by two large black spots on the dorsum, near the posterior end, which are really produced by the excretory organs showing through the cuticle.

Colour yellow-brown, not very dark.

Texture. The whole dorsal surface is covered by a single unbroken chitinous plate; the chitin is rather thin, and is entirely divided up, by very fine lines, into little scale-like or angular spaces, measuring rather more in a direction across the body than in the direction from anterior to posterior end; in the former direction averaging about .02 millim. No other markings; the chitin otherwise smooth, but not polished.

Shape. Almost elliptical, very slightly shouldered in front, rather rounder posteriorly.

Mandibles of male (fig. 13). Fixed chela with a single sharp spine (the poison spine?); no other teeth. Movable chela not much longer than the fixed, broad and divided into three parts, viz.: (1) the principal piece of the arm, which is truncated almost at a right angle to its lower edge, and has its upper distal end a long curved point; there is one strong median tooth; (2) a long narrow slightly undulated accessory piece projecting beyond No. 1; (3) a broad outer lamina about as long as the accessory piece, and having a rounded end with a notch in it.

Epistome (fig. 15) rounded, with a few small triangular or tooth-like projections on its distal edge, and a few fine radiating markings on its proximal dorsal surface.

Legs not very long, second pair the thickest; without apophyses in either sex. All legs furnished with not very numerous hairs, and with roundish caruncles and very fine claws.

Dorsal surface provided with fine hairs, set not very near together in irregular transverse rows, the hairs slightly longer than the space between the rows; they all point

backward, except two anterior, which point forward over the rostrum; length of hairs increases a little from the anterior to the posterior part of the body, the former about .04 millim., the latter about .06 millim.

Ventral surface. All the chitinized plates divided up into scale-like spaces similar to those of the dorsal surface, but even more distinct. In female anal plate triangular, much broader than long, broad end forward, the two anterior angles very sharp, the posterior truncated; anus in the middle of the plate. Ventral plate large and broad, only divided from anal by a narrow line. Ventral and genital plates anchylosed. Two large triangular metapodic plates behind the fourth leg. Stigmal plates much expanded posteriorly. Sternal plate normal. The whole ventral surface is haired like the dorsal, but the hairs are not so large.

Habitat. Frequent in the nests of the Common Mole (*Talpa europæa*) in England. The male rarer than the female.

LÆLAPS (?) LIGONIFORMIS, sp. n. * (Pl. XXXII. figs. 16 and 17.)

	♀	♂
Length without rostrum, about	.50 mm.	.37 mm.
Breadth, about	.37 „	.25 „
Length of legs, 1st pair, without unguis, about	.34 „	.28 „
„ „ 2nd „ „ „	.25 „	.25 „
„ „ 3rd „ „ „	.30 „	.27 „
„ „ 4th „ „ „	.44 „	.38 „

Colour dull, light drab-brown.

Texture slightly rough, but without distinct markings, the whole dorsal surface covered by an unbroken chitinous plate.

Shape. Somewhat spade-shaped, whence the name. The female obtusely pointed anteriorly; broadly truncated posteriorly. The male more rounded at both ends, but narrower. Dorsum rather flat, particularly in the female.

Mandibles of male (fig. 17) with the fixed arm of the chela regularly serrated with a number of closely-set sharp teeth, and the end turned suddenly down, forming a large terminal tooth. Movable arm curved, swollen at the base, thin at the distal end, with two large teeth there. From the side of the movable chela springs a large accessory piece which projects forward and downward, is undulated, of varying thickness, and has an almost globular end; it bears a spine about the middle.

Legs without special features; the fourth pair considerably the longest, and stretching much behind the body. No apophyses in either sex; claws of all legs large and distinct; no caruncles.

* I scarcely think that this species can ultimately remain in the genus *Laelaps*; on account of the exceptional conformation of the ventral surface of the male, it may require an entirely new genus; but I think there are other unrecorded species allied to it which it would be well to investigate before defining such a genus; the species may, therefore, be provisionally placed in *Laelaps*.

Dorsal surface with four longitudinal rows of about ten spines on the dorsum, and two rows, or a double row, all round the periphery except just the anterior edge, on which there are two fine spines, near together, projecting forward. The dorsal spines are sparsely set, increase greatly in size from the anterior to the posterior end of the body, and do not overlap, except at the rear. The anterior spines are sharply pointed; those of the female about .03 millim. long: the hind spines are very thick, blunt, and spatulate; those of the female about .12 millim. long.

Ventral surface. Female with anal plate fused with ventral, very large, covering half the abdomen, and divided by fine lines into scale-like spaces. Genital plate almost touching the fused ventral and anal. It is bell-shaped, *i. e.* rounded anteriorly, straight posteriorly; a central triangle with the point forward is the most chitinized part in this plate; the peripheral band is also chitinized, but two lenticular spaces between are slightly chitinized and look almost like foramina. Sternal plate normal; there are two small plates near its anterior edge and two near its posterior corners. Stigmal plate almost straight, but clavate posteriorly. The male, in addition to the usual sexual differences of the plates, has in the middle of the ventro-anal plate, and about midway between the anus and the coxæ of the fourth leg, a large and singular projection standing straight downward and formed like a short length of concave gutter (half tube), from the lower edge of which project two large chitinous apophyses, much more strongly chitinized than any other part of the body. There are three or four spines round it. I do not know the function of this organ. It has the appearance of a male genital armature, but the ordinary male genital aperture is quite plain and unmistakable in its usual place at the other end of the body, *viz.* just in front of the anterior edge of the sternal plate.

Habitat. I found a very small number of specimens near Innsbruck, in Tyrol, on the stumps of dead pine-trees, and while this paper has been passing through the press I have found the species in large numbers in England on the underside of fungi growing on old tree-stumps at Goring in Oxfordshire.

I regret that want of space on the Plates has prevented my figuring the ventral surface of this creature.

EXPLANATION OF THE PLATES.

A. Anus. ac. Acetabula for insertion of the coxæ of the legs. The figures 1, 2, 3, 4 refer to the 1st, 2nd, 3rd, or 4th pair of legs. ap. Anal plate. bp. Bipilate appendage (Winkler's labium, Kramer's "Bauchtäster").	br. Brain (so-called). The small dark cells are the cortical or corpuscular layer, the rest the inner or fibrous mass. cæ. Front cæcum of the ventriculus. cæ¹. Upper hind cæcum of the ventriculus. cæ². Lower hind cæcum of the ventriculus.

- cas.* Capsula spermatis (Spermatic capsule). See page 294.
- cc.* Chitinous cuticle.
- cl.* Cloaca.
- cm.* Soft (flexible) cuticle.
- cms.* Camera spermatis (Sperm-chamber). See pages 299, 300.
- cs.* Cornu sacculi (the horn of the sacculus). See page 301.
- ct.* Connective tissue.
- de.* Ductus ejaculatorius.
- dg.* Cells considered by Winkler to be dermal glands.
- dp.* Dorsal plate.
- ep.* Epistome (dorsal anterior edge of the oral tube, or rostrum).
- epg.* Epigynum (external genital plate of the female).
- eph.* Epipharynx.
- fc.* Fat-cells.
- ga.* Genital aperture of the male.
- gla.* Accessory glands of the male genital system.
- hd.* Hypodermis.
- hg.* Hind gut.
- hs.* Hypostome (ventral anterior portion of the oral tube).
- l.* Leg, the figures 1, 2, 3, 4 indicate the 1st, 2nd, 3rd, or 4th pair.
- la.* Levatores ani muscles.
- m.* Muscles of the legs; the figures show the pair of legs.
- md.* Mandibles.
- mdr.* Retractor muscles of the mandibles.
- mph.* Pharyngeal muscles (page 310).
- mr.* Retractor muscles of the rostrum.
- msp.* Suspensory muscles of the tendinous platform (episternite).
- mv.* Malpighian vessels.
- nm.* Mandibular nerve.
- O.* Ovary (see note on page 297).
- od.* Oviduct (see page 298).
- œ.* Œsophagus.
- ol.* Lyrate organ (see page 298).
- ov.* Ova.
- ove.* Ovum with nearly matured embryo.
- ovs.* Ovum with yolk-segmentation.
- P.* Palpus.
- ph.* Pharynx.
- pt.* Peritreme (stigmatic canal).
- R.* Rostrum.
- rs.* Rami sacculi (see page 301).
- S.* Stigma.
- sa.* Sacculus fœmineus (see page 300).
- sd.* Diverticula of the sacculus fœmineus (see page 302).
- sg.* Salivary gland.
- sm.* Sperm-mother-cells.
- sp.* Stigmal plate.
- spt.* Spermatheca.
- st.* Sternal plate.
- sz.* Spermatozoa.
- T.* Testis.
- ta.* Tubuli annulati (ringed tubes). See page 301.
- tp.* Tendinous plate for attachment of muscles (episternite).
- tr.* Tracheæ.
- V.* Ventriculus.
- va.* Vagina.
- vd.* Vasa deferentia.
- vp.* Ventral plate.

In all drawings of whole organs, or of groups of organs, except figures 40, 49, 66, the top of the Plate is the anterior end. This does not apply to sections.

PLATE XXXII.

- Fig. 1. *Hæmogamasus horridus*, ♀, sp. n. (page 312), dorsal aspect; × 35. Both mandibles are extruded.
- Fig. 2. The same species, ♀, ventral aspect; × 35. The legs are cut short to save space. The ventral surface of the oral tube is shown anteriorly; its front part is the hypostome (maxillary lip), the central fringed triangle being the lacinia of Mégnin, and the two lateral pieces the

galeæ of the same writer. Behind the oral tube is seen the bipilate organ ("Bauchtäster" of Kramer, "labium" of Winkler, &c.); behind this, in the median line, the almost hexagonal sternal plate; then the genital plate with a membranous anterior edge, and a small crescentic metapodic plate on each side of it posteriorly; finally, the anal plate (inversely pyriform). The stigmal plates are along the edges of the anterior part of the body.

- Fig. 3. The same species: last two joints of one of the mandibles of the male seen from the outer side; $\times 350$. The lowest lobe and the two smaller globular lobes of the movable arm of the chela are less chitinized than the other portions. The working or cardinal portion of the movable arm is sunk in the fixed arm, and is only shown, as though the latter were transparent, in order to exhibit its unusual form and elliptical foramen.
- Fig. 4. The same species: chela of one of the mandibles of the female seen from the outer side; $\times 350$. The two laminæ on the under-edge of each arm of the chela are less chitinized than the rest of the organ. The fixed arm of the chela in all Gamasid-mandibles is the upper arm.
- Fig. 5. (Near right lower corner.) The same species: epistome of the male seen from above; $\times 350$. It is very hyaline, but the basal triangle is rather more chitinized than the remainder. The radiating lines springing from this are markings, not hairs.
- Fig. 6. *Hæmogamasus nidi*, sp. n. (page 314): chela of one of the mandibles of the male from the outer side; $\times 380$.
- Fig. 7. The same species: epistome seen from above; $\times 350$. It is very hyaline; the radiating lines near the centre are markings, not hairs.
- Fig. 8. *Hæmogamasus hirsutus*, ♀, dorsal aspect; $\times 45$. This drawing is to show the course of the alimentary canal and Malpighian vessels; it is taken from a young, but mature, living specimen, the internal organs showing through the dorsal surface in consequence of the transparency of the cuticle. The spines and hairs are omitted for the sake of clearness; the legs also are omitted. The oral tube (rostrum) is in outline; the epistome is seen between and above the extruded mandibles; the irregular-shaped plate of thin chitin on the dorsal surface is semi-transparent, and varies in form in every specimen. The small ventriculus, with its great anterior and upper posterior cæca, is dark (as in nature when the creature is recently fed); the Malpighian vessels are almost white (as in nature when seen by reflected light). The cloaca is shown, looking globular near the centre of the posterior end, between the distal terminations of the cæca. The striated part is the soft cuticle.
- Fig. 9. (Near centre of lower part of Plate.) The same species; chela of one of the mandibles of the male from the outer side; $\times 650$. The lobate proximal portion of the fixed arm is flat (a blade) and is very hyaline.
- Fig. 10. The same species: epistome of female seen from above; $\times 350$.
- Fig. 11. The same species: epistome of another female seen from the same direction and magnified to the same extent, in order to show the variation in different specimens. Both this and fig. 10 are carefully made from actual examples.
- Fig. 12. *Lalaps oribatoides*, sp. n. (page 315), ♀, dorsal aspect; $\times 65$. The two conspicuous black spots arise from the excretory organs showing through the cuticle. The mandibles are retracted.
- Fig. 13. The same species: chela of one of the mandibles of the male seen from the outer side; $\times 355$. The sharp spine on the fixed arm of the chela is probably a poison-fang.
- Fig. 14. The same species: chela of one of the mandibles of the female seen from the outer side; $\times 300$.
- Fig. 15. The same species: epistome of female seen from above; $\times 300$. The organ is very hyaline, but the sub-triangular space at the base is rather more chitinized than the rest. The lines running forward from it are markings, not hairs.

- Fig. 16. (Left lower corner.) *Laelaps* (?) *ligoniformis*, sp. n. (page 316), ♀, dorsal aspect; × 75. Mandibles almost entirely retracted.
- Fig. 17. (Near 16.) The same species; chela of one of the mandibles of the male from the outer side; × 600.
- Fig. 18. *Holotaspis montivagus* (page 287): distal portion of the epistome of the female; × 345. This organ is hyaline; the smaller, central, Y-shaped projection not so much as the rest. The basal part of the organ, which is not drawn, is simply a continuation of what is drawn, forming a sub-triangular lamina with slightly rough edges. The wavy lines shown on the figure are markings, not hairs.
- Fig. 19. (Middle of right edge.) *Gamasus crassus*, Kramer (page 296): chela of one of the mandibles of the male seen from the outer side; × 190. The obovate foramen is seen in the movable arm, and the extremely minute poison-fang and duct (?) in the fixed arm.
- Fig. 20. The same species: epistome of male, three-quarter view from above; × 190. It is placed in this position in order to show how the distal end curves downward. The large central triangle is hollow and contains fine cellular matter; the peripheral part is dense chitin.
- Fig. 21. The same species: epistome of female seen from above; × 190. Wholly chitinized.
- Fig. 22. Expulsory vesicle from one side of *Hemogamasus horridus* (page 311); × 65.
- Fig. 23. Pharyngeal chitinous skeleton of *Hemogamasus horridus* (page 310), three-quarter view from below; × 170. The truncated (upper) end is the oral end; the small piece of tube springing from the lower (pointed) end is the commencement of the œsophagus.
- Fig. 24. Four ripe spermatocysts taken from the proximal end of the testis of a male *Hemogamasus hirsutus* immediately after dissection in water containing a small quantity of potassium acetate; × 400.
- Fig. 25. Five spermatocysts taken from the sacculus fœmineus of the female of the same species immediately after dissection in similar liquid; same amplification.

PLATE XXXIII.

- Fig. 26. Four spermatocysts taken from the ripest found in the testis of an adult male *Laelaps acuta* immediately after dissection; × 150.
- Fig. 27. Group of spermatocysts taken from the testis of an adult male *Laelaps vacua* immediately after dissection; × 300. The lowest are the ripest.
- Fig. 28. Three spermatocysts taken from the cornu sacculi of a female of the same species immediately after dissection; × 300.
- Fig. 29. The whole genital system of an adult male *Gamasus terribilis* (page 290), × 55, showing the two testes joined by a thin central band, the azygous ductus ejaculatorius, and the great accessory gland.
- Fig. 30. (Right middle.) The whole genital system of an adult male *Laelaps cuneifer* (page 292); × 70. The organs are in the natural position, as seen from above; the testis transversely across the body. The testis and vasa deferentia nearly hide the great accessory gland.
- Fig. 31. (Left top corner.) The same system from the same species; × 150. The testis is turned over backward, and the vasa deferentia extended, so as to show the bifid distal end of the accessory gland.
- Fig. 32. (Below fig. 30.) Terminal portion of the testis of the same species, × 300, showing the early condition of the sperm-mother-cells and their development into riper spermatocysts. The mature spermatocyst is very large in this species.

- Fig. 33. The anterior portion of the genital system, but without the testis, of an adult male *Laelaps laevis*; to show the two elliptical accessory glands (page 292) with the constriction at their points of discharge. A single spermatocyst is travelling down one vas deferens. This was so in the actual specimen.
- Fig. 34. (Left middle.) Sagittal median section through the testis of *Laelaps laevis*, $\times 150$ (page 292), showing the developmental history of the sperm-mother-cell and spermatozoa, so far as it takes place in the testis.
- Fig. 35. (The lowest line.) This is a series of drawings, numbered consecutively in small figures 1 to 11. It illustrates the history of the development of the sperm-mother-cells and spermatozoa in *Gamasus crassus* (page 293); all drawn, $\times 170$. These are all taken from the testis of the male; the latest stage is to the right, but those found in the female are usually even more filamentous.
- Fig. 36. (Right lower corner.) Outline of the rostrum (oral tube) and sternum, &c., seen from below, of a male *Hæmogamasus hirsutus* killed *in coitu* in an early stage of the coitus; showing the spermatic capsule still in the genital aperture of the male, and not yet full-sized nor containing spermatocysts; $\times 70$ (page 295).
- Fig. 37. Spermatic capsule of the same species at its largest size, $\times 70$; taken from a male killed *in coitu* at the moment when the spermatic capsule is about to be removed from the genital opening of the male by the mandibles of the male; it is now full of spermatocysts, &c.
- Fig. 38. Rostrum, mandibles, &c., seen from the side, of a male of the same species killed *in coitu*, at the end of the coitus, after the contents of the spermatic capsule had been discharged and after it had contracted by its elasticity; $\times 145$. The rostrum has been dissected out; the empty capsule is seen with its small end held between, and attached to, the two mandibles. The lower part of the rostrum is shown in optical section, in order to exhibit the pharynx and its musculature.
- Fig. 39. Rostrum, mandibles, &c., seen from the side, of a male *Laelaps cuneifer*, killed *in coitu*, at the same stage of the coitus as the last figure; $\times 150$ (page 295). The spermatic capsule is in a similar condition.
- Fig. 40. One mandible, seen from the side, of a male *Gamasus crassus* killed *in coitu*; $\times 110$ (page 296). The greater part of the mandible and the whole spermatic capsule were inside the vagina of female, and were dissected out after death. The spermatic capsule is seen passing through the foramen in the movable arm of the chela of the male mandible. The anterior end of the capsule has been fractured and destroyed; its contents have been discharged.
- Fig. 41. Genital aperture of a male of the same species killed *in coitu* at the moment when the spermatic capsule is about to leave the aperture, $\times 110$, showing the mode in which the end of the spermatic capsule, which emerges last, is folded over.
- Fig. 42. Spermatheca of the female of the same species; $\times 100$. It arises as a stalked diverticulum from the anterior end of the vagina, above and between the inner chitinized labia.
- Fig. 43. (Left lower corner.) The whole genital system of an adult female *Sejus togatus* seen from above; $\times 300$. It shows the racemose form of the ovary, the paired oviducts, and azygous vagina, all resembling *Uropoda* (page 309). The large central sac is the spermatheca, from which proceed muscles which are omitted in this figure.
- Fig. 44. (A little to the right.) The spermatheca of the same creature seen from the side; $\times 100$. This is to show the muscles attached to its hind and ventral surfaces.
- Fig. 45. (Left top corner.) A group of ripe spermatozoa taken from the testis of an adult male *Sejus togatus*; $\times 300$. Those taken from spermatheca of the female are precisely similar. Compare these with the penultimate form in *Gamasus crassus*, fig. 35, 10.
- Fig. 46. (Right lower corner.) General view of the alimentary and genital systems of an adult female

Laelaps laevis; $\times 75$. The figure is to give an idea of the arrangement of the parts relatively to one another; it is taken looking down from above, after the dorsal plate, the fatty matter underlying it, and such muscles as interfered with the view had been removed. The whole of the ventriculus and its anterior and its upper posterior cæca are seen (light); the lower posterior cæca just show (dark). The hind gut is hidden by the sacculus fœmineus and its cornu, which are in the centre, and by the ova. The most mature ovum, *ovs*², must in one respect be considered diagrammatic, because it should have been drawn very much larger; but to have made it the full size would have greatly increased the size of the figure without corresponding advantage, and would have obscured parts of the cæca which I wished to show. The distal ends of the lyrate organ are seen (dark) projecting from below the anterior lateral cæca of the ventriculus. The œsophagus and pharynx are not drawn, as they are not in the part of the body where the genital and alimentary systems interlock.

Fig. 47. Terminal portion of one of the arms of the lyrate organ of an adult female *Hæmogamasus hirsutus*; $\times 275$. To show the cellulation, &c.

PLATE XXXIV.

Fig. 48. The sacculus fœmineus, with its upper appendages, &c., and the lyrate organ and ovary of an adult female *Hæmogamasus horridus* (page 297), *in situ*, seen from above; $\times 60$. The central part of the lyrate organ and of the ovary, and the camera spermatis and oviduct, are hidden by the sacculus, which contains numerous spermatocysts, &c. The cornu sacculi contains similar bodies, and passes underneath to the camera spermatis. The rami and ringed tubes are extended.

Fig. 49. The same organs and the oviduct of an adult female of the same species seen from the side; $\times 85$. This is taken from a specimen which, although adult, was young, *i. e.* comparatively recently mature. The sacculus fœmineus is fuller of spermatocysts than in the last figure, but the ova are very little developed. The arms of the lyrate organ would not descend so much *in situ*—they would be almost horizontal; they have been turned slightly downward to show the whole of the sacculus.

Fig. 50. (Right lower corner.) The same species. One of the tubuli annulati (ringed tubes) seen from the side, $\times 175$, to show the form of the mouth of the tube in the acetabulum of the third leg. The acetabulum is not shown, only the tube.

Fig. 51. The sacculus fœmineus, with its appendages and the lyrate organ, of an adult female *Laelaps stabularis*, seen from above; $\times 100$. These organs are not quite *in situ*: the cornu is straightened out to show its whole form; it would really curl more round the base of the lyrate organ. A few spermatocysts are seen in the rami (as after recent coition).

Fig. 52. (Right margin.) The lyrate organ and ovary of an adult female *Hæmogamasus hirsutus* (page 302) seen from above; $\times 85$. The camera spermatis is seen in the centre.

Fig. 53. The sacculus fœmineus, with its two great globular diverticula and other appendages, of an adult female of the same species, seen from above; $\times 75$. Drawn from a specimen dissected in very dilute glycerine. The sacculus, cornu, and both diverticula contain numerous spermatocysts; there are not any in the rami.

Fig. 54. (Left, near top.) Proximal termination of one of the tubuli annulati of an adult female of the same species in the corresponding ramus, seen from the side; $\times 400$. The globular termination within the ramus is extremely delicate and hyaline.

Fig. 55. (Lower left corner.) Distal termination, or mouth, of the same tube of the same creature, in the acetabulum of the leg, seen from the side, to show its form; $\times 250$.

- Fig. 56. (Centre of Plate.) The sacculus fœmineus, with its appendages and the lyrate organ, and ovary of an adult female *Laelaps oribatoides* (page 303), *in situ*, seen from above; $\times 175$. The two globular diverticula of the sacculus are seen, close together, above the lyrate organ. The long winding cornu sacculi contains all the spermatocysts, as is usual in oldish specimens; the eggs are attaining a considerable degree of development, but the most mature egg is not shown in the drawing. The ringed tubes are not shown quite to their distal extremities.
- Fig. 57. (Left margin.) Sacculus fœmineus and cornu and part of the ringed tubes of an adult female *Laelaps vacua* (page 303); $\times 300$. The cornu contains numerous spermatocysts, and some remains are seen in the sacculus; compare the former with figs. 27, 28. The tube between the sacculus and cornu is extended; *in situ* it is wound round the organ.
- Fig. 58. (Right lower margin.) Sacculus fœmineus, cornu sacculi, &c. of an adult female of a small *Laelaps*, sp. ? (page 303); $\times 150$. The spermatocysts are here seen both in the sacculus and cornu.
- Fig. 59. (Centre of upper margin.) Sacculus fœmineus, cornu, ringed tubes, lyrate organ, and camera spermatis of an old adult female *Laelaps acuta* (page 303), seen from above; $\times 100$. The tube from the cornu to the camera is extended by the sacculus and lyrate organ being moved further from each other than in nature. *In situ* the sacculus lies between and above the arms of the lyrate organ, and the tube winds round the former. The ringed tubes are cut short. The sacculus is turned a little on its side; *in situ*, its cornu is downward.
- Fig. 60. Sacculus fœmineus, cornu sacculi, ringed tubes, lyrate organ, and a few ova of an adult female *Holotaspis marginatus* (page 304), seen from above; $\times 160$. The short and singular cornu is between the sacculus and the lyrate organ.
- Fig. 61. (Centre of lower margin.) Sacculus and ringed tubes of an adult female of a small *Holotaspis*, sp. ? (page 304), seen from above; $\times 150$. There is not any cornu sacculi.
- Fig. 62. (Left margin.) Sacculus fœmineus, ringed tubes, and lyrate organ of an adult female *Laelaps cuneifer*, seen from above; $\times 150$.
- Fig. 63. (Right lower corner.) Sacculus fœmineus, ringed tubes, and lyrate organ of an adult female *Holotaspis montivagus* (page 304), seen from above. Organs *in situ*; $\times 150$.
- Fig. 64. The trefoil of three small sacs, &c., dissected out of the sacculus fœmineus of an adult female of the same species; $\times 300$.
- Fig. 65. (Near middle of left margin.) The lyrate organ, camera spermatis, ringed tubes, &c. of an adult female *Laelaps* (?) *ligoniformis* (page 305), seen from above; $\times 150$. There are not any sacculus fœmineus, cornu, &c.
- Fig. 66. Stalked receptacle (spermatheca ?) found close to the mouth of the ringed tube on each side of the body of the adult female of the same species, seen from the side; $\times 340$ (page 305). A small piece of the cuticle of the acetabulum of the third leg is seen on edge at the left of the drawing.

PLATE XXXV.

- Fig. 67. Horizontal, rather thick, section of an adult female *Hæmogamasus horridus*, almost at the level of the dorsal surface of the brain, where that organ comes nearest to the dorsum; $\times 75$. The sacculus fœmineus, with part of its cornu, and a portion of each ramus are seen in the centre of the figure; the rami are practically shortened (*i.e.* the whole length is not seen), because from their bending downward the section cuts the dorsal wall of each ramus at its bend; thus the apparent cæcal anterior ending of each ramus is not the termination of that organ, but the place where the section strikes the curve. The ringed tubes would lie below the section. The distal portion of one arm of the lyrate organ is seen on each side, crossed by the rami. The

most mature egg is seen in a form which it often assumes, sometimes to a much greater degree, when about being forced through a narrow passage—*e. g.* the entrance of the oviduct. A small part of the upper portion of the brain is seen with the dark cells of the cortical layer at the edge; behind it is the œsophagus, cut through and looking like a ring, surrounded by numerous portions of the tendinous platform (episternite) and the muscles proceeding from it. The cloaca is cut about where the two Malpighian vessels enter, and shows a septum between. The greater part of the portion of these vessels which lies in the hinder half of the body is shown, but not the bend by which they reach the cloaca. Almost the whole of the upper hind cæca of the ventriculus is shown, and nearly the whole of the right anterior cæcum; the left is cut across.

Fig. 68. Sagittal section of the posterior portion of an adult female *Hæmogamasus horridus*, cut a little to one side of the median line; $\times 75$. It shows the sacculus fœmineus, with a portion of its cornu and the commencement of one ramus lying above the ventriculus. Nearly the whole length of the oviduct and vagina is shown.

Fig. 69. Transverse section of an adult female of the same species, cut through the insertion of the fourth pair of legs; $\times 85$. It cuts the sacculus fœmineus just at the point where the rami are starting from it. The arms of the lyrate organ are cut across.

Fig. 70. Sagittal section of an adult female *Hæmogamasus hirsutus*, cut a trifle to the side of the median line; $\times 70$. It shows one of the great diverticula of the sacculus fœmineus cut through and also a part of the sacculus itself. The camera spermatis is full of filamentous spermatozoa, and the point where the arms of the lyrate organ join it is seen. Almost the whole length of the oviduct and vagina are exhibited.

Fig. 71. Sagittal section of an adult female *Holotaspis montivagus*(?), cut precisely in the median line; $\times 125$. The sacculus is shown with the central and a portion of one lateral chamber of the trefoil (pages 304–305); also the tube proceeding from the central chamber and wholly contained within the sacculus. The lyrate organ, cut through, is just behind the sacculus. An egg has the embryo highly developed; a portion of the Malpighian vessels of the embryo is already functional and is filled with the usual contents; it is near the dorsum of the embryo. This large egg has, as generally happens, flattened the cloaca and pushed some of the surrounding organs somewhat out of position. The whole length of the œsophagus is seen passing right through the brain and entering the ventriculus; one mandibular nerve is also indicated. A salivary, or poison, gland lies below the mandibular muscles. The tendinous platform (episternite) is seen almost surrounded by the ventriculus. The tendency to the dome-like recess in the vagina is seen.

Fig. 72. Transverse section of an adult male *Hæmogamasus horridus*, cut between the third and fourth legs; $\times 120$. The two great accessory glands are cut just where the large columnar cells of the hinder parts of the organs are fading into the more reticulated anterior portion; the testis is seen above in several blocks, showing sperm-mother-cells in various stages.

Fig. 73. Transverse section through the rostrum of an adult female *Hæmogamasus horridus*, cut near the anterior end of the pharynx; $\times 280$. It shows the tricarinate (in section triradiate) form of the pharyngeal skeleton (page 310) and the distensor and constrictor muscles.

Fig. 74. Transverse section of an adult female *Hæmogamasus hirsutus*, cut through the anus; $\times 200$. It shows the levatores and constrictor ani muscles, &c. The creature was killed with boiling water, and consequently the anus is protruded rather more than would be natural.

Ad Michael ad nat. del.

A.D. Michael ad nat. del.

A.D. Michael ad nat. del.

G. Freytag & Berndt, Vienna.