long continuance, powerful means are called for, to arrest the progress of high morbid action; this remedy, carried to such a length as to produce local effects upon the salivary glands, is generally entitled to our confidence.

That combination of tetanic symptoms, which is sometimes connected with hysteria, with a complex nervous affection denominated by a late writer,* 'Mymosis Urgens,' has been generally thought, from its comparatively mild character, to be a distinct disease from tetanus. But I am disposed to think, that the similar symptoms in each affection, arise from the same proximate cause, and constitute the same disease. The third case above related was unquestionably a case of traumatic tetanus. When the symptoms of this disease were nearly subdued by remedies, convulsions took place. Here the disease, from first to last, was probably the same, differing only in degree. At one end of the scale appears tetanus, and at the other, convulsion.

Beverly, Nov. 1823.

Case of Tetanus. By WARREN ABBOT, M. D.

[Communicated for the New-England Journal of Medicine and Surgery.]

NOVEMBER 18th, 1823, I was called to visit Mr. R—— a dispensary patient aged thirty-three. When I entered his room he was resting on a chair with his right arm and leg extended and firmly fixed by the violent contraction of the extensor muscles, his mouth, was forcibly drawn to the right side, and appeared to be held in doubtful contest by the spastic action of the levator and depressor anguli oris and other muscles of the mouth and lips, while his whole countenance exhibited a forbidding and almost terrific appearance. I asked the cause of his suffering but he could not answer so long as the paroxysm continued. I enquired of his wife, she replied, 'I dont know what is the matter with him; he appears dreadful silly, snappish and crazy of late; sometimes he wants drink and when I bring it to him he knocks it away; sometimes he wont speak, and then he makes up such awful looking faces; then again he wants something done to his hand where it was hurt, but if I go to touch him he screeches out like a mad man; he is all the time getting worse, he wont eat, nor drink, what shall I do with him?' On farther inquiry it appeared that a few days previous a rusty nail had by accident been thrust into the right hand near the lower end of

* Hall,

the metacarpal bone of the little finger. The wound had so far healed that nothing appeared externally but a small cicatrix. The precise nature and extent of the wound therefore could not easily be ascertained. Neither could I learn the direction the nail penetrated, or the relative state of the skin and loose cellular integument at the time the wound was inflicted, it is probable however that one of the tendons of the little finger was injured. Shortly after the accident when the wound began to heal, he had, as he expressed it, a 'strange kind of feeling beginning in his little finger, and running up his arm to the shoulder, then down his side to his leg, at last it went clear through to his stomach and insides he could'n't tell how. It was sometimes so bad he could neither talk, eat nor drink—he did not dare to drink for fear he should be strangled.'

On examination the large muscles on the right side, particularly the latissimus dorsi, seemed contracted into a form like the segment of a globe. The spasmodic affection was most severe on the right side, but the whole system was becoming more subject to tetanic influence, the paroxysms were also occurring more frequently. As there could be no doubt as to the character of the disease, and it being necessary that something should be done without delay, I thought that opium was a remedy best suited to his case. He could take no liquid, he was therefore directed to take two grains of opium in pills immediately to be followed by one grain every hour till the spasmodic action should be overcome.

19th—In a free perspiration, had taken twelve grains of opium in ten hours; after an hour or two of sleep appeared perfectly easy, and was able to wipe the sweat from his face with his *right hand*. A cathartic of salis and senna was directed to be taken in the course of the day.

20th—The cathartic had operated well. The patient was sitting at the table with his family at breakfast as usual.

I have thought proper to communicate this case, not on account of any thing new, but because every fact in clinical medicine, relating to such an alarming disease as tetanus, is worth preserving and retaining till indelibly fixed on the mind of every one who has any thing to do with life and health.

1. It is worthy of remark that owing to the constriction of the muscles of the throat, the symptoms of hydrophobia were distinctly manifest in this case.

2. The wound, which was the exciting cause, was completely healed, when the nervous irritation and spastic action was the greatest. There is an instance related of a man who died of

this disease, after amputation of the shoulder joint, in which case, the life of the patient and the perfect healing of the wound were terminated on the same day.

3. In giving opium in diseases of this kind, we must regard the effect produced, rather than the quantity given.*

Boston, Dec. 5th, 1823.

A Case of Organic Disease of the uterus; and a Case of Imperforated Hymen. By C. G. ADAMS, M. D.

[Communicated for the New-England Journal in a letter to one of the Editors.]

ABOUT ten years since, when a pupil, I recollected to have dissected a human uterus with its appendages, which had, for two or three years, been kept in alcohol. In the almost shapeless mass of disease which the preparation presented, several tumours varying in size, the largest being about that of a man's fist, and of fleshy consistence, were found in the parietes of the uterus and involved in the broad ligament of one side. A tumour, apparently buried in the thickened parietes of the uterus at its fundus, when disengaged from the cyst which enveloped it, was found to be very firm bone. Its shape was circular, flattened on one side and convex on the opposite—its surface had a rough granulated appearance. On passing a saw through it, to the action of which it afforded very strong resistance, it appeared to be constituted of many very compact osseous portions irregularly blended together, and defined by small interstices, the structure of the bone itself being in no degree cellular. I do not recollect any remarkable appearance of the ovaria. There was no tumour on the internal surface of the uterus. The bone, which I send you, weighs in its present dry state 2½ ounces, about one fourth part of it having been removed by the saw, and not in my possession.

I am informed by my father (Daniel Adams, M. D.) in whose practice the case occurred, that the subject of it was a married black woman of about forty-five. She never was pregnant. Painful and irregular menstruation—pain and weight in the uterine region together with distressing tympanites had for eight or ten years back been increasing in severity and constancy.

* "Twenty grains of opium have been given every three hours, in conjunction with musk, cinnamon and other medicines; and continued with little abatement for seventeen days; in which time the patient took five hundred grains of this narcotic." *Good's Study of Medicine Vol. iii. page 226.*