of respiration. After a time they should be recommenced, and the process continued until the urine becomes alkaline.

9. Subcutaneous injections of carbonate of sodium are not to be commended, on account of the pain and the deep-seated inflammation they produce.

THE NIGHT-SWEATS OF PHTHISIS TREATED WITH CAMPHORIC ACID.

LEU (*Charité Annalen*, 1889) has employed camphoric acid—produced by the oxidation of camphor with sulphuric acid—against the night-sweats of phthisis, in 13 patients; the drug being administered 155 times in all. In 60 per cent. complete success was obtained, this consisting in complete dryness of the skin. There were no good results obtained in 18 per cent., and partial success in 22 per cent. The dose employed varied from 30 grains given in the evening, to 75 grains administered in two doses. The good effects were often not observed until the second night after the drug was given; and the action of the medicine often then persisted during several nights.

The author found the value of camphoric acid greater than that of nupropine, and the effects more lasting. The drug was also free from disagreeable secondary effects, except that in one case an attack of urticaria developed, which, however, had probably nothing to do with the medicament. He tried its alcoholic solution externally for bathing the body, and found it decidedly superior to vinegar in preventing local sweating.

SURGERY.

UNDER THE CHARGE OF

J. WILLIAM WHITE, M.D.,

PROFESSOR OF CLINICAL SURGERY IN THE UNIVERSITY OF PENNSYLVANIA; SURGEON TO THE UNIVERSITY
PHILADELPHIA, AND GERMAN HOSPITALS.

THE TREATMENT OF CICATRICIAL AND CANCEROUS ŒSOPHAGEAL STRICTURES.

LE FORT (*Bulletin générale de Thérapeutique*, January 15, 1890) divides the contractions of the œsophagus which may require treatment into two great classes—the cicatricial and the carcinomatous—and arrives at the following conclusions:

1. As to the cicatricial strictures: He believes gastrostomy to be a dangerous remedy, which should only be employed in the last extremity, by which he does not mean that we should wait until the patient is dying from inanition, but that we should at first employ all other means, especially dilatation. It is necessary, in his opinion, to use sounds of a special form (which he figures), passing, in the first place, the very finest bougies, which enter

freely into the stricture, and which sufficiently dilate it to enable the patient to swallow liquids. After this all urgency is passed, and we can take the time required to accomplish a complete cure, and without risking the life of the patient.

2. As to gastrostomy in cancerous contractions of the œsophagus, the mortality is enormous (72 in 100 at least) and the average life of those who survive the operation is hardly more than two months, or three months at the most. It seems, therefore, that even in carcinomatous cases continuous dilatation should be attempted. Gastrostomy is only justified by the impossibility of introducing nourishing drinks, and in those patients who, having retained their strength and reason, suffer from the fear of death by starvation. In cicatricial contractions dilatation in the manner which has been indicated should be tried with all the more energy and perseverance, because even after gastrostomy it is necessary to return to the use of the sound in order to dilate the œsophagus. In either class of cases Professor Le Fort refuses to consider gastrostomy as the routine treatment of contractions of the œsophagus.

THE SURGICAL TREATMENT OF HEPATIC ABSCESS.

MR. RICKMAN J. GODLEE summarizes (*The British Medical Journal*, January 11, 18, 25, 1890) as follows the essential part of his views on the treatment of hepatic abscess, recently published in some extremely valuable lectures which embrace the whole subject, and embody the latest and most authoritative opinions as to its surgical aspect:

1. Pyæmic abscesses do not call for surgical interference, or, if in rare cases one should point, it is only opened to relieve symptoms, but without hope of doing permanent good.

2. The same observations apply to abscesses resulting from suppurative phlebitis of the portal vein.

3. Multiple abscesses associated with dysentery or ulceration of the bowels are very unfavorable for surgical treatment. They must, however, be opened and treated on the same lines as the single or tropical abscesses, because they cannot be certainly diagnosed.

4. Single abscess of the liver, whether tropical or not, must, if it approach the surface, be opened, the following precautions being adopted:

(a) If it present at the epigastrium, the presence of adhesions must be ascertained before incising the liver.

(b) If through the chest-wall, a spot must be chosen below the normal limit of the pleura; but if by chance either pleura or peritoneum be opened, the opening must be closed with a double row of stitches before incising the liver.

(c) Strict antiseptic precautions must be throughout adopted, either carbolic acid or some slightly soluble salt of mercury being employed for the dressing.

(d) The tube must be of large size at first, and a tube of some sort must be kept in until the discharge is reduced to a very minute quantity.

If the abscess have burst into the lung, pleura, pericardium, peritoneum, or kidney, and the position of the abscess can be clearly determined, it must

be opened without delay. If the position of an abscess be only suspected and the patient be losing ground, it is right to puncture the liver in the most likely situation, bearing in mind that, although usually quite harmless, a slight amount of risk accompanies this very trivial operation.

This rule applies to cases in which the abscess has ruptured into any of the cavities enumerated above. If, on the other hand, whether the abscess has ruptured or not, there are no means of diagnosing the whereabouts of the matter, and the patient is not losing or is even gaining ground, the surgeon should hold his hand for a time.

5. Hydatids of the upper and back part of the liver are to be treated upon the same lines; but in cases of this sort, and in those of subdiaphragmatic abscess, it must be remembered that the diaphragm may be pushed up to a very great height, thus closely simulating intrapleural suppuration.

6. Empyema, pericarditis, and peritonitis caused by rupture of an hepatic abscess or hydatid must be promptly dealt with on general principles.

IMMEDIATE SUTURE OF THE KIDNEY AFTER NEPHRO-LITHOTOMY.

M. LE DENTU reports a case in which he removed a renal calculus from a patient who had suffered for fifteen years—having concluded that the calculus was seated in the pelvis of the kidney, although there were no symptoms of suppurative pyelo-nephritis. Some time before, he had removed in the same way a renal calculus which did not present any trace of suppuration, and attempted to obtain primary union of the renal substance, but without success. Nevertheless, he repeated the attempt in the second case. After cutting the inferior half of the convex edge and extracting the calculus, he put in seven stitches of catgut. The renal tissue was so friable that it was not possible to draw these threads very tight, but those that were put in were sufficient to arrest capillary hemorrhage. The result was excellent: not a drop of urine escaped from the wound. By way of precaution, a drain was left in until the fifteenth day. In spite of this success, he does not believe, however, that primary union is easy to obtain after nephro-lithotomy.

SURGICAL TREATMENT OF TUBERCULAR PERITONITIS.

CECCHENELLI reports (*Centralblatt für Chirurgie*, No. 39, 1889) four cases of tubercular peritonitis treated with incision and irrigation. He divides such cases into two groups:

1. A dry inflammatory process, in which surgical interference is useless, since it is impossible to separate all adhesions; and, therefore, the whole peritoneal cavity cannot be cleansed.

2. Peritonitis with ascites, in which incision is indicated, especially in sacculated peritonitis.

In free effusion, paracentesis, followed by moderate irrigation, is often sufficient.

THE SURGICAL TREATMENT OF PERITYPHLITIS.

At the last French Surgical Congress (*Archives Générales de Médecine*, Nov. 1889), M. ROUX, of Lausanne, stated his opinion that there is too great a

tendency to abstain from all active treatment in suppurative perityphlitis, from the incorrect belief that spontaneous cure is the rule. There are, on the contrary, a great number of accidents and relapses, without speaking of the frequent cases of fatal peritonitis, which can be avoided by surgical interference at the proper time. In a series of twenty-one perityphlitic abscesses treated medically, M. Roux has had six imperfect cures, with fistulas and repeated abscesses, and nine cases of peritonitis of which eight terminated fatally. He does not claim that it is necessary to operate in all cases of perityphlitis, but that it is better not to continue too long the expectant treatment. As soon as there is pus, it is well to evacuate it. One can be assured that the pericæcal inflammation is suppurative, when by palpation and percussion an evident infiltration of the superior part of the cæcum and of the ascending colon is found, the intestine being empty. Abscess once diagnosed, M. Roux does not advise, in order to discover the focus of suppuration, the exploratory incision of Sonnenburg, nor exploratory punctures, which frequently give negative results, and are not without danger. M. Roux makes a very large incision, similar to that for the ligature of the external iliac, separates the fascia transversalis, and carries his finger behind the cæcum, where ordinarily the pus is collected. Even if an error in diagnosis is made, such error is only possible in cases which practically demand the same treatment as perityphlitis, and no additional risk is run. If the purulent focus is not found, it will empty itself spontaneously, more easily toward the exterior than into the cavity of the peritoneum, owing to the presence of this wound.

DISEASE OF THE VERMIFORM APPENDIX.

DR. CHARLES MCBURNEY (*The New York Medical Journal*, December 21, 1889), in a valuable paper on the above subject, reviews the general clinical history of cases of appendicitis, and lays especial stress on the important aid to diagnosis to be derived from a symptom which he has frequently observed, namely, the ascertaining, by the pressure of a single finger-tip, that the point of greatest tenderness is, in the average adult, almost exactly two inches from the anterior iliac spine, on a line drawn from this process through the umbilicus. Much greater tenderness at this point than at others, taken in connection with the history of the case and other well-known signs, he looks upon as almost pathognomonic of appendicitis. This point indicates the situation of the base of the appendix, where it arises from the cæcum, but does not by any means demonstrate, as one might conclude, that the chief point of disease is there. The abscess, or concretion, or cyst, may be at quite a little distance, but the greatest pain, on pressure with one finger, will be felt at the point described. The incision should be a liberal one, for much room may be required, and a five-inch cut in the adult is not too much. It should follow as nearly as possible the right edge of the rectus muscle, and the centre of the incision should lie opposite to or a little below the anterior iliac spine, on a line drawn to the umbilicus. When the external oblique aponeurosis is cut through by this incision, the aponeurotic structure, in which the other abdominal muscles end, comes into view, and is easily divided without cutting muscular fibre. Then the fascia transversalis, the subperitoneal fat, and the peritoneum are cut in succession. If pus has

formed close against the anterior abdominal wall, these last-mentioned will be found infiltrated with serum, and even thickened so as to look like cheesy tubercle. Otherwise these parts may appear perfectly normal. On opening the peritoneum the appendix may at once be seen, or adhesions and inflammatory exudations may have so distorted the parts that a careful and difficult search may be required to find the appendix at all. It may be flattened out and glued firmly to the inflamed surface of the cæcum by old and recent adhesions, or it may be coiled upon itself and hurried out of view in a mass of lymph. The finger is often quicker than the eye to detect the appendix in these conditions, as it is very certain to be found where the greatest thickening, as felt by the finger, exists.

OSTEOPLASTIC EXARTICULATION OF THE FOOT.

DR. W. RASUMOWSKY (*Archiv für Klinische Chirurgie*, vol. xxxix., 1889) discusses the various operations on the foot (Pirogoff, Wladimiroff, Mikulicz, Tauher, Malgaigne), and reports a modification of his own, which he has employed in one case, and which he thinks justifies the following conclusions:

1. In children it is possible to perform an osteoplastic foot operation without the removal of the lower epiphysis of the tibia—that is to say, amputation of the leg can often be replaced by osteoplastic exarticulation of the foot, a procedure which possesses unmistakable advantages.

2. When the cutaneous covering of the heel is intact—i. e., in cases which in adults would indicate Pirogoff's operation, in children, according to the method of Rasumowsky, the posterior portion of the os calcis may be thrust into the inter-malleolar niche. Similar modifications may be made according to the region and amount of uninjured skin which remains. The details are described with great fulness.

TREATMENT OF ANEURISMS OF THE EXTREMITIES.

The discussion at the French Surgical Congress on the question of operative interference in cases of aneurism embraced certain facts brought forward by M. KIRMISSON (*Archives Générales de Médecine*, November, 1889). Of three spontaneous aneurisms related by this surgeon, one only was cured by digital compression after a painful attempt to use an elastic bandage; in the two other cases, compression having failed, it was necessary to have recourse to the ligature.

M. VASLIN had had the same experience, and advocated in aneurisms of the inferior extremity the application of an aseptic ligature at a distance from the tumor.

M. TRÉLAT, who employs only the operative method, declared himself in favor of the direct extirpation of the sac, rejecting the aseptic ligature upheld by Lister, Boeckel, Lucas-Championnière, and Verneuil. The advantage of cutting out the sac is to avoid subsequent accidents—that is to say, inflammation of the sac, producing circulatory and nervous disorders. Arterio-venous aneurisms must be treated at the earliest possible moment; if the sac is small, ligate the arteries and veins; if the sac is bulky, extirpate it.

M. RECLUS rejected all of the milder measures, not alone because cure is only obtained in about one-half the cases, but also because these procedures

are extremely painful, and, above all, because operative methods applied later are liable to be followed by grave accidents, which are due to the irritation excited by the earlier attempts.

M. VERNEUIL, who was formerly an advocate of the "mild" treatment, avoids surgical interference in old people in cases of aneurism of the trunk. With this exception, he adopts the operative method, preferring the ligature to extirpation, often a most difficult operation.

M. PEYROT reported an interesting case of popliteal aneurism, the removal of which was very long and difficult. He believes that while there are cases suitable for ligature, and others which indicate extirpation, that the latter process will become the chosen one.

M. PÉAN (*Revue de Chirurgie*, November, 1889) calls attention to the fact that in nearly all aneurisms of the extremities operation is clearly indicated, as they are very accessible and differ only according to the importance of the artery involved, popliteal and femoral aneurisms being the most interesting surgically. Their cure can be accomplished not only by arresting the blood current by ligature, but also, and preferably, according to M. Péan, by occlusion of the artery by means of forceps. If the aneurism is small and circumscribed, one need not hesitate to open the sac and to empty it, compressing both the superior and the inferior ends of the artery. The forceps are allowed to remain for forty-eight hours, after which they are removed and the wound is closed. For the femoro-popliteal aneurisms the same method could be used, but experience has demonstrated that immediate union is difficult to obtain, and that secondary hemorrhage may result. In these conditions it is preferable to look for the artery above and below the sac, and to place forceps at each of these two points. Pressure has a great advantage over the ligature, in not necessitating the denudation of the arteries and in not permitting foreign bodies to be an obstacle to union by first intention.

NERVE SUTURES.

E. ETZOLD records (*Deutsche Zeitschrift für Chirurgie*, Bd. 29, Hefte 5 und 6, 1889) a number of cases occurring at the Dorpat clinic, in which various nerves, chiefly the ulnar, radial, median, and musculo-cutaneous, were sutured at different intervals after their division, and on the whole with great success. After considering the general subject, he arrives at the following conclusions:

Nerves do not unite by either primary adhesion or second intention. The axis-cylinders are the extension of the cells of the ganglia, and their re-formation or their union by means of an exudation of cellular elements of mesodermal origin is, for anatomical reasons, not to be expected.

Divided nerves are regenerated by means of a proliferation from the proximal stump. This was established by experiments upon animals, and has been confirmed by clinical observation, which shows beyond all doubt that the proximal end of a divided nerve is regenerated earlier and more completely than the distal end.

The return of sensibility is of no value in the diagnosis of nerve regeneration.

The symptoms indicating its occurrence are, *a*, native muscle contraction; *b*, disappearance of atrophy, especially of muscular atrophies; *c*, the slow appearance of this improvement; *d*, the return of faradic excitability in

muscles previously paralyzed. The galvanic current is not of much importance in the diagnosis of nerve regeneration.

Spontaneous union of divided nerves in the extremities is very rare. In high injuries of nerves the prognosis is unfavorable in spite of nerve sutures.

Regeneration of nerves is prevented by the extensive formation of cicatricial tissue.

Nerve suturing is not only a justifiable operation, but in every traumatic case of nerve section it is the duty of the surgeon to adopt it.

The essentials of success are: absolute antisepsis, complete hæmostasis, avoidance of irritation. If, after nerve injuries, a congested condition of the limb results, it should be elevated, and massage employed as soon as the wound is healed. Direct galvanization of the nerve scar should be employed, as well as massage soon after cicatrization in order to diminish the scar.

It is not proven that electric treatment of the organs supplied by the cut nerves either limits the atrophy or favors the nerve regeneration.

Massage and passive gymnastics constitute the rational treatment for peripheral paryses.

The most extensive use of the extremity that is found possible after nerve section, appears to have a favorable influence upon the healing.

ACUTE MERCURIAL POISONING.

KAPOSI (*Centralblatt für Chirurgie*, December 18, 1889) reports the case of a vigorous woman forty years old, who on account of a maculo-papular syphiloderm had taken, by hypodermic injection within seven weeks, a considerable quantity of "oleum cinereum," which resulted in the appearance of an acute stomatitis, followed by dysentery and death six weeks later.

The autopsy showed the well-known signs of mercurial poisoning. The chemical analysis of the tissues of the last place of injection showed that scarcely a third of the amount deposited there had been absorbed. This fact made the case still more noteworthy, as usually gray oil is exceedingly well tolerated.

In four other cases Kaposi found a greater or less degree of acute stomatitis after similar injections, and in one case there was coexistent albuminuria.

OTOLOGY.

UNDER THE CHARGE OF

CHARLES H. BURNETT, M.D.,

AURAL SURGEON, PRESBYTERIAN HOSPITAL, ETC., PHILADELPHIA.

CREOLIN IN AURAL THERAPEUTICS.

In the report of the work in the ear clinic at Göttingen, BÜRKNER communicates the experience with creolin (*Archiv für Ohrenheilkunde*, vol. xxviii., Part 4, October, 1889). Preference has been given to Pearson's creolin, which